

)

١

U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460

EPA Reg. Number:

Date of Issuance:

9198-148

SEP 23 1994

NOTICE OF PESTICIDE:

<u>x</u> Registration _ Reregistration

(under FIFRA, as amended)

Term of Issuance: Until Reregistration

Name of Pesticide Product:

The Anderson's Pest Arrest 10% Dust

Name and Address of Registrant (include 21P Code):

The Andersons D/B/A Free Flow Fertilizer P.C. Box 119 Maumee, Ohio 43537



Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration/ reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
- Add the phrase "EPA Registration No. 9198-148" to your label before you release the product for shipment.

Signature of Approving Official:

Date:

SEP 23 1994

Dennie H. Edward fr

Dennis H. Edwards, Jr.

Product Manager 19

Insecticide-Rodenticide Branch

Registration Division

EPA Form 85/0-6

- 3. Change the Statement of Practical Treatment for oral exposure to read something like this: "If swallowed: Never give anything by mouth to an unconscious person. If conscious and not convulsing, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger." If you do not want to change your currently proposed statement, please submit a rationale for not wanting to do so.
- 4. The format for the whole Statement of Practical Treatment should be changed, with separate subheading for each exposure route. You may also add the paragraph titled GENERAL from your other carbaryl labels such as Pest Arrest Flea and Tick Killer (EPA Reg. No. 9198-142) to this label.
- 5. On page 5, under Imported Fire Ant control, change "8 level teaspoonfuls" to "12 level teaspoonfuls." Two ounces of powder equals 12 teaspoons.
- 6. Submit five copies of the revised labeling for this product before you release this product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

Dennis H. Edwards, Jr.

Product Manager 19

Insecticide-RodenticideBranch Registration Division (7505C)

Dennis H. Edwards

Enclosure

)

)

1

THE ANDERSONS PEST ARREST 10% DUST

> EPA REG. NO. 9198-(TO BE ASSIGNED) EPA EST. NO. 9198-OH-1

KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel For Additional Precautionary Statements

ACCEPTED with COMMENTS in EPA Letter Dated		
/ SEP 2 3 1994	NET WEIGHT	LBS.
Under the Pederal Insecticide, Fungicide, and Redensicide Act as amended, for the proticide registered under EPA Reg. No.	TREATS UP TO	SQ. FT.

)

)

)

(BACK PANEL)

NET WEIGHT LBS.

TREATS UP TO SQ. FT.

THE ANDERSONS

PEST ARREST 10% DUST

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

HARMFUL IF SWALLOWED. Causes eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly after handling. Avoid contamination of food, feeding troughs and water receptacles.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, give water or milk to drink and telephone for medical advice. Consult medical personnel before inducing vomiting. If medical advice cannot be obtained, then take person and PRODUCT CONTAINER to the nearest medical emergency treatment center or hospital.

In case of eye contact wash eyes immediately with fresh water for at least 15 minutes. Get medical attention if irritation persists. For skin, wash with plenty of soap and water. If inhaled, remove person from exposure area.

NOTE TO PHYSICIAN:

Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning equipment or by disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

VEGETABLES AND FRUIT

Pest Arrest 10% Dust can be applied with any suitable hand or power duster. For best results, make application in the morning or evening when the air is still. Treat both upper and lower surfaces of leaves. Applications should be started when insects first appear and repeated at 7 to 10 day intervals, or 5 to 7 days in rainy weather. Apply to deposit a light visible dusting to all plant and fruit parts or 4 to 8 ounces per 1,000 sq. ft. which is equal to approximately 10 to 20 lbs./acre. Use the low rate when insects are first seen or are anticipated; the high rate when insects are present in damaging numbers.

Crops with the pests are listed below. Observe number of days to stop treatment before harvest.

Asparagus:

Asparagus Beetles - 1 day before harvest. Do not apply more often than once every 3 days.

Beans:

)

)

Mexican Bean Beetle, Bean Leaf Beetle, Leafhoppers, Flea Beetles, Corn Earworms, Lygus Bugs, Stink Bugs, Lime Bean Pod Borer, Japanese Beetle - Can be applied day of harvest.

Beets (Table):

Armyworms, Flea Beetles, Leafhoppers - 3 days. 14 days if tops are used for food or feed.

Brussels Sprouts, Broccoli, Cabbage & Cauliflower:

Small Cabbage Looper (treat every 5 to 7 days for control), Armyworms, Flea Beetles -

3 days before harvest.

Carrots:

Leafhoppers, Flea Beetles - Can be applied day of harvest.

Citrus:

Black Scale, California Red Scale, Soft Brown Scale, Orange Tortrix, California Orange Dog, Fruit Tree Leaf Rollers - For Scale Insects best results are obtained when crawlers are present. 5 days before harvest.

Collards, Kale, Mustard Greens, Radishes, Rutabagas, Chinese Cabbage, Horseradish & Turnips:

Flea Beetles, Parlequin Bugs, Leafhoppers, Armyworms, Imported Cabbageworms, Tarnished Plant Bugs - 3 days of harvest on Horseradish, Radishes, Rutabegas, and Turnip Roots. 14 days on Chinese Cabbage, Collards, Kale, Mustard Greens and Turnip Tops.

Corn:

)

)

)

(Sweet and Field) Corn Earworm, Flea Beetles, Leafhoppers, Fall Armyworms - For Corn Earworm control treat corn stalks every 3 days until silks dry. Can apply day of harvest. Wait 7 days before foraging.

Cucumbers, Melons, Squash & Pumpkins:

Squash Bugs, Leafhoppers, Cucumber Beetles, Flea Beetles - Do not apply to Melons or Cucumbers when temperatures exceed 80° F or when vines are wet. Can be applied day of harvest.

Eggplant:

Corn Earworm, Flea Beetles, Hornworms, Cucumber Beetles - Can be applied day of harvest.

Grapes:

Grape Leafhoppers, Grape Leaf Folder, Grape Berry Moth, European Fruit Lecanium, Grape Leaf Skeletonizer, Japanese Beetles, June Beetles, Red Banded Leaf Roller - Day of Harvest.

Head and Leaf Leatuce:

Armyworms, Small Cabbage Looper (treat every 5 to 7 days for control), Leafhoppers, Salt Marsh Caterpillers - 3 days before harvest on Head Lettuce; 14 days before harvest on Leaf Lettuce.

Okra:

Corn Earworm, Flea Beetle, Japanese Beetle, Stink Bugs - May be applied day of harvest.

Peanuts and Cowpeas:

Blister Beetles, Mexican Bean Beetles, Thrips, Leafhoppers, Three Cornered Alfal'a Hopper, Cucumber Beetles, Bean Leaf Beetles, Japanese Beetles, Alfalfa Caterpillar, Velvetbean Caterpillar, Green Cloverworm, Armyworms, Webwarms, Corn Earworms, Stink Bugs, Grasshoppers - On Cowpeas do not apply within 3 days of harvest; on peanuts, no limitation.

Peppers:

(White) Flea Beetles, Leafhoppers - Day of harvest.

Tomatoes:

Tomato Fruit Worm, Tomato Hornworm, Flea Beetles, Stink Bug, Leafhoppers - Apply day of harvest.

Strawberries:

Leaf Rollers, Lygus Bugs, Salt Marsh Caterpillar - One day before harvest.

Sugar Beets:

)

)

Leafhoppers, Armyworms, Webworms, Grasshoppers - 14 day pre-harvest interval.

ORNAMENTALS, FLOWERS, SHRUBS AND TREES

Apply this product to thoroughly cover the plant foliage for control of the following pests. Repeat as needed for continued control.

Apple Aphid
Blister Beetle
Boxelder Bug
Elm Leaf Beetle
Flea Beetle
June Beetle
Leafhoppers
Leaf Rollers
Mealy Bugs
Mimosa Webworm
Oak Leaf Miner
Orange Tortrix

Periodical Cicadae
Psyllids
Rose Aphids
Rose Slug
-SCALESoft Brown
European Elm
Oak Kermes
Oystershell
Black Cottony

Maple Thrips (Exposed)

For best scale insect control, apply when newly hatched "crawlers" are present. Do not apply to Boston Ivy as plant injury will result.

TURF AND PERIMETER INSECTS

Apply 4 oz. to cover 250 sq. ft. of smaller turf areas adjacent to patios and home recreational areas or foundation perimeter areas to control Ants, Chiggers, Earwigs, Fleas, Millipedes, Spiders, Sod Webworms, Sowbugs and Leafhoppers. Area should be lightly sprinkled after application to wash dust off foliage. Repeat as necessary. Keep children and pets off treated areas until grass has been sprinkled and area is dry.

IMPORTED FIRE ANT CONTROL

12

Apply 2 oz. (8 level teaspoonfuls) evenly over and around each mound. After application, gently sprinkle the treated area with a minimum of 1 quart of water per 6 inches of mound diameter. For best results, do not disturb the mound before or during the application. Apply in cool weather, 65°-80°, or in early morning or late evening. Treat new mounds as they appear.

NON-COMMERCIAL POULTRY (Chickens, Partridges, Pheasants, Pigeons): Northern Fowl Mite, Lice and as a supplement to premise treatments for Chicken Mite and Fleas

8 ounces per 100 birds. Apply thoroughly by means of a shaker can or hand duster. Repeat in 4 weeks if necessary. Do not use on Poultry or game birds within 7 days of slaughter.

IN OR AROUND POULTRY HOUSES: Chicken Mite, Fleas and Bed Bugs

8 ounces per 40 sq. ft. of litter or floor space and to roosts and adjacent areas by means of a shaker can, hand duster or power duster. Force dust into cracks and crevices. Repeat as necessary. Do not treat nest litter. Avoid direct contamination of eggs. Do not use in premise treatment within 7 days of slaughter.

DOGS AND CATS: Ticks and Fleas

•

)

)

Dust lightly over the animal and rub thoroughly into the skin. Apply also to the feet between the toes and on the legs. Apply at weekly intervals if needed. Avoid treatment of kittens and puppies under 4 weeks of age.

NOTE: Do not allow treated pets back on carpets and rugs in the house as this product may stain or discolor rugs and wall-to-wall carpets. Dust lightly in and around kennels and dog houses by means of a shaker can, hand or power duster. Force dust into cracks and crevices. Repeat as necessary, but not more often than once weekly. DO NOT USE ON PREGNANT DOGS.

DO NOT USE IN THE HOME: DO NOT APPLY TO RUGS AND CARPETS AS STAINING MAY OCCUR.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area.

DISPOSAL: PRODUCT - Partially filled bag may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash.

CONTAINER - Do not reuse bag. Discard bag in trash.

WARRANTY STATEMENT

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions For Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

)

Manufactured by:

THE ANDERSONS LAWN FERTILIZER DIVISION P.O. BOX 119 MAUMEE, OHIO 43537