919	8-158		1-11	12002		030 000	
SEPA	Environmenta	Inited States		Form Appro	Registra Amendn	tion	O. Approval expires 2-28-99 OPP Identifier Number
		Application	on for Pestic	ide - Sectio	n I		
1. Company/Product Number 9198-158			2. EPA Product Manager 3. Pro LaRocca			oposed Classification	
4. Company/Product (Name) The Andersons Fertilizer Plus 0.058% Talstar			РМ# 13	PM#			
5. Name and Address of Applicant (Include ZIP Code) The Andersons Lawn Fert. Div., Inc. d/b/a Free F Fert. PO Box 119 Maumee, OH 43537 Check if this is a new address			6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. JAN 1 1 2002 Product Name				
<u> </u>			Section -				
Amendment - Explain below. Resubmission in response to Agency letter dated Notification - Explain below.			Final printed labels in repsonse to Agency letter dated "Me Too" Application. Other - Explain below.				
This notification is consistent we confidentical statement of form notification is not consistent will penalties under sections 12 an	ula of this product. I under th the terms of PR Notice 9	thad that it is a vi	iolation of 18 U.S.C. Se	ec. 1001 to wilfully in nay be in violation of	ake any false stater	nent to EPA	A. I further undersant that if this
1. Material This Product Wil	T	<u> </u>			T		
Child-Resistant Packeging Yes No * Certification must be submitted	Unit Packaging Yes No If "Yes" Unit Packaging wgt. container		Water Soluble Packaging Yes No If "Yes" No. per Package wgt container		2. Type of C	Metal Plastic Glass Paper Other (S	pecify)
3. Location of Net Contents	Information	4. Size(s) Re	tail Container	5.	Location of Labe On Label	l Directio	ns
6. Manner in Which Lebel is	Affixed to Product	Lithog Paper Steno	raph glued iled	Other _			
			Section - I	V			
1. Contact Point Complete	items directly below t	or identification	n of individual to b	e contected, if n	ecessary, to pro	cess this	application.)
Name Dawn L. Walters			II			F elephone 334-262-2	No. (Include Area Code) 766
I certify that the state I acknowledge that an both under applicable	y knowlinglly false or		all attachments th				6. Date / pilication Received (Stamped)
2. Signatura Daun G. Waltura			3. Title Regulatory Administrator			4 * * C C	
4. Typed Name Dawn L. Walters			5. Date 11/20/01			**************************************	



The Andersons Fertilizer Plus 0.058% Talstar®

The Committee of the property of the committee of the com

This product is a unique combination of insecticide and fertilizer to control damaging insects while replenishing your lawn with essential nutrients. It is designed to control a variety of insects on lawns.

Only for use, and storage by commercial applicators for use to control ants (including imported fire ants), mole crickets and other insect pest on lawns.

Active Ingredients Bifenthrin: * Other Ingredients	<u>99.942%</u>
*Cis isomers 97% minimum, trans isomers 3% maximum US Patent No. 4,238,505	

Talstar-Registered Trademark of FMC Corporation

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Net Weight 50 lbs. (22.68 kg) Covers Up To 25,000 Sq. Ft.

EPA Reg. No. 9198-158
EPA Est. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A
Underlined letter is first letter used in run code on bag





The Andersons Fertilizer Plus 0.058% Talstar

Net Weight 50 lbs. (22.68 kg) Covers 12,500 Sq. Ft.

Guaranteed Analysis		
Total Nitrogen (N)	XX %	
Available Phosphate (P ₂ 0 ₅)		
Soluble Potash (K ₂ 0) XX%		
Plant nutrients derived from:		
For products being sold into Washington state:		
Information regarding the contents and levels of metals in this product is available on the internet		
at http://www.regulatory-info-ap.com./		

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or when disposing of equipment washwaters.

FIRST AID	
If swallowed	Call a poison control center or doctor for treatment advice
	Have person sip a glass of water if able to swallow
	Do not induce vomiting unless told to by a poison control center or doctor
	Do not give anything by mouth to an unconscious person.
lf on skin	Take off contaminated clothing
	Rinse skin immediately with plenty of water for 15-20 minutes
	Call a poison control center or doctor for treatment advice
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	Remove contacts, if present, after the first 5 minutes, then continue rinsing
	Call a poison control center or doctor for treatment advice
If inhaled	Move person to fresh air
	If person is not breathing, call 911 or an ambulance, then give artificial person is not breathing and the person is not been give artificial. The person is not breathing and the person is not been give artificial. The person is not breathing and the person is not breathing and the person is not breathing and the person is not breathing. The person is not breathing and the person is not breathing and the person is not breathing and the person is not breathing. The person is not breathing and the
	respiration, preferably mouth-to mouth if possible
	Call a poison control center or doctor for treatment advice
	uct container or label with you when calling a poison control center or doctor or 💥
	ment. You may also contact 1-800-757-8951 for emergency medical treatment
information.	₱ ₱ ₱ ₱ ₱ ₱ ₱ ₱ ₱ ₱ ₱ ₱ ₱ ₱ ₱ ₱ ₱ ₱ ₱
	Note to Physician:
This product is	s a pyrethroid. If large amounts have been ingested, milk cream ååd cther digestible

fats and oils may increase absorption and so should be avoided.

150 . 4.

READ ALL DIRECTIONS PRIOR TO APPLICATION OF THE PRODUCT

But the control of the state of

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or grass grown for seed, or for research purposes.

In New York State, the product may NOT be applied to any grass or turf areas within 100 feet of a water body (lake, pond, river, stream, wetland or drainage ditch). In New York State, do make a single repeat application of this product if there are signs of renewed insect activity, but not sooner that two weeks after the first application. Do not apply more than 0.2 lbs. active ingredient (345 lbs. of this product) per acre per year.

Do not apply more than 800 lbs. of this product per acre per year.

Do not allow children and pets to enter the treated area until dry.

This product can be used to control the following lawn insects:

Annual Bluegrass Weevil (Hyperodes) (Adult)

Armyworms Black Turfgrass Ataenius (Adult)

Chinch bugs Fire Ants

Leafhoppers

Millipedes
Pavement Ants

Red Ants

Sowbugs Weevils Ants

Billbugs (Adult) Centipedes Cutworms

Imported Fire Ants

Mealybugs Mole Crickets Pillbugs

Sod Webworms (Lawn Moth)

Ticks

Treat when insects first appear. Apply this product at a rate of 2 to 4 pounds per 1000 square feet evenly over the turfgrass area using a properly calibrated drop or rotary spreader. Check frequently to make sure the equipment is working properly and the granules are being evenly distributed. A more uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. Best results will occur if the treated area is thoroughly irrigated with water immediately after application.

For more difficult to control pests such as ants, fleas, imported fire ants, mole crickets and ticks, a higher application rate may be needed. In this case, apply 4 to 8 pounds per 1000 square feet.

Comments:

Armyworms, Cutworms; and Sod Webworms: For best results, the treated area should be irrigated immediately after application with up to 0.1 inches of water to activate the insecticide.

Annual Bluegrass Weevil: Treatment should be made to control the adult weevils as they migrate from their overwintering sites and move into grass areas. Contact your State Cooperative Extension Service for more specific information.

9498-455 Page 4975 5/4

Billbugs: Apply when adult billbugs are first observed during April and May. Consult your State Cooperative Extension Service for information on degree days and timing in your region.

Part 3 Table 1 County makes to option and part appropriate factories 4.

Black Turfgrass Ataenius: Apply treatments during May and July to control both generations of adults. The May application should be made at the same time as the full bloom stage of Vanhoutte spiraea and the Horse chestnut. The July application should coincide with the blooming of Rose of Sharon.

Chinch Bugs: These bugs infest the base of the plant and are usually found in the thatch layer. Watering the treated area with up .25 inches immediately after application will result in quicker control.

Flea Larvae: These larvae develop in the soil and shady areas. Irrigate the treated area with up to 0.5 inches of water immediately after application.

Imported Fire Ants: The best control will be reached by a combination of broadcast applications and mound drenching. When mound drenching, a product such as Talstar Lawn and Tree Flowable, must be used. Irrigation should be performed prior to application if the soil is not moist. For best results, apply in cool weather. The early morning or late evening hours should suffice.

Mole Cricket Adults: These adults can be difficult to control because of their extremely active stage in the early spring. Applications should be made as late in the day as possible and be watered up to 0.5 inches immediately after treatment. If the soil is not moist prior to application, it should be irrigated to ensure best control.

Mole Cricket Nymphs: Optimum control can be reached by treating young nymphs because they are more susceptible to the insecticide. Irrigate soil if it is not moist before application. Treatment should be made as late in the day as possible. Water the treated area with up to 0.5 inches directly after application.

Ticks: Do not make spot applications. Treat the entire area where ticks may occur. The higher application rate might be needed if heavy leaf litter or dense ground cover exists. Retreatment might also be necessary due to animals reintroducing new populations.

Deer Ticks: These ticks have a life cycle that ranges over a two year period and involves four life stages. Treatments should be applied in mid- to late-spring to control larvae and nymphs that are present on the soil.

American Dog Ticks: These ticks tend to gather along paths or roadways where humans are likely to be found.

SUGGESTED SPREADER SETTINGS

Note: following will be a list of application spreaders and appropriate settings. This list will vary dependent on market of spreaders. (this note will note appear on final printed label).

Note: These spreader settings were established using standard equipment available from the spreader manufacturer at swath widths and speeds typically used within the industry. It is recommended that all spreader equipment be calibrated at the time of application to achieve the desired application rate.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

6/6

Back Panel (for internal uso only, will not appear on final label)

STORAGE: Keep out of reach of children and animals. Store in original container only. Store in cool, dry place and avoid excess heat.

PESTICIDE DISPOSAL: Wastes resulting from the use of the product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.
(AP031300/RVnot11901)

Manufactuered by: The Andersons Lawn Fertilizer Division d/b/a Free Flow Fertilizer PO Box 119 Maumee, OH 43537

Date Printed	11-19/01 notification
Date Approved	08. 9.00
Supersedes	3/13/00
Label Changes	Notification: WA metals statement, revised warranty
Eilename	9798-158 doc
	1 AP031300/RVnot11301

