

9198-163

04/13/2000

1/14

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460	EPA REGISTRATION NO. 9198-163	DATE OF ISSUANCE APR 13 2000
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	TERM OF ISSUANCE Conditional NAME OF PESTICIDE PRODUCT The Andersons Fertilizer with 0.38% Barricade®	
NAME AND ADDRESS OF REGISTRANT (Include ZIP code) <div style="text-align: center;"> <div>┌</div> <div>└</div> </div> The Andersons Lawn Fert. Division d/b/a Free Flow Fertilizer P.O. Box 119 Maumee, OH 43537		
NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.		
On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.		
A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith. Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.		
This product is registered in accordance with FIFRA section 3(c)(7)(A), provided that you:		
<ol style="list-style-type: none"> 1. Submit "Discussion of Impurities", EPA Guideline 830-1670 and "Corrosion Characteristics", EPA Guideline 830-6320 within one year from the date of this registration. 2. Add the phrase "EPA Registration No. 9198-163" to the label before you release the product for shipment. 3. Submit one (1) copy of the final printed label, before you release the product for shipment under the stamped enclosed label. 		
If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.		
<div style="text-align: right;"> Joanne I. Miller Product Manager (23) Herbicide Branch Registration Division (7505C) </div> <div style="text-align: left;"> Enclosure </div>		
<input type="checkbox"/> ATTACHMENT IS APPLICABLE		
SIGNATURE OF APPROVING OFFICIAL		DATE APR 13 2000
<i>Joanne I. Miller</i>		

2/14

9198-XXXC
Page 1 of 13

(Front Panel)
Commercial Use
(for internal use only, will not appear on final label)

The Andersons Fertilizer with 0.38% Barricade®

Active Ingredient:	
Prodiamine (CAS No. 29091-21-2)	0.38%
Other Ingredients:	99.62%
Total	100.00%

Barricade – trademark of Novartis

For selective preemergence control of grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals
- established perennial & wildflower plantings

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional Precautionary Statements and First Aid

EPA Reg. No. 9198-

EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

Underlined letter is first letter used in run code

Net Weight
Covers

ACCEPTED
with COMMENTS
In EPA Letter Dated

APR 13 2000

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

9198-163

(Back Panel)
Commercial Use
(for internal use only, will not appear on final label)

GUARANTEED ANALYSIS

Total Nitrogen (N) XXX.XX%
Available Phosphours (K₂O) XXX.XX%
Soluble Potash (P₂O₅) XXX.XX%
Plant Nutrients Derived From:

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals**CAUTION**

Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact, may cause allergic reactions in some individuals.

First Aid

If on skin: Wash with soap and water. Rinse thoroughly.

If inhaled: Remove victim to clear air.

If in eyes: Flush thoroughly with water for several minutes. Contact a physician if irritation persists.

Personal Protective Equipment (PPE)

Mixers and loaders who handle this pesticide must wear:

- Waterproof gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- After handling this product, immediately wash the outside of gloves before removing them, then remove gloves and all other PPE. Immediately wash thoroughly and change into clean clothing.

4/14

(Back Panel)
Commercial Use
(for internal use only, will not appear on final label)

Environmental Hazards

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations substantially above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

The Andersons Fertilizer with 0.38% Barricade is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals
- established perennial & wildflower plantings

The Andersons Fertilizer with 0.38% Barricade controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when The Andersons Fertilizer with 0.38% Barricade is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.


ESTABLISHED TURF

The Andersons Fertilizer with 0.38% Barricade is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of The Andersons Fertilizer with 0.38% Barricade that may be applied per year is given for each turf grass species in the **Maximum Annual Rates** section of this label.

Most effective weed control in turf grasses will be obtained when The Andersons Fertilizer with 0.38% Barricade is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

(Back Panel)
Commercial Use
(for internal use only, will not appear on final label)

CRABGRASS SEED GERMINATION DATES



Rates of Application

The Andersons Fertilizer with 0.38% Barricade may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. The Andersons Fertilizer with 0.38% Barricade will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by The Andersons Fertilizer with 0.38% Barricade is related to the rate applied.

Maximum Annual Rates

The Andersons Fertilizer with 0.38% Barricade is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Application Rate of The Andersons Fertilizer with 0.38% Barricade/ Calendar Year by Turf Grass Species ¹			
Turf Species	lbs. AI/A	lbs. product /A	lbs. product/ 1000 sq. ft.
Bermudagrass ²	1.5	395	9.1
Bahiagrass			
Centipedegrass			
Seashore Paspalum			
St. Augustinegrass			
Tall Fescue (including turf-type)			
Zoysia			
Buffalograss	1.0	263	6.0
Kentucky Bluegrass			
Perennial Ryegrass			
Creeping Red Fescue	0.75	197	4.5
Creeping Bentgrass	0.65	171	3.9

¹These are the maximum rates per calendar year by species limitations.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 0.5 lb. AI/A (3.14 LB./1000 sq. ft.). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded.

6/14

(Back Panel)
Commercial Use
(for internal use only, will not appear on final label)**Weeds Controlled**

When used in accordance with this label The Andersons Fertilizer with 0.38% Barricade is approved for control of the following weeds:

Barnyardgrass	Kochia
Bluegrass, Annual (<i>Poa annua</i>)	Lambsquarter, Common
Carpetweed	Lovegrass
Chickweed, Common	Panicum, (Texas, Fall, Browntop)
Chickweed, Mouseear (from seed)	Pigweed
Crabgrass (Large, Smooth)	Purslane, Common
Crowfootgrass	Pusley, Florida
Cupgrass, Woolly	Rescuegrass ³⁾
Foxtails, Annual	Shepherd's Purse ²⁾
Goosegrass ¹⁾	Signalgrass, Broadleaf
Henbit	Speedwell, Persian
Itchgrass	Sprangletop
Johnsongrass (from seed)	Spurge, Prostrate
Junglerice	Witchgrass
Knotweed	Woodsorrel, Yellow (from seed)

¹⁾In many areas a single application of 171 to 395 lbs./A (3.9 to 9.1 lbs./1,000 sq.ft.) The Andersons Fertilizer with 0.38% Barricade will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 171 to 263 lbs./A (3.9 to 6.0 lbs./1,000 sq. ft.) followed after 60 to 90 days by a second application at doses that would not exceed those given in the **Maximum Application Rates Table**. Do not exceed the maximum rate for turf grass species listed in the **Maximum Application Rates Table** above.

²⁾Applications for this weed should be made in late summer, fall, or winter prior to germination.

³⁾Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

When to Apply After Overseeding Turf

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if The Andersons Fertilizer with 0.38% Barricade is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

When to Overseed After Application

The Andersons Fertilizer with 0.38% Barricade will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.]

7/14

(Back Panel)
Commercial Use
(for internal use only, will not appear on final label)

The Andersons Fertilizer with 0.38% Barricade Rate (lbs/A)		Interval Before Overseeding		
Product	ai	North	Transition	South
131.5	0.50	4 mo	4 mo	4 mo
171	0.65	5 mo	4 mo	4 mo
197	0.75	6 mo	5 mo	5 mo
213	0.81	---	6 mo	6 mo
258	0.98	---	7 mo	7 mo
302	1.15	---	---	9 mo
342	1.30	---	---	10 mo
395	1.50	---	---	12 mo

Use Precautions*For final printed labels which do include ornamental uses:*

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

For final printed labels which do not include ornamental uses:

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

The following precautions apply to the use of The Andersons Fertilizer with 0.38% Barricade in turf grasses and lawns:

1. Application of The Andersons Fertilizer with 0.38% Barricade may thin emerged annual bluegrass and newly over-seeded grasses.
2. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if The Andersons Fertilizer with 0.38% Barricade is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
3. Do not apply to newly set sod until the following year.
4. Application of The Andersons Fertilizer with 0.38% Barricade to turf stressed by drought, low fertility, or pest damage may result in turf injury.
5. Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
6. Do not apply The Andersons Fertilizer with 0.38% Barricade to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa annua*) are desirable species.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift.

Do not graze or feed livestock forage cut from areas treated with The Andersons Fertilizer with 0.38% Barricade.

(Back Panel)
Commercial Use
(for internal use only, will not appear on final label)

Do not apply this product through any type of irrigation system unless otherwise instructed by this label.

Do not apply aerially.

Do not apply to golf course putting greens.

LANDSCAPE ORNAMENTALS

The Andersons Fertilizer with 0.38% Barricade may be applied for residual preemergence weed control in ornamentals (including established perennial and wildflower plantings).

USE RATES

Apply The Andersons Fertilizer with 0.38% Barricade at 171 to 395 lbs./A in fall and/or spring. Equivalent applications for smaller areas are 3.9 to 9.1 lbs./1000 sq. ft. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 395 lbs./A (9.1 lbs./1,000 sq. ft.).

APPLICATION TIMING AND INFORMATION

All Applications

The Andersons Fertilizer with 0.38% Barricade is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when The Andersons Fertilizer with 0.38% Barricade is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when The Andersons Fertilizer with 0.38% Barricade is applied to soil free of clods, weeds, and debris such as leaves and mulch. Prior to application of The Andersons Fertilizer with 0.38% Barricade, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

Tolerant Ornamental Species

The Andersons Fertilizer with 0.38% Barricade will not harm most trees, shrubs, vines, and flowers. The species listed below in Table 1 are tolerant to The Andersons Fertilizer with 0.38% Barricade. The Andersons Fertilizer with 0.38% Barricade is approved for application, except in California, to the species in Table 2. The Andersons Fertilizer with 0.38% Barricade may be applied over the top of the listed species.

When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to The Andersons Fertilizer with 0.38% Barricade.

Table 1 - Tolerant Ornamental Species - All States

Scientific name	Common name
<i>Abies</i> spp.	Fir species** (Balsam, Fraser, Noble, etc.)
<i>Acer palmatum</i>	Japanese Maple

(Back Panel)
Commercial Use
(for internal use only, will not appear on final label)

Acer platanoides
Actinidia chinensis
Agapanthus africanus
Arctostaphylos densiflora
Arctotheca calendula
Aucuba japonica
Berberis gladwynensis
Berberis julianae
Berberis mentorensis
Berberis thunbergii
Berberis verruculosa
Buxus microphylla
Callistemon viminalis
Calluna vulgaris
Carpobrotus edulis
Cassia artemisioides
Ceanothus rigidus
Chamaecyparis pisifera
Cleyera japonica
Citrus spp.
Cornus florida
Cornus stolonifera
Cortaderia selloana
Cotoneaster apiculatus
Cotoneaster buxifolius
Cotoneaster dammeri
Cotoneaster microphyllus
Crataegus spp.
Cupressus sempervirens
Delosperma alba
Dodonaea viscosa
Elaeagnus pungens
Euonymus fortunei
Euonymus japonica
Euonymus kiautschovicks
Fatsia japonica
Forsythia intermedia
Forsythia viridissima
Gardenia jasminoides
Gladiolus spp.
Hedera helix
Hibiscus
Hibiscus Rosa-sinensis
Ilex cornuta
Ilex crenata
Ilex opaca
Ilex pernyi
Ilex vomitoria
Iris spp.
Jasminium nudiflorum
Juniperus chinensis
Juniperus conferta

Norway Maple
Kiwi*
Lily-of-the-Nile (African Lily)
Vine Hill Manzanita
Cape Weed
Japanese Aucuba
Barberry
Wintergreen Barberry
Mentor Barberry
Japanese Barberry
Warty Barberry
Japanese Boxwood
Weeping Bottlebrush
Scotch Heather
Hottentot Fig (Ice Plant)
Feathery Cassia
Wild Lilac
False Cypress
Cleyera
Citrus species*
Flowering Dogwood
American Dogwood
Pampas Grass
Cranberry Cotoneaster
Cotoneaster
Bearberry Cotoneaster
Rockspray Cotoneaster
Hawthorne
Italian Cypress
White Trailing Ice Plant
Hop Bush
Silverberry
Wintercreeper
Japanese Spindle Tree (Evergreen Euonymus)
Spreading Euonymus
Japanese Aralia
Border Forsythia
Greenstem Forsythia
Gardenia, Cape-Jasmine
Gladiolus species**
English Ivy
Rose of Sharon**
Chinese Hibiscus
Chinese Holly
Japanese Holly
American Holly
Holly
Yaupon Holly
Iris species**
Winter Jasmine
Chinese Juniper
Shore Juniper

(Back Panel)
Commercial Use
(for internal use only, will not appear on final label)

<i>Juniperus horizontalis</i>	Creeping Juniper
<i>Juglans</i> spp.	Walnut*
<i>Justicia brandegeana</i>	Shrimp Plant
<i>Lagerstromia indica</i>	Crape Myrtle
<i>Ligustrum amurense</i>	Amur Privet
<i>Ligustrum japonicum</i>	Japanese Privet
<i>Ligustrum lucidum</i>	Glossy Privet (Wax-Leaf)
<i>Liriope muscari</i>	Big Blue Lillyturf
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>Lonicera tatarica</i>	Tatarian Honeysuckle
<i>Magnolia</i> spp.	Magnolia species**
<i>Malephora luteola</i>	Ice Plant
<i>Malus</i> spp.	Crabapple*
<i>Nandina domestica</i>	Heavenly Bamboo
<i>Narcissus</i> spp.	Narcissus species**
<i>Nerium</i> spp.	Oleander
<i>Olea europaea</i>	Olive*
<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendrum arboreum</i>	Sourwood
<i>Persea americana</i>	Avocado*
<i>Photinia fraseri</i>	Fraser's Photinia (Redtip)
<i>Picea</i> spp.	Spruce species** (Colorado Blue, Norway, etc.)
<i>Pieris japonica</i>	Lily-of-the-Valley Shrub
<i>Pinus brutia</i>	Calabrian Pine
<i>Pinus canariensis</i>	Canary Island Pine
<i>Pinus elliotii</i>	Slash Pine
<i>Pinus halepensis</i>	Aleppo Pine
<i>Pinus nigra</i>	Austrian Black Pine
<i>Pinus palustris</i>	Longleaf Pine
<i>Pinus radiata</i>	Monterey Pine
<i>Pinus strobus</i>	Eastern White Pine
<i>Pinus sylvestris</i>	Scotch Pine
<i>Pinus taeda</i>	Loblolly Pine
<i>Pinus thunbergiana</i>	Japanese Black Pine
<i>Pinus virginiana</i>	Virginia Pine
<i>Pistacia</i> spp.	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>Pittosporum tobira</i>	Japanese Pittosporum
<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Prunus laurocerasus</i>	English Laurel
<i>Prunus</i> spp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Pseudotsuga menziesii</i>	Douglas Fir**
<i>Pyracantha coccinea</i>	Firethorn Scarlet
<i>Pyracantha fortuneana</i>	Firethorn
<i>Pyracantha koidzumii</i>	Firethorn
<i>Pyrus</i> spp.	Bradford Pear spp.
<i>Quercus rubra</i>	Oak species
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Rhododendron</i> (including Azalea)	'Coral Bells' 'Formosa'

(Back Panel)
Commercial Use
(for internal use only, will not appear on final label)

<i>Rosa banksiae</i>	'Hino-crimson'
<i>Rosmarinus officinalis</i>	'PJM'
<i>Rumohra adiantiformis</i>	'Roseum Elegans'
<i>Santolina virens</i>	Lady Bank's Rose
<i>Sedum album</i>	Rosemary*
<i>Syzygium paniculatum</i>	Leatherleaf Fern
<i>Taxus cuspidata</i>	
<i>Taxus media</i>	Stonecrop
<i>Thuja occidentalis</i>	Japanese Boxcherry
<i>Trachelospermum asiaticum</i>	Japanese Yew
<i>Tsuga canadensis</i>	Yew
<i>Tulipa</i> spp.	American Arborvitae
<i>Viburnum japonicum</i>	Star Jasmine
<i>Viburnum odoratissimum</i>	Canada Hemlock
<i>Viburnum plicatum</i>	Tulip species
<i>Viburnum rigidum</i>	Japanese Viburnum
<i>Viburnum tinus</i>	Sweet Viburnum
<i>Viburnum trilobum</i>	Japanese Snowball
<i>Viburnum wrightii</i>	Canary Island Viburnum
<i>Vinca major</i>	Laurustinus
<i>Vinca minor</i>	Cranberry Bush
<i>Vitis</i> spp.	Leatherleaf Viburnum
<i>Weigela florida</i>	Vinca
<i>Yucca aloifolia</i>	Dwarf Periwinkle
<i>Yucca filamentosa</i>	Grape*
	Old Fashioned Weigela
	Spanish Bayonet
	Yucca, Adam's Needle

*Do not use on food producing trees, vines, or plants.

**Not for use on container grown plants.

Table 2 - Tolerant Ornamental Species/Varieties - All States Except CA

Scientific name	Common name
<i>Abelia grandiflora</i>	Abelia: Sherwood
<i>Achillea</i> spp.	Yarrow: King Edward
<i>Agapanthus orientalis</i>	
<i>Akebia quinata</i>	Five-Leaf or Chocolate Vine
<i>Allium cernuum</i>	Lady's Leek, Nodding Onion
<i>Anemone hybrida</i>	Japanese Anemone
<i>Aquilegia</i> spp.	Aquilegia: Red and Gold
<i>Artemisia</i> spp.	Wormwood; Silver Mound, Castle
<i>Aster</i> spp.	Aster: Bonny Blue, Purple Dome
<i>Aster X frikartii</i>	
<i>Athyrium filix-femina</i>	Lady Fern; Fern Lady
<i>Begonia</i> spp.	Fibrous Begonia: Hardy Grandis
<i>Bergenia cordifolia</i>	
<i>Boltonia asteroides</i>	Snowbank
<i>Bougainvillea</i> spp.	Bougainvillea

(Back Panel)
Commercial Use
(for internal use only, will not appear on final label)

Buddleia davidii
Callistemon citrinus
Campanula carpatica
Campis X tagliabuana
Ceratostigma plumbaginoides
Chrysanthemum nipponicum
Coreopsis spp.

Crocosmia spp.
Delosperma spp.
Delphinium spp.
Dianthus deltoides
Dianthus gratianopolitanus
Echinacea pupurea
Forsythia suspensa
Gaillardia spp.
Gaura spp.
Gentiana dahurica
Geranium cinereum
Gypsophila repens
Helianthemum spp.
Hemerocallis spp.
Heucherella spp.
Hibiscus spp.
Hosta plantaginea
Hosta sieboldiana
Houttuynia cordata var. *variegata*
Hydrangea macrophylla
Inula ensifolia
Iris ensata
Iris siberica
Juniperus davurica
Lagerstromia indica x fauriei
Lantana montevidensis
Lavender spp.
Leontopodium alpinum
Ligustrum sinense
Lilium spp.
Lirope muscari var. *variegata*
Lirope spicata
Lobelia cardinalis
Loropetalum chinense
Lythrum spp.
Miscanthus sinensis
Oenothera missourensis
Osmanthus heterophyllus
Paeonia suffruticosa
Pennisetum setaceum
Perovskia atriplicifolia
Physostegia virginiana
Quercus shumardii
Rhaphiolepis umbellata

Butterfly-Bush (Dwarf Blue); Royal Red
Crimson Bottlebrush
Tussock Bellflower; (White Clips)
Trumpet Creeper, Trumpet Flower; Madame Galen

Coreopsis (*Calliopsis*): Early Sunrise,
Moonbeam

Lucifer
Cooperi Pink
Larkspur; Blue Elf
Dianthus, Maiden Pinks 'Zing'
Cheddar Pink
Coneflower, Purple; Magnus
Weeping Forsythia
Gaillardia, Blanket Flower: 'Goblin'

Gentian
Cranesbill
Baby's Breath
Sunrose
Daylily: Aztec Gold, Stella De Oro, Tender Love
Coral Bell; Bridget Bloom
Mallow; Disco Belle White
Hosta, Plantain Lily (Fragrant)
Hosta, 'Searsucker'

Bigleaf Hydrangea

Sword-Leaved Iris; Jodlesong
Siberian Iris; Cabernet
Parsoni
Crape Myrtle; Tuscarora
Weeping Lantana
Lavender; Munstead
Edelweiss
Chinese Privet; Variegata
Lily: Jazz
Lirope, Variegated
Lirope, Creeping
Cardinal Flower, Indian Pink
Burgundy
Loosestrife; Modern Pink
Yaku Jima, Silberfeder
Evening Primrose
Osmanthus (False Holly): Gulf Tide
Tree Peony
Fountain Grass (Dwarf)

Dragonhead, False; Vivid
Oak, Shumard's Red
Yedda Hawthorne

(Back Panel)
Commercial Use
(for internal use only, will not appear on final label)

Rhododendron
(including azalea)

'Delaware Valley White'
'Flame Creeper'
'Girard Crimson'
'George L. Tabor'
Wakeiebisu

Rudbeckia spp.

White Gumpo

Saxifraga spp.

Black-Eyed Susan: Goldstrum

Scabiosa spp.

Saxifrage; Purple Dome

Sedum caudicola

Pincushion Flower

Sedum dasyphyllum

Stonecrop; Lidakense

Sedum spurium

Stonecrop

Spiraea bumalda

Stonecrop; Dragon's Blood

Syzyglum paniculatum

Spirea: Anthony Waterer

Teucrium spp.

Australian Brushcherry

Thalictrum dipterocarpum

Germander

Veronica spp.

Meadow Rue

Viburnum suspensum

Veronica, Speedwell; Sunny Border

Arrowood Viburnum

*Do not use on food producing trees, vines or plants.

**Not for use on container grown plants.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container away from fertilizer, feed, or food stuffs and separated from other pesticides.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty container into application equipment. Then dispose of empty bag or box in a sanitary landfill or at an incineration facility; or, if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

(AP)

14/14

9198-XXXC
Page 13 of 13

(Back Panel)
Commercial Use
(for internal use only, will not appear on final label)

Manufactured by:
The Andersons Lawn Fertilizer Division
D/b/a Free Flow Fertilizer Division
PO Box 119
Maumee, OH 43537

Date Printed: 02/08/00
Date Approved:
Supersedes:
Label changes:
Filename: barricade