910	98 - 163	ina form.	8-30	-200 J	nnroved	I. OMB No.	 2070-00 <i>6</i>	30. Approved expires 2-28-9
United States Environmental Protection Age Washington, DC 20460				ency A		Registra Amenda Other	ation	OPP Identifier Number
		Application	n for Pe	sticide - Se	ction	1		
1. Company/Product Number 9198-163			ł	EPA Product Me iller	anager		1 -	roposed Classification None Restricted
4. Company/Product (Name) The Andersons Fertilizer with 0.38	8% Barricade		P 1	и# З				
5. Name and Address of Applicant (Include ZIP Code) The Andersons Lawn Fert. Div., Inc. d/b/a Free F Fert. PO Box 119 Maumee, OH 43537			6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No					
Check if this is a	new address		P	roduct Name				
			Sectio	n - II				
Amendment - Explain beling Resubmission in response		Final printed (abels in repsonse to Agency letter dated "Me Too" Application.				NOTIFICATION		
Notification - Explain below.				Other - Explain below. AUG 3 0 2001				AUG 3 0 2001
This notification is consistent with the confidentical statement of formula of notification is not consistent with the penalties under sections 12 and 14 c	this product. undert terms of PR Notice 98	had that it is a vi-	plation of 18 U.S	i.C. Sec. 1001 to will duct may be in viola	tully make	e anv false stati	ement to EF	A. I further undersant that if this
1. Material This Product Will Be	Packaged In:						_	
Yes No	Yes No Yes' it Packaging wgt.	No. per container	Water Soli Yes No If "Yes" Package w	No. per		2. Type of	Metal Plastic Glass Peper	Specify)
3. Location of Net Contents Info	ļ	4. Size(s) Ret	ail Container		5. Lo	cation of Lal		ons
8. Manner in Which Lebel is Affin	xed to Product	Lithog Paper Stenci	raph glued lad	[] Oth	ner		<u>-</u> -	
			Section	ı - IV				
1. Contact Point (Complete item	s directly below fo	or identificatio	n of individua	i to be contacted	d, if nec	essery, to pi	ocess this	s application.)
Name Dawn L. Walters						Telephor 334-262-	ne No. (Include Area Code) 2766	
I certify that the statement I acknowledge that any kn both under applicable law.			all attachmer					6. Date Application Received (Stamped;
2. Signature Dawn Hullturs			3. Title Regulatory A	dministrator				
4. Typed Name Dawn L. Walters			5. Date 8/22/01				1	

9195-163 Page 1 of 12

(Front Panel)

The Andersons Fertilizer with 0.38% Barricade®

 Active Ingredient:
 0.38%

 Prodiamine (CAS No. 29091-21-2)
 0.38%

 Other Ingredients:
 99.62%

 Total
 100.00%

Barricade - trademark of Novartis

NOTIFICATION

AUG 3 0 2001

For selective preemergence control of grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals
- · established perennial & wildflower plantings

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional Precautionary Statements and First Aid

EPA Reg. No. 9198-163 EPA Est. No. 9198-OH-1<u>M</u>, 9198-OH-2<u>B</u>, 9198-AL-001<u>A</u> Underlined letter is first letter used in run code

Net Weight Covers

GUARANTEED ANALYSIS

Total Nitrogen (N) XXX,XX% Available Phosphours (K₂O) XXX.XX% Soluble Potash (P₂O₅) XXX.XX% Plant Nutrients Derived From:

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact, may cause allergic reactions in some individuals.

First Aid

If on skin: Wash with soap and water. Rinse thoroughly.

If inhaled: Remove victim to clear air.

If in eyes: Flush thoroughly with water for several minutes. Contact a physician if irritation persists.

Personal Protective Equipment (PPE)

Mixers and loaders who handle this pesticide must wear:

Waterproof gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- After handling this product, immediately wash the outside of gloves before removing them, then remove gloves and all other PPE. Immediately wash thoroughly and change into clean clothing.

Environmental Hazards

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations substantially above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

The Andersons Fertilizer with 0.38% Barricade is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens) and lawns
- · landscape ornamentals
- established perennial & wildflower plantings


The Andersons Fertilizer with 0.38% Barricade controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when The Andersons Fertilizer with 0.38% Barricade is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

ESTABLISHED TURF

The Andersons Fertilizer with 0.38% Barricade is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of The Andersons Fertilizer with 0.38% Barricade that may be applied per year is given for each turf grass species in the **Maximum Annual Rates** section of this label.

Most effective weed control in turf grasses will be obtained when The Andersons Fertilizer with 0.38% Barricade is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

CRABGRASS SEED GERMINATION DATES



Rates of Application

The Andersons Fertilizer with 0.38% Barricade may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. The Andersons Fertilizer with 0.38% Barricade will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by The Andersons Fertilizer with 0.38% Barricade is related to the rate applied.

Maximum Annual Rates

The Andersons Fertilizer with 0.38% Barricade is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Application Rate of The Andersons Fertilizer with 0.38% Barricade/ Calendar Year by Turf Grass Species ¹				
Turf Species	lbs. Al/A	lbs. product /A	lbs. product/ 1000 sq. ft.	
Bermudagrass ² Bahiagrass Centipedegrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type) Zoysia	1.5	395	9.1	
Buffalograss Kentucky Bluegrass Perennial Ryegrass	1.0	263	6.0	
Creeping Red Fescue Creeping Bentgrass	0.75 0.65	197 171	3.9	

¹These are the maximum rates per calendar year by species limitations.

Weeds Controlled

When used in accordance with this label The Andersons Fertilizer with 0.38% Barricade is approved for control of the following weeds:

Barnyardgrass

Bluegrass, Annual (Poa annua)

Carpetweed

Chickweed, Common

Chickweed, Mouseear (from seed)

Crabgrass (Large, Smooth)

Crowfootgrass Cupgrass, Woolly Foxtails, Annual Goosegrass¹⁾ Henbit

Itchgrass

Johnsongrass (from seed)

Junglerice Knotweed Kochia

Lambsquarter, Common

Lovegrass

Panicum, (Texas, Fall, Browntop)

Piaweed

Purslane, Common Pusley, Florida Rescuegrass³⁾ Shepherd's Purse²⁾ Signalgrass, Broadleaf Speedwell, Persian

Sprangletop Spurge, Prostrate

Witchgrass

Woodsorrel, Yellow (from seed)

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 0.5 lb.Al/A (3.14 LB./1000 sq. ft.). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded.

¹ In many areas a single application of 171 to 395 lbs./A (3.9 to 9.1 lbs./1,000 sq.ft.) The Andersons Fertilizer with 0.38% Barricade will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial

application of 171 to 263 lbs./A (3.9 to 6.0 lbs./1,000 sq. ft.) followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Application Rates Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Application Rates Table above.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

When to Apply After Overseeding Turf

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if The Andersons Fertilizer with 0.38% Barricade is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

When to Overseed After Application

The Andersons Fertilizer with 0.38% Barricade will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.]

The Andersons Fe Barricade F		Interval Before Overseeding			
Product	ai	North	Transition	South	
131.5	0.50	4 mo	4 mo	4 mo	
171	0.65	5 mo	4 mo	4 mo	
197	0.75	6 mo	5 mo	5 mo	
213	0.81		6 mo	6 mo	
258	0.98		7 mo	7 mo	
302	1.15			9 mo	
342	1.30			10 mo	
395	1.50			12 mo	

Use Precautions

For final printed labels which do include ornamental uses:

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

For final printed labels which do not include ornamental uses:

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

The following precautions apply to the use of The Andersons Fertilizer with 0.38% Barricade in turf grasses and lawns:

²⁾Applications for this weed should be made in late summer, fall, or winter prior to germination.

³⁾Suppression only.

- 1. Application of The Andersons Fertilizer with 0.38% Barricade may thin emerged annual bluegrass and newly over-seeded grasses.
- Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if The Andersons Fertilizer with 0.38% Barricade is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
- 3. Do not apply to newly set sod until the following year.
- 4. Application of The Andersons Fertilizer with 0.38% Barricade to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
- 6. Do not apply The Andersons Fertilizer with 0.38% Barricade to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa annua*) are desirable species.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift.

Do not graze or feed livestock forage cut from areas treated with The Andersons Fertilizer with 0.38% Barricade.

Do not apply this product through any type of irrigation system unless otherwise instructed by this label.

Do not apply aerially.

Do not apply to golf course putting greens.

LANDSCAPE ORNAMENTALS

The Andersons Fertilizer with 0.38% Barricade may be applied for residual preemergence weed control in ornamentals (including established perennial and wildflower plantings).

USE RATES

Apply The Andersons Fertilizer with 0.38% Barricade at 171 to 395 lbs./A in fall and/or spring. Equivalent applications for smaller areas are 3.9 to 9.1 lbs./1000 sq. ft. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 395 lbs./A (9.1 lbs./1,000 sq. ft.).

APPLICATION TIMING AND INFORMATION

All Applications

The Andersons Fertilizer with 0.38% Barricade is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when The Andersons Fertilizer with

0.38% Barricade is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when The Andersons Fertilizer with 0.38% Barricade is applied to soil free of clods, weeds, and debris such as leaves and mulch. Prior to application of The Andersons Fertilizer with 0.38% Barricade, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

Tolerant Ornamental Species

The Andersons Fertilizer with 0.38% Barricade will not harm most trees, shrubs, vines, and flowers. The species listed below in Table 1 are tolerant to The Andersons Fertilizer with 0.38% Barricade. The Andersons Fertilizer with 0.38% Barricade is approved for application, except in California, to the species in Table 2. The Andersons Fertilizer with 0.38% Barricade may be applied over the top of the listed species.

When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to The Andersons Fertilizer with 0.38% Barricade.

Table 1 - Tolerant Ornamental Species - All States

Scientific name	Common name
Abies spp.	Fir species** (Balsam, Fraser, Noble, etc.)
Acer palmatum	Japanese Maple
Acer platanoides	Norway Maple
Actinidia chinensis	Kiwi*
Agapanthus africanus	Lily-of-the-Nile (African Lily)
Arctostaphylos densiflora	Vine Hill Manzanita
Arctotheca calendula	Cape Weed
Aucuba japonica	Japanese Aucuba
Berberis gladwynensis	Barberry
Berberis julianae	Wintergreen Barberry
Berberis mentorensis	Mentor Barberry
Berberis thunbergii	Japanese Barberry
Berberis verruculosa	Warty Barberry
Buxus microphylla	Japanese Boxwood
Callistemon viminalis	Weeping Bottlebrush
Calluna vulgaris	Scotch Heather
Carpobrotus edulis	Hottentot Fig (Ice Plant)
Cassia artemisioides	Feathery Cassia
Ceanothus rigidus	Wild Lilac
Chamaecyparis pisifera	False Cypress
Cleyera japonica	Cleyera
Citrus spp.	Citrus species*
Cornus florida	Flowering Dogwood
Cornus stolonifera	American Dogwood
Cortaderia selloana	Pampas Grass
Cotoneaster apiculatus	Cranberry Cotoneaster
Cotoneaster buxifolius	Cotoneaster
Cotoneaster dammeri	Bearberry Cotoneaster

Cotoneaster microphyllus

Crataegus spp.

Cupressus sempervirens

Delosperma alba Dodonaea viscosa Elaeagnus pungens Euonymus fortunei Euonymus japonica

Euonymus kiautschovicks

Fatsia japonica Forsythia intermedia Forsythia viridissima Gardenia jasminoides

Gladiolus spp. Hedera helix Hibiscus

Hibiscus Rosa-sinensis

Ilex cornuta
Ilex crenata
Ilex opaca
Ilex pernyi
Ilex vomitoria
Iris spp.

Jasminium nudiflorum Juniperus chinensis Juniperus conferta Juniperus horizontalis

Juglans spp.

Justicia brandegeana Lagerstromia indica Ligustrum amurense Ligustrum japonicum Ligustrum lucidum Lirope muscari Lonicera japonica Lonicera tatarica Magnolia spp.

Malus spp.

Nandina domestica Narcissus spp. Nerium spp. Olea europaea

Malephora luteola

Ophiopogon japonicus

Osteospermum fruticosum Oxydendrum arboreum

Persea americana Photinia fraseri Picea spp. Pieris japonica

Pinus brutia

Pinus canariensis

Rockspray Cotoneaster

Hawthorne Italian Cypress

White Trailing Ice Plant

Hop Bush Silverberry Wintercreeper

Japanese Spindle Tree (Evergreen Euonymus)

Spreading Euonymus Japanese Aralia Border Forsythia Greenstem Forsythia Gardenia, Cape-Jasmine

Gladiolus species**

English Ivy

Rose of Sharon**
Chinese Hibiscus
Chinese Holly
Japanese Holly
American Holly

Holly

Yaupon Holly Iris species** Winter Jasmine Chinese Juniper Shore Juniper Creeping Juniper

Walnut*
Shrimp Plant
Crape Myrtle
Amur Privet
Japanese Privet

Glossy Privet (Wax-Leaf)

Big Blue Lillyturf Japanese Honeysuckle Tatarian Honeysuckle Magnolia species**

Ice Plant Crabapple*

Heavenly Bamboo Narcissus species**

Oleander Olive*

Mondo Grass

Trailing African Daisy

Sourwood Avocado*

Frasier's Photinia (Redtip)

Spruce species** (Colorado Blue, Norway, etc.)

Lily-of-the-Valley Shrub

Calabrian Pine Canary Island Pine

9198-163 Page 9 of 12

(Back Panel)

Pinus elliottii
Pinus halepensis
Pinus nigra
Pinus palustris
Pinus radiata
Pinus strobus
Pinus sylvestris
Pinus taeda
Pinus thunbergiana
Pinus virginiana
Pistacia spp.

Pittosporum rhombifolium Pittosporum tobira Podocarpus macrophyllus Prunus laurocerasus

Prunus spp.

Pyracantha coccinea Pyracantha fortuneana Pyracantha koidzumii Pyrus spp. Quercus rubra Rhaphiolepis indica Rhododendron

(including Azalea)

Pseudotsuga menziesii

Rosa banksiae Rosmarinus officinalis Rumohra adiantiformis Santolina virens

Santolina virei Sedum album

. Syzygium paniculatum Taxus cuspidata Taxus media Thuja occidentalis

Trachelospermum asiaticum

Tsuga canadensis

Tulipa spp.

Weigela florida

Viburnum japonicum
Viburnum odoratissimum
Viburnum plicatum
Viburnum rigidum
Viburnum tinus
Viburnum trilobum
Viburnum wrightii
Vinca major
Vitis spp.

Slash Pine
Aleppo Pine
Austrian Black Pine
Longleaf Pine
Monterey Pine
Eastern White Pine
Scotch Pine
Loblolly Pine

Japanese Black Pine

Virginia Pine Pistachio*

Queensland Pittosporum Japanese Pittosporum

Japanese Yew English Laurel

Almond, Apricot, Nectarine, Peach, Plum and Prune*

Douglas Fir**
Firethorn Scarlet
Firethorn

Firethorn

Bradford Pear spp. Oak species Indian Hawthorne 'Coral Bells' 'Formosa'

'Hino-crimson' 'PJM'

'Roseum Elegans' Lady Bank's Rose Rosemary* Leatherleaf Fern

Stonecrop

Japanese Boxcherry
Japanese Yew

Yew

American Arborvitae

Star Jasmine Canada Hemlock Tulip species Japanese Viburnum Sweet Viburnum

Japanese Snowball
Canary Island Viburnum
Laurustinus

Cranberry Bush
Leatherleaf Viburnum

Vinca

Dwarf Periwinkle

Grape*

Old Fashioned Weigela

Yucca aloifolia Yucca filamentosa Spanish Bayonet Yucca, Adam's Needle

*Do not use on food producing trees, vines, or plants.

Table 2 - Tolerant Ornamental Species/Varieties - All States Except CA

Scientific name Common name Abelia grandiflora Abelia: Sherwood Achillea spp. Yarrow: King Edward Agapanthus orientalis Akebia quinata Five-Leaf or Chocolate Vine Allium cernuum Lady's Leek, Nodding Onion Anemone hybrida Japanese Anemone Aquilegia spp. Aquilegia: Red and Gold Artemisia spp. Wormwood; Silver Mound, Castle Aster spp. Aster: Bonny Blue, Purple Dome Aster X frikartii Athyrium filix-femina Lady Fern; Fern Lady Fibrous Begonia: Hardy Grandis Begonia spp. Bergenia cordifolia

Bergenia cordifolia
Boltonia asteroides
Bougainvillea spp.
Buddleia davldii
Butterfly-Bush (Dwarf Blue); Royal Red
Callistemon citrinus
Crimson Bottlebrush

Campanula carpatica Tussock Bellflower; (White Clips)
Campis X tagliabuana Trumpet Creeper, Trumpet Flower; Madame Galen

Ceratostigma plumbaginoides Chrysanthemum nipponicum

Coreopsis spp.

Crocosmia spp. Luci
Delosperma spp. Coo
Delphinium spp. Lark
Dianthus deltoides Diar

Dianthus gratianopolitanus

Echinacea pupurea Forsythia suspensa

Gaillardia spp. Gaura spp.

Gentiana dahurica Geranium cinereum Gypsophila repens Helianthemum spp.

Hemerocallis spp. Heucherella spp. Hibiscus spp. Hosta plantaginea

Hosta sieboldiana Houttuynia cordata var. variegata

Hydrangea macrophylla

Coreopsis (Calliopsis): Early Sunrise,

Moonbeam Lucifer Cooperi Pink Larkspur; Blue Elf

Dianthus, Maiden Pinks 'Zing'

Cheddar Pink

Coneflower, Purple; Magnus

Weeping Forsythia

Gaillardia, Blanket Flower: 'Goblin'

Gentian Cranesbill Baby's Breath Sunrose

Daylily: Aztec Gold, Stella De Oro, Tender Love

Coral Bell; Bridget Bloom Mallow; Disco Belle White Hosta, Plantain Lily (Fragrant)

Hosta, 'Searsucker'

Bigleaf Hydrangea

^{**}Not for use on container grown plants.

Inula ensifolia Iris ensata Iris siberica

Juniperus davurica

Lagerstromia indica x fauriei Lantana montevidensis

Lavender spp.

Leontopodium alpinum

Ligustrum sinense

Lilium spp.

Lirope muscari var. variegata Lirope spicata

Lobelia cardinalis

Loropetalum chinense

Lythrum spp.

Miscanthus sinensis
Oenothera missourensis
Osmanthus botarophyllus

Osmanthus heterophyllus

Paeonia suffruticosa

Pennisetum setaceum Perovskia atriplicifolia

Physostegia virginiana Quercus shumardii Rhaphiolepis umbellata

Rhododendron

(including azalea)

Rudbeckia spp. Saxifraga spp.

Scabiosa spp. Sedum cauticola

Sedum dasyphyllum

Sedum spurium Spiraea bumalda Syzyglum paniculatum

Teucrium spp.

Thalictrum dipterocarpum

Veronica spp.

Viburnum suspensum

Sword-Leaved Iris; Jodlesong

Siberian Iris; Cabernet

Parsoni

Crape Myrtle; Tuscarora Weeping Lantana Lavender; Munstead

Edelweiss

Chinese Privet; Variegata

Lily: Jazz

Lirope, Variegated Lirope, Creeping

Cardinal Flower, Indian Pink

Burgundy

Loosestrife; Modern Pink
Yaku Jima, Silberfeder
Evening Primrose

Osmanthus (False Holly): Gulf Tide

Tree Peony

Fountain Grass (Dwarf)

Dragonhead, False; Vivid Oak, Shumard's Red Yedda Hawthorne 'Delaware Valley White'

'Flame Creeper'
'Girard Crimson'
'George L. Tabor'
Wakeiebisu

White Gumpo
Black-Eyed Susan: Goldstrum
Saxifrage; Purple Dome
Pincushion Flower
Stonecrop; Lidakense

Stonecrop

Stonecrop; Dragon's Blood Spirea: Anthony Waterer Australian Brushcherry

Germander

Meadow Rue

Veronica, Speedwell; Sunny Border

Arrowood Viburnum

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container away from fertilizer, feed, or food stuffs and separated from other pesticides.

^{*}Do not use on food producing trees, vines or plants.

^{**}Not for use on container grown plants.

9198-163

9198-163 Page 12 of 12

(Back Panel)

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty container into application equipment. Then dispose of empty bag or box in a sanitary landfill or at an incineration facility; or, if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (APO41300/RVnot082101)

Manufactured by:

The Andersons Lawn Fertilizer Division, Inc. d/b/a Free Flow Fertilizer Division PO Box 119
Maumee, OH 43537

Date Frented	6.42.00	-
Flate Auground	4.7300	
Supervieles	43 3 00	
Label changes	footfication - warrant,	
F 95 (1) 11:00	0193/163	
su Morre (3.09)	APCHINE Ryneps	