

9198-163

04/29/2004

1/24

9198-163
Page 1 of 12
Homeowner Version

(Front Panel)

The Andersons Fertilizer with 0.38% Barricade® Herbicide

Active Ingredient:

Prodiamine* 0.38%

Other Ingredients: 99.62%

Total 100.00%

*(CAS No. 29091-21-2)

Barricade – trademark of Syngenta Crop Protection

For preemergence control of grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals
- established perennial & wildflower plantings

KEEP OUT OF REACH OF CHILDREN**CAUTION**

See back panel for additional Precautionary Statements and First Aid

EPA Reg. No. 9198-163

EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

Underlined letter is first letter used in run code

Net Weight

Covers

ACCEPTED

APR 29 2004

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

9198-163

2/24

(Back Panel)

The Andersons Fertilizer with 0.38% Barricade® Herbicide

GUARANTEED ANALYSIS

Total Nitrogen (N) XXX.XX%
Available Phosphorus (K₂O) XXX.XX%
Soluble Potash (P₂O₅) XXX.XX%
Plant Nutrients Derived From:

For products being sold into states which require metals information:

Information regarding the contents and levels of metals in this product is available on the internet at:
<http://www.regulatory-info-ap.com/>.

For products containing the micronutrient iron:

Notice: this product contains the micronutrient iron to help prevent iron deficiency. Iron may stain concrete surfaces and should not be applied on water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash this product from concrete surfaces with water.

PRECAUTIONARY STATEMENTS

CAUTION

Hazards to Humans and Domestic Animals

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID

- | | |
|------------------------|---|
| If in eyes | Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. |
| If swallowed | Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| If on skin or clothing | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. |
| If inhaled | Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice. |

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

Environmental Hazards

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations substantially above the level of water solubility, it may be toxic to fish. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

3/24

(Back Panel)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For final printed labels which do include landscape ornamental uses:

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

For final printed labels which do not include landscape ornamental uses:

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

GENERAL INFORMATION

The Andersons Fertilizer with 0.38% Barricade Herbicide is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in turfgrasses and landscapes. This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

APPLICATION DIRECTIONS


Apply uniformly through suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a preemergence herbicide that, when properly applied as directed, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the **Maximum Annual Rates** section of this label.

Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

CRABGRASS GERMINATION DATES



Approx. Date

- 1 - After May 30
- 2 - After May 10
- 3 - After April 20
- 4 - After March 20
- 5 - Jan. 1 to March 20

(Back Panel)

Use Restrictions and Precautions

The following precautions apply to the use of this product in turf grasses and lawns:

- Application of this product may thin emerged annual bluegrass and newly over seeded grasses.
- Do not apply to over seeded turf within 60 days after seeding or until after the second mowing, whichever is longer.
- Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
- Do not apply to newly set sod until the following year.
- Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
- Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa annua*) are desirable species.
- Do not graze or feed livestock forage cut from areas treated with this product.
- Do not apply this product through any type of irrigation system unless otherwise instructed by this label. Do not apply aerially.
- Do not apply to golf course putting greens.

Rates of Application

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds.

This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

Maximum Annual Rates

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Annual Rates Table

| Maximum Application Rate of this product per Calendar Year by Turf Grass Species ¹ | |
|--|-------------------------------|
| Turf Species | lbs. product/ 1000 sq. ft. |
| Bermudagrass ² , Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia | 9.1 |
| Buffalograss, Kentucky Bluegrass, Perennial Ryegrass | 6.0 |
| Creeping Red Fescue | 4.5 |
| Creeping Bentgrass | 3.9 |

¹These are the maximum rates per calendar year by species limitations.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 3.0 LB./1000 sq. ft. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

Do not apply more than 9.1 lbs./1,000 sq. ft. of this product per calendar year. Use higher rates of this product to achieve higher levels of fertility and weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates table.

5/24

(Back Panel)

Weeds Controlled

When used at the recommended rates this product will control:

| | | | |
|---|-----------------------------|-------------------------------|-----------------------------------|
| Barnyardgrass | Cupgrass, Woolly | Lambsquarter, Common | Signalgrass, Broadleaf |
| Bluegrass, Annual (<i>Poa annua</i>) | Foxtails, Annual | Lovegrass | Speedwell, Persian |
| Carpetweed | Goosegrass ¹ | Oxalis | Sprangletop |
| Chickweed, Common | Henbit | Panicum | Spurge, Prostrate |
| Chickweed, Mouseear (from seed) | Itchgrass | (Texas, Fall, Browntop) | Witchgrass |
| Crabgrass | Johnsongrass (from seed) | Pigweed | Woodsorrel, Yellow (from seed) |
| (Large, Smooth) | Junglerice | Purslane, Common | |
| Crowfootgrass | Knotweed | Pusley, Florida | |
| | Kochia | Rescuegrass ³ | |
| | | Shepherd's Purse ² | |

¹ In many areas a single application of 3.9 to 9.1 lbs./1,000 sq.ft. of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 3.9 to 6.0 lbs./1,000 sq. ft. followed after 60 to 90 days by a second application at doses that would not exceed those given in the **Maximum Annual Rates Table**. Do not exceed the maximum rate for turf grass species listed in the **Maximum Annual Rates Table** above.

² Applications for this weed should be made in late summer, fall, or winter prior to germination. Do not exceed the maximum application rates specified in the Maximum Annual Rates table.

³ Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum application rates specified in the Maximum Annual Rates table. All applications must be made prior to germination of the weed seeds.

Overseeding/Reseeding

This product will inhibit the germination of turf species if overseed/reseeded too soon after application. The interval between application and overseeding/reseeding is dependent on the application rate, soil temperature, and soil moisture. A shorter interval is possible if growing conditions are warm and wet, while a longer interval is needed if conditions are cool and dry.

Use the Overseeding/Reseeding Intervals Table below as a guide when choosing the appropriate overseeding/reseeding interval for each situation.

Overseeding/Reseeding Intervals Table:

| This Product Rate (lb/1,000 sq. ft) | Overseeding/Reseeding Interval (months) Based Upon Seeding Method and Predominant Soil Conditions for the Season | | | |
|--|---|--------------|---------------|--------------|
| | Broadcast Seeding | | Drill Seeding | |
| | Wet and Warm | Cool and Dry | Wet and Warm | Cool and Dry |
| 2.8 | 4 | 6 | 3 | 4 |
| 6.0 | 8 | 12 | 6 | 8 |
| 9.1 | 12 | 18 | 9 | 12 |

SPREADER SETTINGS

{available at final printing – will vary among subregistrants and market availability of equipment}

0/24

(Back Panel)

Broadcast Spreader Settings

For one pass application: For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of wheels will ensure complete coverage.

| Spreader | 3.9 lb. / 1000sq. ft. | 4.5 lb. / 1000 sq. ft. | 6.0 lb. / 1000 sq. ft. | 9.1 lb. / 1000 sq. ft. |
|----------|--------------------------|---------------------------|---------------------------|---------------------------|
| aa | X | X | X | X |
| bb | X | X | X | X |
| cc | X | X | X | X |

Drop Spreader Settings

For one pass application: First apply 2 border strips along the edge of each end of the lawn. Then apply products along the length of the lawn, overlapping wheels slightly and shutting off the spreader before turning as you reach the edge of the border strips.

| Spreader | 3.9 lb. / 1000sq. ft. | 4.5 lb. / 1000 sq. ft. | 6.0 lb. / 1000 sq. ft. | 9.1 lb. / 1000 sq. ft. |
|----------|--------------------------|---------------------------|---------------------------|---------------------------|
| aa | X | X | X | X |
| bb | X | X | X | X |
| cc | X | X | X | X |

It is important to follow directions as indicated on this quality lawn care product as over-application may cause injury to the lawn. Measure the size of the area to be treated and do not over or under apply. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Calibrate spreader prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. Close spreader when filling, stopping or turning. Do not apply by hand or with a hand held spreader.

LANDSCAPE ORNAMENTALS

This product may be applied for residual preemergence weed control in ornamentals (including established perennial and wildflower plantings).

Use Rates

Apply this product at 3.9 to 9.1 lbs./1,000 sq. ft in fall and/or spring. Use higher rate of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 9.1lbs./1,000 sq. ft..

Application Timing and Information

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves and mulch. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

Established Landscape Ornamentals

This product may be applied to established ornamentals as a broadcast, over-the-top, or directed application. Overhead irrigation or rainfall soon after application will wash granules off plant foliage and activate the product in the soil.

7/24

(Back Panel)

Newly-Transplanted Landscape Ornamentals

This product may be applied to established ornamentals as a broadcast, over-the-top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

Tolerant Ornamental Species

This product will not harm trees, shrubs, vines, and flowers listed below. The species listed below in Table 1 are tolerant to this product. This product is also approved for application, except in California, to the species in Table 2, below. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of the ornamental species listed below.

When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product. Avoid application to plants that are under stress.

Table 1 - Tolerant Ornamental Species - All States

| Scientific name | Common name |
|----------------------------------|--|
| <i>Abies</i> spp. | Fir species** (Balsam, Fraser, Noble, etc.) |
| <i>Acer palmatum</i> | Japanese Maple |
| <i>Acer platanoides</i> | Norway Maple |
| <i>Actinidia chinensis</i> | Kiwi* |
| <i>Agapanthus africanus</i> | Lily-of-the-Nile (African Lily) |
| <i>Arctostaphylos densiflora</i> | Vine Hill Manzanita |
| <i>Arctotheca calendula</i> | Cape Weed |
| <i>Aucuba japonica</i> | Japanese Aucuba |
| <i>Berberis gladwynensis</i> | Barberry |
| <i>Berberis julianae</i> | Wintergreen Barberry |
| <i>Berberis mentorensis</i> | Mentor Barberry |
| <i>Berberis thunbergii</i> | Japanese Barberry |
| <i>Berberis verruculosa</i> | Warty Barberry |
| <i>Buxus microphylla</i> | Japanese Boxwood |
| <i>Callistemon viminalis</i> | Weeping Bottlebrush |
| <i>Calluna vulgaris</i> | Scotch Heather |
| <i>Carpobrotus edulis</i> | Hottentot Fig (Ice Plant) |
| <i>Cassia artemisioides</i> | Feathery Cassia |
| <i>Ceanothus rigidus</i> | Wild Lilac |
| <i>Chamaecyparis pisifera</i> | False Cypress |
| <i>Cleyera japonica</i> | Cleyera |
| <i>Citrus</i> spp. | Citrus species* |
| <i>Cornus florida</i> | Flowering Dogwood |
| <i>Cornus stolonifera</i> | American Dogwood |
| <i>Cortaderia selloana</i> | Pampas Grass |
| <i>Cotoneaster apiculatus</i> | Cranberry Cotoneaster |
| <i>Cotoneaster buxifolius</i> | Cotoneaster |
| <i>Cotoneaster dammeri</i> | Bearberry Cotoneaster |
| <i>Cotoneaster microphyllus</i> | Rockspray Cotoneaster |
| <i>Crataegus</i> spp. | Hawthorne |
| <i>Cupressus sempervirens</i> | Italian Cypress |
| <i>Delosperma alba</i> | White Trailing Ice Plant |
| <i>Dodonaea viscosa</i> | Hop Bush |

8/24

9198-163
Page 8 of 12
Homeowner version

(Back Panel)

| | |
|--------------------------------|--|
| <i>Elaeagnus argentea</i> | Silverberry |
| <i>Euonymus fortunei</i> | Wintercreeper |
| <i>Euonymus japonica</i> | Japanese Spindle Tree (Evergreen Euonymus) |
| <i>Euonymus alatus</i> | Spreading Euonymus |
| <i>Fatsia japonica</i> | Japanese Aralia |
| <i>Forsythia intermedia</i> | Border Forsythia |
| <i>Forsythia viridissima</i> | Greenstem Forsythia |
| <i>Gardenia jasminoides</i> | Gardenia, Cape-Jasmine |
| <i>Gladiolus</i> spp. | Gladiolus species** |
| <i>Hedera helix</i> | English Ivy |
| <i>Hibiscus</i> | Rose of Sharon** |
| <i>Hibiscus Rosa-sinensis</i> | Chinese Hibiscus |
| <i>Ilex cornuta</i> | Chinese Holly |
| <i>Ilex crenata</i> | Japanese Holly |
| <i>Ilex opaca</i> | American Holly |
| <i>Ilex pernyi</i> | Holly |
| <i>Ilex vomitoria</i> | Yaupon Holly |
| <i>Iris</i> spp. | Iris species** |
| <i>Jasminum nudiflorum</i> | Winter Jasmine |
| <i>Juniperus chinensis</i> | Chinese Juniper |
| <i>Juniperus conferta</i> | Shore Juniper |
| <i>Juniperus horizontalis</i> | Creeping Juniper |
| <i>Juglans</i> spp. | Walnut* |
| <i>Justicia brandegeana</i> | Shrimp Plant |
| <i>Lagerstromia indica</i> | Crape Myrtle |
| <i>Ligustrum amurense</i> | Amur Privet |
| <i>Ligustrum japonicum</i> | Japanese Privet |
| <i>Ligustrum lucidum</i> | Glossy Privet (Wax-Leaf) |
| <i>Liriodendron tulipifera</i> | Big Blue Lillyturf |
| <i>Lonicera japonica</i> | Japanese Honeysuckle |
| <i>Lonicera tatarica</i> | Tatarian Honeysuckle |
| <i>Magnolia</i> spp. | Magnolia species** |
| <i>Malephora luteola</i> | Ice Plant |
| <i>Malus</i> spp. | Crabapple* |
| <i>Nandina domestica</i> | Heavenly Bamboo |
| <i>Narcissus</i> spp. | Narcissus species** |
| <i>Nerium</i> spp. | Oleander |
| <i>Olea europaea</i> | Olive* |
| <i>Ophiopogon japonicus</i> | Mondo Grass |
| <i>Osteospermum fruticosum</i> | Trailing African Daisy |
| <i>Oxydendrum arboreum</i> | Sourwood |
| <i>Persea americana</i> | Avocado* |
| <i>Photinia fraseri</i> | Fraser's Photinia (Redtip) |
| <i>Picea</i> spp. | Spruce species** (Colorado Blue, Norway, etc.) |
| <i>Pieris japonica</i> | Lily-of-the-Valley Shrub |
| <i>Pinus brutia</i> | Calabrian Pine |
| <i>Pinus canariensis</i> | Canary Island Pine |
| <i>Pinus elliotii</i> | Slash Pine |
| <i>Pinus halepensis</i> | Aleppo Pine |
| <i>Pinus nigra</i> | Austrian Black Pine |
| <i>Pinus palustris</i> | Longleaf Pine |
| <i>Pinus radiata</i> | Monterey Pine |
| <i>Pinus strobus</i> | Eastern White Pine |

9/24

(Back Panel)

Pinus sylvestris
Pinus taeda
Pinus thunbergiana
Pinus virginiana
Pistacia spp.
Pittosporum rhombifolium
Pittosporum tobira
Podocarpus macrophyllus
Prunus laurocerasus
Prunus spp.

Pseudotsuga menziesii
Pyracantha coccinea
Pyracantha fortuneana
Pyracantha koidzumii
Pyrus spp.
Quercus rubra
Raphiolepis indica
Rhododendron
(including *Azalea*)

Rosa banksiae
Rosmarinus officinalis
Rumohra adiantiformis
Santolina virans
Sedum album
Syzygium paniculatum
Taxus cuspidata
Taxus media
Thuja occidentalis
Trachelospermum asiaticum
Tsuga canadensis
Tulipa spp.
Viburnum japonicum
Viburnum odoratissimum
Viburnum plicatum
Viburnum rigidum
Viburnum tinus
Viburnum trilobum
Viburnum wrightii
Vinca major
Vinca minor
Vitis spp.
Weigela florida
Yucca aloifolia
Yucca filamentosa

*Do not use on food producing trees, vines, or plants.

**Not for use on container grown plants.

Scotch Pine
Loblolly Pine
Japanese Black Pine
Virginia Pine
Pistachio*
Queensland Pittosporum
Japanese Pittosporum
Japanese Yew
English Laurel
Almond, Apricot, Nectarine,
Peach, Plum and Prune*
Douglas Fir**
Firethorn Scarlet
Firethorn
Firethorn
Bradford Pear spp.
Oak species
Indian Hawthorne
'Coral Bells'
'Formosa'
'Hino-crimson'
'PJM'
'Roseum Elegans'
Lady Bank's Rose
Rosemary*
Leatherleaf Fern

Stonecrop
Japanese Boxcherry
Japanese Yew
Yew
American Arborvitae
Star Jasmine
Canada Hemlock
Tulip species
Japanese Viburnum
Sweet Viburnum
Japanese Snowball
Canary Island Viburnum
Laurustinus
Cranberry Bush
Leatherleaf Viburnum
Vinca
Dwarf Periwinkle
Grape*
Old Fashioned Weigela
Spanish Bayonet
Yucca, Adam's Needle

(Back Panel)

Table 2 - Tolerant Ornamental Species/Varieties - All States Except CA

| Scientific name | Common name |
|---|---|
| <i>Abelia grandiflora</i> | Abelia: Sherwood |
| <i>Achillea</i> spp. | Yarrow: King Edward |
| <i>Agapanthus orientalis</i> | |
| <i>Akebia quinata</i> | Five-Leaf or Chocolate Vine |
| <i>Allium cernuum</i> | Lady's Leek, Nodding Onion |
| <i>Anemone hybrida</i> | Japanese Anemone |
| <i>Aquilegia</i> spp. | Aquilegia: Red and Gold |
| <i>Artemisia</i> spp. | Wormwood; Silver Mound, Castle |
| <i>Aster</i> spp. | Aster: Bonny Blue, Purple Dome |
| <i>Aster X frikartii</i> | |
| <i>Athyrium filix-femina</i> | Lady Fern; Fern Lady |
| <i>Begonia</i> spp. | Fibrous Begonia: Hardy Grandis |
| <i>Bergenia cordifolia</i> | |
| <i>Boltonia asteroides</i> | Snowbank |
| <i>Bougainvillea</i> spp. | Bougainvillea |
| <i>Buddleia davidii</i> | Butterfly-Bush (Dwarf Blue); Royal Red |
| <i>Callistemon citrinus</i> | Crimson Bottlebrush |
| <i>Campanula carpatica</i> | Tussock Bellflower; (White Clips) |
| <i>Campis X tagliabuana</i> | Trumpet Creeper, Trumpet Flower; Madame Galen |
| <i>Ceratostigma plumbaginoides</i> | |
| <i>Chrysanthemum nipponicum</i> | |
| <i>Coreopsis</i> spp. | Coreopsis (Calliopsis): Early Sunrise, Moonbeam |
| <i>Crocasmia</i> spp. | Lucifer |
| <i>Delosperma</i> spp. | Cooperi Pink |
| <i>Delphinium</i> spp. | Larkspur; Blue Elf |
| <i>Dianthus deltoides</i> | Dianthus, Maiden Pinks 'Zing' |
| <i>Dianthus gratianopolitanus</i> | Cheddar Pink |
| <i>Echinacea pupurea</i> | Coneflower, Purple; Magnus |
| <i>Forsythia suspensa</i> | Weeping Forsythia |
| <i>Gaillardia</i> spp. | Gaillardia, Blanket Flower: 'Goblin' |
| <i>Gaura</i> spp. | |
| <i>Gentiana dahurica</i> | Gentian |
| <i>Geranium cinereum</i> | Cranesbill |
| <i>Gypsophila repens</i> | Baby's Breath |
| <i>Helianthemum</i> spp. | Sunrose |
| <i>Hemerocallis</i> spp. | Daylily: Aztec Gold, Stella De Oro, Tender Love |
| <i>Heucherella</i> spp. | Coral Bell; Bridget Bloom |
| <i>Hibiscus</i> spp. | Mallow; Disco Belle White |
| <i>Hosta plantaginea</i> | Hosta, Plantain Lily (Fragrant) |
| <i>Hosta sieboldiana</i> | Hosta, 'Searsucker' |
| <i>Houttuynia cordata</i> var. <i>variegata</i> | |
| <i>Hydrangea macrophylla</i> | Bigleaf Hydrangea |
| <i>Inula ensifolia</i> | |
| <i>Iris ensata</i> | Sword-Leaved Iris; Jodlesong |
| <i>Iris siberica</i> | Siberian Iris; Cabernet |
| <i>Juniperus davurica</i> | Parsoni |
| <i>Lagerstromia indica x fauriei</i> | Crape Myrtle; Tuscarora |
| <i>Lantana montevidensis</i> | Weeping Lantana |
| <i>Lavender</i> spp. | Lavender; Munstead |

(Back Panel)

| | |
|--|------------------------------------|
| <i>Leontopodium alpinum</i> | Edelweiss |
| <i>Ligustrum sinense</i> | Chinese Privet; Variegata |
| <i>Lilium</i> spp. | Lily; Jazz |
| <i>Liriope muscari</i> var. <i>variegata</i> | Liriope, Variegated |
| <i>Liriope spicata</i> | Liriope, Creeping |
| <i>Lobelia cardinalis</i> | Cardinal Flower, Indian Pink |
| <i>Loropetalum chinense</i> | Burgundy |
| <i>Lythrum</i> spp. | Loosestrife; Modern Pink |
| <i>Miscanthus sinensis</i> | Yaku Jima, Silberfeder |
| <i>Oenothera missouriensis</i> | Evening Primrose |
| <i>Osmanthus heterophyllus</i> | Osmanthus (False Holly): Gulf Tide |
| <i>Paeonia suffruticosa</i> | Tree Peony |
| <i>Pennisetum setaceum</i> | Fountain Grass (Dwarf) |
| <i>Perovskia atriplicifolia</i> | |
| <i>Physostegia virginiana</i> | Dragonhead, False; Vivid |
| <i>Quercus shumardii</i> | Oak, Shumard's Red |
| <i>Raphiolepis umbellata</i> | Yedda Hawthorne |
| <i>Rhododendron</i> (including azalea) | 'Delaware Valley White' |
| | 'Flame Creeper' |
| | 'Girard Crimson' |
| | 'George L. Tabor' |
| | Wakeiebisu |
| | White Gumpo |
| <i>Rudbeckia</i> spp. | Black-Eyed Susan; Goldstrum |
| <i>Saxifraga</i> spp. | Saxifrage; Purple Dome |
| <i>Scabiosa</i> spp. | Pincushion Flower |
| <i>Sedum caudicola</i> | Stonecrop; Lidakense |
| <i>Sedum dasyphyllum</i> | Stonecrop |
| <i>Sedum spurium</i> | Stonecrop; Dragon's Blood |
| <i>Spiraea bumalda</i> | Spirea; Anthony Waterer |
| <i>Syzygium paniculatum</i> | Australian Brushcherry |
| <i>Teucrium</i> spp. | Germander |
| <i>Thalictrum dipterocarpum</i> | Meadow Rue |
| <i>Veronica</i> spp. | Veronica, Speedwell; Sunny Border |
| <i>Viburnum suspensum</i> | Arrowwood Viburnum |

STORAGE AND DISPOSAL*For homeowner products:*

Do not contaminate water, food, or feed by storage or disposal.

Storage:

Store in original container away from fertilizer, feed, or food stuffs and separated from other pesticides.

Disposal:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

12/24

(Back Panel)

IMPORTANT! READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (RVamd030804)

Manufactured by:

The Andersons Lawn Fertilizer Division, Inc.

d/b/a Free Flow Fertilizer

PO Box 119

Maumee, OH 43537

| | |
|---------------|--|
| Date Printed | 3/8/04 |
| Date Approved | |
| Supersedes | 11/5/97 |
| Label changes | Amendment – to match Syngenta's basic label (Barricade F & 65WG) |
| Filename | 9198-163 |
| Label code | RVamd030804 |

13/24

(Front Panel)

The Andersons Fertilizer with 0.38% Barricade® Herbicide

| | |
|--------------------------|---------------|
| Active Ingredient: | |
| Prodiamine* | 0.38% |
| Other Ingredients: | <u>99.62%</u> |
| Total | 100.00% |

*(CAS No. 29091-21-2)

Barricade – trademark of Syngenta Crop Protection

For preemergence control of grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals
- established perennial & wildflower plantings

KEEP OUT OF REACH OF CHILDREN


CAUTION

See back panel for additional Precautionary Statements and First Aid

EPA Reg. No. 9198-163
EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A
Underlined letter is first letter used in run code

Net Weight
Covers

ACCEPTED
APR 29 2004
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
9198-163



14/24

(Back Panel)

The Andersons Fertilizer with 0.38% Barricade® Herbicide

GUARANTEED ANALYSIS

Total Nitrogen (N) XXX.XX%
Available Phosphorus (K₂O) XXX.XX%
Soluble Potash (P₂O₅) XXX.XX%
Plant Nutrients Derived From:

For products being sold into states which require metals information:

Information regarding the contents and levels of metals in this product is available on the internet at:
<http://www.regulatory-info-ap.com/>.

For products containing the micronutrient iron:

Notice: this product contains the micronutrient iron to help prevent iron deficiency. Iron may stain concrete surfaces and should not be applied on water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash this product from concrete surfaces with water.

PRECAUTIONARY STATEMENTS

CAUTION

Hazards to Humans and Domestic Animals

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID

- | | |
|------------------------|---|
| If in eyes | Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. |
| If swallowed | Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| If on skin or clothing | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. |
| If inhaled | Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice. |

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

Environmental Hazards

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations substantially above the level of water solubility, it may be toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

15/24

(Back Panel)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For final printed labels which do include landscape ornamental uses:

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

For final printed labels which do not include landscape ornamental uses:

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

GENERAL INFORMATION

The Andersons Fertilizer with 0.38% Barricade Herbicide is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in turfgrasses and landscapes. This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

APPLICATION DIRECTIONS


Apply uniformly through suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a preemergence herbicide that, when properly applied as directed, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the **Maximum Annual Rates** section of this label.

Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

CRABGRASS GERMINATION DATES



Approx. Date

- 1 - After May 30
- 2 - After May 10
- 3 - After April 20
- 4 - After March 20
- 5 - Jan. 1 to March 20

(Back Panel)

Use Restrictions and Precautions

The following precautions apply to the use of this product in turf grasses and lawns:

- Application of this product may thin emerged annual bluegrass and newly over seeded grasses.
- Do not apply to over seeded turf within 60 days after seeding or until after the second mowing, whichever is longer.
- Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
- Do not apply to newly set sod until the following year.
- Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
- Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa annua*) are desirable species.
- Do not graze or feed livestock forage cut from areas treated with this product.
- Do not apply this product through any type of irrigation system unless otherwise instructed by this label.
- Do not apply aerially.
- Do not apply to golf course putting greens.

Rates of Application

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

Maximum Annual Rates

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Annual Rates Table

| Maximum Application Rate of this product per Calendar Year by Turf Grass Species¹ | | | |
|--|------------------|------------------------|-----------------------------------|
| Turf Species | lbs. AI/A | lbs. product /A | lbs. product/ 1000 sq. ft. |
| Bermudagrass ² , Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia | 1.5 | 395 | 9.1 |
| Buffalograss, Kentucky Bluegrass, Perennial Ryegrass | 1.0 | 263 | 6.0 |
| Creeping Red Fescue | 0.75 | 197 | 4.5 |
| Creeping Bentgrass | 0.65 | 171 | 3.9 |

¹These are the maximum rates per calendar year by species limitations.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 132 lb./acre or 3.0 LB./1000 sq. ft. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

Do not apply more than 395 lbs. per acre (9.1 lbs./1,000 sq. ft.) of this product per calendar year. Use higher rates of this product to achieve higher levels of fertility and weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates table.

(Back Panel)

Weeds Controlled

When used at the recommended rates this product will control:

| | | | |
|---|-------------------------|-------------------------------|-----------------------------------|
| Barnyardgrass | Cupgrass, Woolly | Lambsquarter, Common | Signalgrass, Broadleaf |
| Bluegrass, Annual (<i>Poa annua</i>) | Foxtails, Annual | Lovegrass | Speedwell, Persian |
| Carpetweed | Goosegrass ¹ | Oxalis | Sprangletop |
| Chickweed, Common | Henbit | Panicum | Spurge, Prostrate |
| Chickweed, Mouseear (from seed) | Itchgrass | (Texas, Fall, Browntop) | Witchgrass |
| Crabgrass | Johnsongrass | Pigweed | Woodsorrel, Yellow (from seed) |
| (Large, Smooth) | (from seed) | Purslane, Common | |
| Crowfootgrass | Junglerice | Pusley, Florida | |
| | Knotweed | Rescuegrass ³ | |
| | Kochia | Shepherd's Purse ² | |

¹ In many areas a single application of 3.9 to 9.1 lbs./1,000 sq.ft. (171 to 395 lbs./A) of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 3.9 to 6.0 lbs./1,000 sq. ft. (171 to 263 lbs./A) followed after 60 to 90 days by a second application at doses that would not exceed those given in the **Maximum Annual Rates Table**. Do not exceed the maximum rate for turf grass species listed in the **Maximum Annual Rates Table** above.

² Applications for this weed should be made in late summer, fall, or winter prior to germination. Do not exceed the maximum application rates specified in the Maximum Annual Rates table.

³ Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum application rates specified in the Maximum Annual Rates table. All applications must be made prior to germination of the weed seeds.

When to Apply After Overseeding Turf

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

When to Overseed After Application – All States*

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

| The Andersons Fertilizer with 0.38% Barricade Herbicide Rate | | | Interval Before Overseeding | | |
|---|--------|------|-----------------------------|------------|-------|
| lbs/1000 sq. ft | lbs./A | ai/A | North | Transition | South |
| 3.0 | 131.5 | 0.50 | 4 mo | 4 mo | 4 mo |
| 3.9 | 171 | 0.65 | 5 mo | 4 mo | 4 mo |
| 4.5 | 197 | 0.75 | 6 mo | 5 mo | 5 mo |
| 4.9 | 213 | 0.81 | --- | 6 mo | 6 mo |
| 5.9 | 258 | 1.00 | --- | 7 mo | 7 mo |
| 6.9 | 302 | 1.15 | --- | --- | 9 mo |
| 7.8 | 342 | 1.30 | --- | --- | 10 mo |
| 9.1 | 395 | 1.50 | --- | --- | 12 mo |

18/24

(Back Panel)

*Note: In AZ, CA, NV and TX, the overseeding interval can be shorter in established bermudagrass that has been overseeded with perennial ryegrass. See the next section " *Poa annua* Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ, CA, NX and TX only)".

***Poa annua* Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ, CA, NX and TX only)**

Use on golf courses (excluding golf course putting greens) and lawns when overseeding with perennial ryegrass (minimum seed rate of 350 lbs./A).

How Much and When to Apply

| Amount to Apply | | When to Apply | Expected Control | Use Precautions |
|-----------------|-----------|---|--|--|
| Lbs./Acre | Lbs./1000 | | | |
| 99 – 171 | 2.3 – 3.9 | 6 to 8 weeks before ryegrass overseeding Second application 4-8 weeks after overseeding date or when perennial ryegrass roots are in the second inch of soil | One application for 70% or greater control of <i>Poa annua</i> Second application should enhance the residual under normal weather conditions | 1. Some seed mortality and temporary reduction in root growth of new seedlings may occur 2. To reduce the potential for seeding mortality, maintain a moist seedbed with light, frequent irrigation. 3. Make no more than 2 applications per year for this use, and do not exceed 222 lbs./A (5.1 lbs./1000) for these two applications for <i>Poa annua</i> control in ryegrass overseeded turf 4. Do not make a second application if any injury to the ryegrass is observed after the first application 5. Do not make a second application unless the product was first applied before overseeding |

Recommended Spreader Settings

These suggested spreader settings are not intended to replace calibration.
Please calibrate your spreader before applying product.

___ lbs. treats ___ ft² at the 171 lbs. product/acre – LIGHT RATE (3.9 lbs./1,000 sq. ft.)
___ lbs. treats ___ ft² at the 197 lbs. product/acre – LOW RATE (4.5 lbs./1,000 sq. ft.)
___ lbs. treats ___ ft² at the 263 lbs. product/acre – MEDIUM RATE (6.0 lbs./1,000 sq. ft.)
___ lbs. treats ___ ft² at the 395 lbs. product/acre – HIGH RATE (9.1 lbs./1,000 sq. ft.)

| Spreader | Ground or PTO Speed | Width of Coverage | Spreader Settings | | | |
|----------|---------------------|-------------------|-------------------|----------|-------------|-----------|
| | | | Light Rate | Low Rate | Medium Rate | High Rate |
| aa | X mph | XX ft/ | X | X | X | X |
| bb | X mph | XX ft | X | X | X | X |
| cc | X mph | XX ft | X | X | X | X |

19/24

(Back Panel)

LANDSCAPE ORNAMENTALS

This product may be applied for residual preemergence weed control in ornamentals (including established perennial and wildflower plantings).

Use Rates

Apply this product at 171 to 395 lbs./A (3.9 to 9.1 lbs./1,000 sq. ft) in fall and/or spring. Use higher rate of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 395 lbs./A (9.1 lbs./1,000 sq. ft.).

Application Timing and Information

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves and mulch. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

Established Landscape Ornamentals

This product may be applied to established ornamentals as a broadcast, over-the-top, or directed application. Overhead irrigation or rainfall soon after application will wash granules off plant foliage and activate the product in the soil.

Newly-Transplanted Landscape Ornamentals

This product may be applied to established ornamentals as a broadcast, over-the-top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

Tolerant Ornamental Species

This product will not harm trees, shrubs, vines, and flowers listed below. The species listed below in Table 1 are tolerant to this product. This product is also approved for application, except in California, to the species in Table 2, below. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of the ornamental species listed below.

When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product. Avoid application to plants that are under stress.

Table 1 - Tolerant Ornamental Species - All States

| Scientific name | Common name |
|----------------------------------|--|
| <i>Abies</i> spp. | Fir species** (Balsam, Fraser, Noble, etc.) |
| <i>Acer palmatum</i> | Japanese Maple |
| <i>Acer platanoides</i> | Norway Maple |
| <i>Actinidia chinensis</i> | Kiwi* |
| <i>Agapanthus africanus</i> | Lily-of-the-Nile (African Lily) |
| <i>Arctostaphylos densiflora</i> | Vine Hill Manzanita |
| <i>Arctotheca calendula</i> | Cape Weed |
| <i>Aucuba japonica</i> | Japanese Aucuba |
| <i>Berberis gladiolensis</i> | Barberry |

(Back Panel)

| | |
|---------------------------------|--|
| <i>Berberis julianae</i> | Wintergreen Barberry |
| <i>Berberis mentorensis</i> | Mentor Barberry |
| <i>Berberis thunbergii</i> | Japanese Barberry |
| <i>Berberis verruculosa</i> | Warty Barberry |
| <i>Buxus microphylla</i> | Japanese Boxwood |
| <i>Callistemon viminalis</i> | Weeping Bottlebrush |
| <i>Calluna vulgaris</i> | Scotch Heather |
| <i>Carpobrotus edulis</i> | Hottentot Fig (Ice Plant) |
| <i>Cassia artemisioides</i> | Feathery Cassia |
| <i>Ceanothus rigidus</i> | Wild Lilac |
| <i>Chamaecyparis pisifera</i> | False Cypress |
| <i>Cleyera japonica</i> | Cleyera |
| <i>Citrus</i> spp. | Citrus species* |
| <i>Cornus florida</i> | Flowering Dogwood |
| <i>Cornus stolonifera</i> | American Dogwood |
| <i>Cortaderia selloana</i> | Pampas Grass |
| <i>Cotoneaster apiculatus</i> | Cranberry Cotoneaster |
| <i>Cotoneaster buxifolius</i> | Cotoneaster |
| <i>Cotoneaster dammeri</i> | Bearberry Cotoneaster |
| <i>Cotoneaster microphyllus</i> | Rockspray Cotoneaster |
| <i>Crataegus</i> spp. | Hawthorne |
| <i>Cupressus sempervirens</i> | Italian Cypress |
| <i>Delosperma alba</i> | White Trailing Ice Plant |
| <i>Dodonaea viscosa</i> | Hop Bush |
| <i>Elaeagnus pungens</i> | Silverberry |
| <i>Euonymus fortunei</i> | Wintercreeper |
| <i>Euonymus japonica</i> | Japanese Spindle Tree (Evergreen Euonymus) |
| <i>Euonymus kiautschovicks</i> | Spreading Euonymus |
| <i>Fatsia japonica</i> | Japanese Aralia |
| <i>Forsythia intermedia</i> | Border Forsythia |
| <i>Forsythia viridissima</i> | Greenstem Forsythia |
| <i>Gardenia jasminoides</i> | Gardenia, Cape-Jasmine |
| <i>Gladiolus</i> spp. | Gladiolus species** |
| <i>Hedera helix</i> | English Ivy |
| <i>Hibiscus</i> | Rose of Sharon** |
| <i>Hibiscus Rosa-sinensis</i> | Chinese Hibiscus |
| <i>Ilex cornuta</i> | Chinese Holly |
| <i>Ilex crenata</i> | Japanese Holly |
| <i>Ilex opaca</i> | American Holly |
| <i>Ilex pernyi</i> | Holly |
| <i>Ilex vomitoria</i> | Yaupon Holly |
| <i>Iris</i> spp. | Iris species** |
| <i>Jasminum nudiflorum</i> | Winter Jasmine |
| <i>Juniperus chinensis</i> | Chinese Juniper |
| <i>Juniperus conferta</i> | Shore Juniper |
| <i>Juniperus horizontalis</i> | Creeping Juniper |
| <i>Juglans</i> spp. | Walnut* |
| <i>Justicia brandegeana</i> | Shrimp Plant |
| <i>Lagerstromia indica</i> | Crape Myrtle |
| <i>Ligustrum amurense</i> | Amur Privet |
| <i>Ligustrum japonicum</i> | Japanese Privet |
| <i>Ligustrum lucidum</i> | Glossy Privet (Wax-Leaf) |
| <i>Liriope muscari</i> | Big Blue Lillyturf |

(Back Panel)

| | |
|---|---|
| <i>Lonicera japonica</i> | Japanese Honeysuckle |
| <i>Lonicera tatarica</i> | Tatarian Honeysuckle |
| <i>Magnolia</i> spp. | Magnolia species** |
| <i>Malephora luteola</i> | Ice Plant |
| <i>Malus</i> spp. | Crabapple* |
| <i>Nandina domestica</i> | Heavenly Bamboo |
| <i>Narcissus</i> spp. | Narcissus species** |
| <i>Nerium</i> spp. | Oleander |
| <i>Olea europaea</i> | Olive* |
| <i>Ophiopogon japonicus</i> | Mondo Grass |
| <i>Osteospermum fruticosum</i> | Trailing African Daisy |
| <i>Oxydendrum arboreum</i> | Sourwood |
| <i>Persea americana</i> | Avocado* |
| <i>Photinia fraseri</i> | Frasier's Photinia (Redtip) |
| <i>Picea</i> spp. | Spruce species** (Colorado Blue, Norway, etc.) |
| <i>Pieris japonica</i> | Lily-of-the-Valley Shrub |
| <i>Pinus brutia</i> | Calabrian Pine |
| <i>Pinus canariensis</i> | Canary Island Pine |
| <i>Pinus elliotii</i> | Slash Pine |
| <i>Pinus halepensis</i> | Aleppo Pine |
| <i>Pinus nigra</i> | Austrian Black Pine |
| <i>Pinus palustris</i> | Longleaf Pine |
| <i>Pinus radiata</i> | Monterey Pine |
| <i>Pinus strobus</i> | Eastern White Pine |
| <i>Pinus sylvestris</i> | Scotch Pine |
| <i>Pinus taeda</i> | Loblolly Pine |
| <i>Pinus thunbergiana</i> | Japanese Black Pine |
| <i>Pinus virginiana</i> | Virginia Pine |
| <i>Pistacia</i> spp. | Pistachio* |
| <i>Pittosporum rhombifolium</i> | Queensland Pittosporum |
| <i>Pittosporum tobira</i> | Japanese Pittosporum |
| <i>Podocarpus macrophyllus</i> | Japanese Yew |
| <i>Prunus laurocerasus</i> | English Laurel |
| <i>Prunus</i> spp. | Almond, Apricot, Nectarine, Peach, Plum and Prune* |
| <i>Pseudotsuga menziesii</i> | Douglas Fir** |
| <i>Pyracantha coccinea</i> | Firethorn Scarlet |
| <i>Pyracantha fortuneana</i> | Firethorn |
| <i>Pyracantha koidzumii</i> | Firethorn |
| <i>Pyrus</i> spp. | Bradford Pear spp. |
| <i>Quercus rubra</i> | Oak species |
| <i>Raphiolepis indica</i> | Indian Hawthorne |
| <i>Rhododendron</i> (including Azalea) | 'Coral Bells' 'Formosa' 'Hino-crimson' 'PJM' 'Roseum Elegans' |
| <i>Rosa banksiae</i> | Lady Bank's Rose |
| <i>Rosmarinus officinalis</i> | Rosemary* |
| <i>Rumohra adiantiformis</i> | Leatherleaf Fern |
| <i>Santolina virens</i> | |
| <i>Sedum album</i> | Stonecrop |
| <i>Syzygium paniculatum</i> | Japanese Boxcherry |

22/24

(Back Panel)

| | |
|----------------------------------|------------------------|
| <i>Taxus cuspidata</i> | Japanese Yew |
| <i>Taxus media</i> | Yew |
| <i>Thuja occidentalis</i> | American Arborvitae |
| <i>Trachelospermum asiaticum</i> | Star Jasmine |
| <i>Tsuga canadensis</i> | Canada Hemlock |
| <i>Tulipa</i> spp. | Tulip species |
| <i>Viburnum japonicum</i> | Japanese Viburnum |
| <i>Viburnum odoratissimum</i> | Sweet Viburnum |
| <i>Viburnum plicatum</i> | Japanese Snowball |
| <i>Viburnum rigidum</i> | Canary Island Viburnum |
| <i>Viburnum tinus</i> | Laurustinus |
| <i>Viburnum trilobum</i> | Cranberry Bush |
| <i>Viburnum wrightii</i> | Leatherleaf Viburnum |
| <i>Vinca major</i> | Vinca |
| <i>Vinca minor</i> | Dwarf Periwinkle |
| <i>Vitis</i> spp. | Grape* |
| <i>Weigela florida</i> | Old Fashioned Weigela |
| <i>Yucca aloifolia</i> | Spanish Bayonet |
| <i>Yucca filamentosa</i> | Yucca, Adam's Needle |

*Do not use on food producing trees, vines, or plants.

**Not for use on container grown plants.

Table 2 - Tolerant Ornamental Species/Varieties - All States Except CA

| Scientific name | Common name |
|------------------------------------|---|
| <i>Abelia grandiflora</i> | Abelia: Sherwood |
| <i>Achillea</i> spp. | Yarrow: King Edward |
| <i>Agapanthus orientalis</i> | |
| <i>Akebia quinata</i> | Five-Leaf or Chocolate Vine |
| <i>Allium cernuum</i> | Lady's Leek, Nodding Onion |
| <i>Anemone hybrida</i> | Japanese Anemone |
| <i>Aquilegia</i> spp. | Aquilegia: Red and Gold |
| <i>Artemisia</i> spp. | Wormwood; Silver Mound, Castle |
| <i>Aster</i> spp. | Aster: Bonny Blue, Purple Dome |
| <i>Aster X frikartii</i> | |
| <i>Athyrium filix-femina</i> | Lady Fern; Fern Lady |
| <i>Begonia</i> spp. | Fibrous Begonia: Hardy Grandis |
| <i>Bergenia cordifolia</i> | |
| <i>Boltonia asteroides</i> | Snowbank |
| <i>Bougainvillea</i> spp. | Bougainvillea |
| <i>Buddleia davidii</i> | Butterfly-Bush (Dwarf Blue); Royal Red |
| <i>Callistemon citrinus</i> | Crimson Bottlebrush |
| <i>Campanula carpatica</i> | Tussock Bellflower; (White Clips) |
| <i>Campis X tagliabuana</i> | Trumpet Creeper, Trumpet Flower; Madame Galen |
| <i>Ceratostigma plumbaginoides</i> | |
| <i>Chrysanthemum nipponicum</i> | |
| <i>Coreopsis</i> spp. | Coreopsis (Calliopsis): Early Sunrise, Moonbeam |
| <i>Crocosmia</i> spp. | Lucifer |
| <i>Delosperma</i> spp. | Cooperi Pink |
| <i>Delphinium</i> spp. | Larkspur; Blue Elf |
| <i>Dianthus deltoides</i> | Dianthus, Maiden Pinks 'Zing' |
| <i>Dianthus gratianopolitanus</i> | Cheddar Pink |

(Back Panel)

Echinacea purpurea
Forsythia suspensa
Gaillardia spp.
Gaura spp.
Gentiana dahurica
Geranium cinereum
Gypsophila repens
Helianthemum spp.
Hemerocallis spp.
Heucherella spp.
Hibiscus spp.
Hosta plantaginea
Hosta sieboldiana
Houttuynia cordata var. *variegata*
Hydrangea macrophylla
Inula ensifolia
Iris ensata
Iris siberica
Juniperus davurica
Lagerstromia indica x *fauriei*
Lantana montevidensis
Lavender spp.
Leontopodium alpinum
Ligustrum sinense
Lilium spp.
Liriodendron muscari var. *variegata*
Liriodendron spicata
Lobelia cardinalis
Loropetalum chinense
Lythrum spp.
Miscanthus sinensis
Oenothera missouriensis
Osmanthus heterophyllus
Paeonia suffruticosa
Pennisetum setaceum
Perovskia atriplicifolia
Physostegia virginiana
Quercus shumardii
Rhaphiolepis umbellata
Rhododendron
 (including azalea)

Rudbeckia spp.
Saxifraga spp.
Scabiosa spp.
Sedum cauticola
Sedum dasyphyllum
Sedum spurium
Spiraea bumalda
Syzygium paniculatum

Coneflower, Purple; Magnus
 Weeping Forsythia
 Gaillardia, Blanket Flower: 'Goblin'

 Gentian
 Cranesbill
 Baby's Breath
 Sunrose
 Daylily: Aztec Gold, Stella De Oro, Tender Love
 Coral Bell; Bridget Bloom
 Mallow; Disco Belle White
 Hosta, Plantain Lily (Fragrant)
 Hosta, 'Searsucker'

Bigleaf Hydrangea

Sword-Leaved Iris; Jodlesong
 Siberian Iris; Cabernet
 Parsoni
 Crape Myrtle; Tuscarora
 Weeping Lantana
 Lavender; Munstead
 Edelweiss
 Chinese Privet; Variegata
 Lily: Jazz
 Liriodendron, Variegated
 Liriodendron, Creeping
 Cardinal Flower, Indian Pink
 Burgundy
 Loosestrife; Modern Pink
 Yaku Jima, Silberfeder
 Evening Primrose
 Osmanthus (False Holly): Gulf Tide
 Tree Peony
 Fountain Grass (Dwarf)

Dragonhead, False; Vivid
 Oak, Shumard's Red
 Yedda Hawthorne
 'Delaware Valley White'
 'Flame Creeper'
 'Girard Crimson'
 'George L. Tabor'
 Wakeiebisu
 White Gumpo
 Black-Eyed Susan: Goldstrum
 Saxifrage; Purple Dome
 Pincushion Flower
 Stonecrop; Lidakense
 Stonecrop
 Stonecrop; Dragon's Blood
 Spirea: Anthony Waterer
 Australian Brushcherry

(Back Panel)

Teucrium spp.
Thalictrum dipterocarpum
Veronica spp.
Viburnum suspensum

Germander
 Meadow Rue
 Veronica, Speedwell; Sunny Border
 Arrowwood Viburnum

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container away from fertilizer, feed, or food stuffs and separated from other pesticides.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty container into application equipment. Then dispose of empty bag or box in a sanitary landfill or at an incineration facility; or, if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (RVamd030804)

Manufactured by:

The Andersons Lawn Fertilizer Division, Inc.
 d/b/a Free Flow Fertilizer
 PO Box 119
 Maumee, OH 43537

| | |
|---------------|--|
| Date Printed | 3/8/04 |
| Date Approved | |
| Supersedes | 11/5/97 |
| Label changes | Amendment – to match Syngenta's basic label (Barricade F & 65WG) |
| Filename | 9198-163 |
| Label code | RVamd030804 |