

# ProTurf Fluid Fungicide III

Prevents and controls turf diseases on tees, greens, fairways and other turf areas

Net contents 1 Gallon (3.78 liters)

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

**Hazards to Humans and Domestic Animals:** CAUTION, causes eye irritation. Avoid contact with skin, eyes, or on clothing. Wash thoroughly with soap and water after handling. Harmful if swallowed. Consumption of alcoholic beverages increases the toxic effect of thiram.

### FIRST AID

- If in eyes            Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
- If swallowed        Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
- If on skin or clothing    Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
- If inhaled            Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.


**Environmental Hazards:** Do not apply within twenty-five (25) feet of any body of water where fish are taken for human consumption. This pesticide product is toxic to fish and wildlife. Do not contaminate water when disposing of equipment washwaters. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark.

### ACTIVE INGREDIENTS:

Triadimefon, 1-(4-Chlorophenoxy)-3,3-dimethyl-1H-1,2,4-triazol-1-yl)-2-butanone .....	1.59%
Thiram, Tetramethylthiuram disulfate .....	40.76%
INERT INGREDIENTS: .....	57.65%
Total	100.00%

EPA Reg. No. 9198-197  
EPA Est. No. 37429-GA-01  
Product of USA

Recommended for Use by Professional Turfgrass Managers



2 3 4

Back panel:

**DIRECTIONS FOR USE**

Read entire label and use only as directed.

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Prevents and controls the following diseases on the turfs where they occur:

- Leaf spot/melting out (*Helminthosporium* spp./*Dreschsler* spp.)
- Dollar spot (including benzimidazole tolerant strains)
- Brown patch
- Pink snow mold/*Fusarium* patch (*microdochium nivale*, Gerlachia patch) (Washington, Oregon, California only)
- Anthracnose (on *Poa annua*)
- Rust

**Mixing Instructions**

Shake product container well before pouring into spray tank.

Add ProTurf Fluid Fungicide III into the spray tank during filling, while maintaining agitation.

Use with 4-7 gallons of water per 1,000 sq. ft. for tees and greens. When temperatures are above 90°F and humidity is above 70%, apply 7 gallons per 1,000 sq. ft. On fairways and other turf areas, use 4 gallon or higher rate of water.

Thoroughly rinse container when empty and pour remaining contents into spray tank.

Do not combine ProTurf Fluid Fungicide III in the spray tank with surfactants or other adjuvants.

**Application Directions**

Apply with standard pressurized spray equipment.

**To Prevent Diseases** – Treat turf at the NORMAL RATE when weather conditions are favorable for the development of disease. Repeat on a 10-21 day schedule.

**To control Mild Disease Infestations** – Use the NORMAL RATE. Repeat the NORMAL RATE on a 10-14 day schedule until disease is brought under control, then treat at the NORMAL RATE on a preventative schedule.

**To control Moderate to Severe Disease Infestations** –Use the HEAVY RATE. Repeat on a 10-14 day schedule until disease is brought under control, then treat at the NORMAL RATE on a preventative schedule.

Coverage Chart for the Labeled Rates of ProTurf Fluid Fungicide III			
Rate (fl. oz./M)	1 Quart (32 fl. oz.) Will cover	½ Gallon (64 fl. oz.) Will cover	1 Gallon (128 fl. oz.) Will cover
NORMAL (6 fl. oz./M)	5,500 sq. ft.	11,000 sq. ft.	22,000 sq. ft.
HEAVY (9 fl. oz./M)	4,125 sq. ft.	8,250 sq. ft.	16,500 sq. ft.
Use with 4-7 gallons of water per 1,000 sq. ft. for tees and greens.			

3 8 4.

LIMITATIONS: *Not* a control for Pythium, stripe smut, spring dead spot, gray snow mold or Fusarium Blight.

**Note to user**

*Do not* allow spray mixture to stand in the tank for longer than 12 hours.

Spray mixture should be agitated continuously during application.

Use a properly calibrated sprayer.

For best results, *do not* water or mow treated areas within 24 hours of application.

Except in cases of severe infestation, allow one week between applications of this and other control products.

Under some environmental conditions, slight leaf tip bleaching will occur from use of this product.

*Do not* apply this product through any type of irrigation system.

**Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, clean, dry place. Do not store at temperatures exceeding 120°F nor at temperatures below 32°F.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Do not reuse empty container.

**IMPORTANT: READ BEFORE USE:**

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

**WARRANTY DISCLAIMER and LIMITATION of LIABILITY:**

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

**LEGAL RIGHTS:**

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (Rvamd092703)

Sold by:

The Andersons Lawn Fertilizer Division, Inc.

PO Box 119

Maumee, OH 43537

---

Date Printed	9/26/03
Date Approved	
Supersedes	Transfer
Label Changes	Amendment - first aid
Filename	9198-197
Label Code	Rvamd092703