9198-20

12-20-2007

5.804510



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, DC 20460 Use cancellation order publisher S/14/06 71FR47213-47215

> OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Dawn L. Walters The Andersons Lawn Fertilizer Division, Inc. P.O. Box 119 Maumee, OH 43537

DEC 2 0 2007

Subject: Andersons[™] Golf Products Fungicide IX EPA Reg. No. 9198-204 Amendment dated January 19, 2007 EPA Decision Number 377271

Dear Ms Walters:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable provided the following change is made:

- 1. In the column "Use sites" in the table "Maximum Use Rates" on page 3 change the block "Residential and public turf areas: home lawns, parks, athletic fields, schools, day care centers" to "Athletic fields used only by professional athletes"
- 2. In the section "WARRANTY DISCLAIMER and LIMITATION of LIABILIBY:"
 - a. Change the sentence "Where permissible, it is Manufacturer's intent to limit any liability for...." to "To the extent consistent with applicable law, manufacturer's liability for special, consequential or incidental economic damages shall be limited to refund of purchase price or replacement of product at Buyer's choice."
 - b. Change "Manufacturer disclaims any liability...." To "To the extent consistent with applicable law, manufacturer disclaims any liability...."

One copy of the label stamped "Accepted with comments" is enclosed for your records. This label supersedes all labels previously accepted for this product. Please submit one copy of the final printed label that incorporates the required changes before the product is released for shipment.

If you have any questions, please contact Robert Westin by phone at (703) 305-5721 or via email at westin.robert@epa.gov.

Sincerely,

Mary J. Waller

Mary L. Waller Product Manager (21) Fungicide Branch Registration Division (7505P)

Enclosure

Andersons[™] Golf Products Fungicide IX

For use on the following turfgrass sites: golf course tees, greens, collars, aprons and spot treatment on fairways, as well as athletic fields used only by professional athletics. Do not use on residential lawns and turf and lawns and turf at parks and schools.

 Provides protection from and control of gray snow mold, <u>Pythium</u> blight, dollar spot, brown patch, copper spot and pink snow mold

Net Weight 42.24 lbs. (19.15 kg) Covers Up To 22,000 Sq. Ft.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

lf in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove
	contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor for treatment advice. Have person sip a glass
	of water if able to swallow. Do not induce vomiting unless told to by a poison
	control center or doctor. Do not give anything by mouth to an unconscious person.
16	
lf on skin	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-
	20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then
	give artificial respiration, preferably mouth-to mouth if possible. Call a poison
	control center or doctor for treatment advice.
Have the produ	uct container or label with you when calling a poison control contor or dector or going

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

Environmental Hazards: Do not apply directly to water, to areas where surface water is present or to intertidal areas below mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

ACTIVE INGREDIENTS: Chloroneb, (1,4-Dichloro-2,5-dimethoxybenzene Thiophanate-methyl, (dimethyl 4,4'(-o-phenylene		3.26%	
(3-thioallophanate))		1.63%	
INERT INGREDIENTS:		95.11%	
		100.00%	(L
U.S. Patent No. 4,028,464			• •
EPA Reg. No. 9198-204		i i t	(
EPA Est. 538-OH-1 <u>S</u> , 9198-OH-1 <u>M</u> , 9198-OH-2B, 9198-	-AL-001A : ' '		,
Underlined letter is first letter used in run code on bag		(· · · · ·	ı
Product of U.S.A.	•	1. e.	
ACCEPTE with COMME	NTE		•
In EPA Letter	Deset		
DEC 202			•
Under the Federal Is			

Fundicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

9198-204

Gussets: [Note: gusset information is optional, depending on bag type] RIGHT GUSSET Andersons Golf Products Fungicide IX LEFT GUSSET Andersons Golf Products Fungicide IX TOP GUSSET Andersons Golf Products Fungicide IX

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for homeowner use. For use by individual/firms licensed or registered by the state to apply ornamental or turf pest control products.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

For use on the following turfgrass sites: golf course tees, greens, collars, aprons and spot treatment on fairways, as well as athletic fields used only by professional athletics. Do not use on residential lawns and turf and lawns and turf at parks and schools.

Adapted for use in all areas where listed turf diseases attack turfgrasses. Do not exceed maximum application rates and retreatment intervals listed in the Maximum Use Rates table below.

TO PREVENT GRAY AND PINK SNOW MOLD, apply to dry foliage at the NORMAL or MEDIUM RATE prior to the first snow. Repeat in mid-Winter when turf is free of snow and again immediately after the final snow melts in the Spring. In areas where snow cover is present the entire Winter, apply at the HIGH RATE prior to the first snow and follow with a NORMAL RATE when the snow melts in the Spring, if needed.

TO PREVENT PYTHIUM BLIGHT (when weather conditions are favorable: high temperature and humidity, day and night), apply to moist foliage at the NORMAL or MEDIUM RATE and repeat at the minimum retreatment interval while weather conditions persist.

TO CONTROL ACTIVE PYTHIUM BLIGHT, apply to moist foliage at the MEDIUM RATE as soon as symptoms are observed and repeat as needed if weather conditions remain favorable. For severe attack, apply a HIGH RATE application to moist foliage and repeat at a NORMAL RATE as needed. If disease symptoms do not subside, use a second HIGH RATE application at minimum treatment interval.

TO PREVENT DOLLAR SPOT*, COPPER SPOT AND BROWN PATCH, apply to dry or moist foliage at the NORMAL or MEDIUM RATE when disease activity is noted or suspected, repeat at 2-3 week intervals. During periods of high day and night temperatures and humidity, treatment intervals may need to be shortened for best brown patch control.

TO CURE DOLLAR SPOT, COPPER SPOT AND BROWN PATCH, when symptoms indicate advanced stages, apply at the MEDIUM or HIGH RATE and follow with the NORMAL RATE preventive program.

*If dollar spot control is not effective after using this product as recommended, a thiophanate-methyl or benomyl-tolerant strain of fungi may be present. Consult your ProTurf Tech Rep for recommendations of suitable ProTurf Fungicides to use promptly.

ON NEW SEEDINGS, apply at MEDIUM RATE just after sowing seed. After germination or at first sign of disease, apply to moist foliage and repeat at the minimum retreatment interval if evidence of disease persists.

DRY FOLIAGE APPLICATIONS should be made to prevent snow mold. For prevention or control of Pythium blight, prolonged contact with the foliage enhances activity.

ON GREENS AND FAIRWAYS, it is suggested to apply the product to moist foliage in the morning and evening, followed by a light syringing before mowing or allowing heavy play.

Maximum Use Rates						
	Maximum Single Rate		Maximum Annual Rate		Minimum	
Use sites	lbs. product / 1000 sq. ft	lbs. ai / acre	lbs. product / 1000 sq. ft	lbs. ai / acre	Retreatment Interval	
Residential and public turf areas: home lawns, parks, athletic fields, schools, day care centers	3.84	2.72	15.32	10.88	14 days	
Tees Greens Aprons	11.49	8.16	30.70	21.80	14 days	
Fairways Except Florida			7.67	5.45		
Fairways Florida – only during overseeding			<u>3</u> .84	2.72		

Do not apply in a brisk wind.

PRECAUTIONS

Allow one week between the application of this and other control products. (This guideline can be relaxed where severe insect or disease attack requires immediate treatment).

When treating Pythium and brown patch for control on established turf, do not fertilize or water heavily during periods when disease is active.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

STORAGE: Store in clean, dry place. Reseal opened bag by folding top down and securing. PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

	rovide proper di 2.24 lbs. treats 22 42.24 lbs. treat	uggested Spread stribution, calibrat 2,000 ft ² (½ acre) at s 11,000 ft ² (1/4 acr ats 7,333 ft ² (1/8 acr	e spreader befor the RECOMMENT re) at the DOUBL	IDED RATE E RATE	
	GROUND OR	WIDTH OF COVERAGE	SPREADER SETTINGS		
SPREADER	PTO SPEAD		NORMAL RATE	MEDIUM RATE	HIGH RATE
Gandy - 30"	X mph	Overlap wheels	X	X	X
Scotts Rotaries* R-7, R-7X R-8, R-8A (cone 8)	X mph	XX feet	X	X	X
Lely Model	X mph	XX feet	· X	X	X
Vicon	X mph	XX feet	Х	· X	X

For the R-7X model use with adjustable pattern slide ¼ closed.

Lely models WTR, WFR, HR and 1250. PT) a t 450 rpm.

Note to EPA: actual spreader models may vary depending on market demand and availability

PRODUCT RATE: NORMAL = 1.92 lbs. product/1,000 sq. ft (1.0 + 0.5 oz ai/A) MEDIUM = 3.84 lbs. product/1,000 sq. ft (2.0 + 1.0 oz ai/A) HIGH = 5.76 lbs. product/1,000 sq. ft (3.0 + 1.5 oz ai/A)

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (Rvamd011907)

Manufactured by: The Andersons Lawn Fertilizer Division, Inc. d/b/a Free Flow Fertilizer PO Box 119 Maumee, OH 43537

Date Printed	1/19/07				
Date Approved		•			
Supersedes	8/8/02				
Label Changes	Amendment - gc use only				÷
Filename	9198-204		`	· · · · · · · · · · · · · · · · · · ·	
Label Code	RVamd011907				