9198	- 209		2-14-	-200	D2 Form Approved	. OMB No.	2070-006	O. Approval expires 2-28-9
<b>⊕EP</b> A	Environmenta	United States I <b>I Protectic</b> ington, DC 204	_			Registra Amendr Other	tion	OPP Identifier Number
		Application	n for Pe	sticide	- Section	t		
1. Company/Product Numbe 9198-209	r			. EPA Proc ompkins	luct Manager		3. Pro	posed Classification
4. Company/Product (Name) Andersons Fertilizer Plus Pre		rol	PI 2	<b>₩#</b> 5				J
5. Name and Address of App The Andersons Lawn Fert. PO Box 119 Maumee, OH 43537	Fert. Div., Inc. d		ow (b	)(i), my p :: :PA Reg	No	nilar or ident		FIFRA Section 3(c)(3) mposition and labeling
Check if this	is a new address			roduct	Name			
			Sectio	<u>n - II</u>				
Amendment - Explain Resubmission in resp  Notification - Explain	onse to Agency letter	r dated			al printed label ency letter dat le Too" Applica her - Explain be	etion.	<del></del>	DTIFICATION EB 1 4 2002
This notification is consistent with the this product. I underthed that it is a v 40 CFR 152.48, this product may be	violation of 18 U.S.C. Sec. 100	1 to wilfully make any	r false statement to	EPA. I furth penalties und	or undersant that if th	nis notification is no	ling or the conf of consistent wi	fidentical statement of formula of the terms of PR Notice 98-10 and
1. Material This Product Will	Be Packaged In:							
Child-Resistant Packaging Yes No * Certification must	Unit Peckaging Yes No If "Yes" Unit Peckaging wgt.	No. per	Weter Solu Yes No If "Yes" Package w		eging No. per	2. Type of	Metal Plastic Glass Paper Other (S	pecify)
be submitted  3. Location of Net Contents	Information	4. Size(s) Ret	el Conteiner		IS. Lo	cation of Lab	el Direction	ne
r + r +	ontainer				E	On Label		
6. Manner in Which Label is	Affixed to Product	Lithog Paper Stenoi	glued		Other	· · · · · · · · · · · · · · · · · · ·	······································	
			Section	- IV				
1. Contact Point (Complete	items directly below I	or identification	n of Individue	to be co	ntacted, if nece	essary, to pro	cess this	application.)
Name Dawn L. Walters			Title Regulatory /	Administr	ator		Telephone 334-262-2	No. (Include Ares Code) 766
I certify that the stater I acknowledge that am both under applicable I	y knowlinglly false or	Certifical this form and misleading star	all attachmen	its thereto	are true, accu ble by fine or i	irate and corr imprisonment	plete.	5. Date Application Received (Stamped)
2. Signature Dawn X Waltu	W)	:	3. Title Regulatory A	dministrato	)r		6 (	:
1. Typed Name Dawn L. Walters	Allegae of		5. Date	1/11	102			1

# Page 1 of 8

# ANDERSONS Fertilizer Plus Preemergent Weed Control

- preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, knotweed and Poa annua
- Provides a feeding of potassium
- provides quick green-up plus extended feeding with controlled release SREF nitrogen (slow release encapsulated fertilizer)
- Controlled release Poly-S\*® nitrogen provides uniform growth with extended feeding \*A portion of the urea has been coated to provide 2.5% coated slow-release nitrogen.
- readily available potassium
- provides readily available iron
- provides a feeding of nitrogen and potassium
- provides a one-to-one feeding of nitrogen and potassium
- contains pendimethalin for season-long, broad spectrum weed control
- small particle formulation provides high particle count for better herbicide distribution
- premium formula, designed for golf course use
- contains potassium for turf stress resistance

Net Weight XX.XX lbs (XX.X kg) Treats up to XX,XXX sq. ft. **SGN XXX** 

**NOTIFICATION** 

FEB 1 4 2002

KEEP OUT OF REACH OF CHILDREN CAUTION **PRECAUTINARY STATEMENTS** 

Hazards to Humans and Domestic Animals: CAUTION, causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed.

#### FIRST AID

If swallowed: Call a poison control center immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling poison control center or doctor, or going for treatment.

#### Personal Protective Equipment

Applicators and other handlers must wear: Long-sleeved shirt and long pants, chemical resistant gloves such as butyl rubber, barrier laminate, or nitrile rubber > 14 mils (if you want more options follow instructions for Category A on EPA chemical-resistant selection chart), shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protection equipment (PPF). If no such instructions for washables, use detergent and hot water. Keep and wash personal protection equipment separately from other laundry.

# (Front Panel cont'd)

# **User Safety Recommendations:**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

# **Engineering Controls**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

<u>Environmental Hazards:</u> This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. In case of spills, either collect for use or dispose of properly.

#### **ACTIVE INGREDIENT:**

Pendimethalin†	 0.84%
<b>INERT INGREDIENTS</b>	 99.16%
	100.00%

†CAS #40487-42-1 EPA Reg. No. <u>9198-209</u> EPA Est 538-OH-1

ANDERSONS Fertilizer Plus Preemergent Weed Control XX-XX-XX 643-XXXX

#### **Guaranteed Analysis**

Total nitrogen (N)	XX%
0.6% ammoniacal nitrogen	
22.9% urea, methylene ureas nitrogen	
6.5% water insoluble nitrogen	
Available phosphate ((P <sub>2</sub> Q <sub>5</sub> )X%	ı
Soluble potash (K <sub>2</sub> O)X%	
Derived from: urea, methylene ureas, monoammonium phosphate and potassiu	m sulfate

Information regarding the contents and levels of metals in this product is available on the internet at http://www.regulatory-info-ap.com

# Product of USA

U.S. Patent No's 3,705,794; 3,989,470

\*Licensed under one or more of the following U.S. Patents: 3,920,742; 4,066,441

# The Andersons Lawn Fertilizer Division, Inc.

PO Box 119

Maumee, OH 43537

Recommended for Use by Professional Turfgrass Managers

**RIGHT GUSSET** 

XXXX

Andersons XX-XX-XX

Logo ANDERSONS Fertilizer

Plus Preemergent
Weed Control

**LEFT GUSSET** 

XXXX

Andersons XX-XX-XX

Logo ANDERSONS Fertilizer

Plus Preemergent
Weed Control

#### BACK:

Prevents sprouting of the following grassy and dicot weeds and provides [a feeding of potassium –OR— a controlled-release nitrogen feeding]:

# Grassy Weeds

# Dicot Weeds

crabgrass
foxtail
goosegrass
Poa annua
barnyardgrass
fall panicum

chickweed oxalis knotweed prostrate spurge

purslane cudweed hop clover henbit

eveningprimrose

#### Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Workers Protection Standard 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, chemical resistant gloves such as butyl rubber, barrier laminate; or nitrile rubber ≥ 14 mils (if you want more options follow instructions for Category A on EPA chemical-resistant selection chart), shoes plus socks.

## **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to use of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The Worker Protection Standard applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment are during application. Do not enter treatment areas until spray has dried or dusts have settled.

For use on established turfgrasses including bentgrass, bermudagrass, and *Poa annua* (except putting greens and tees), Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of the grasses. See "Bermudagrass, Bentgrass and *Poa annua* Precautions" and "Limitation of Liability" statements.

<u>Not</u> for use on bentgrass, *Poa annua* and bermudagrass putting greens and tees, or on dichondra. Where bentgrass, bentgrass-containing turf and *Poa annua* are the desired species, a light rate application only is recommended.

# How/When to Use

For control of annual grassy and broadleaf weeds such as crabgrass, goosegrass, foxtail, barnyardgrass, fail panicum, *Poa annua*, oxalis, knotweed, purslane, and prostrate spruge, apply in spring before they sprout.

To prevent *Poa annua* and fall germinating chickweed, apply product to dry foliage in late summer/early fall (before germination). Repeat application yearly for continued control. A repeat application made after *Poa annua* treatment should not be sooner than 4 months.

Where lower infestations are expected, the lighter rates may be used.

Apply product uniformly to dry foliage at the suggested rate with a properly calibrated spreader. For best results, water after application ( $^1/_4 - \frac{1}{2}$  inch) to remove granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying as a segment to application. Allow at least a four-month space between the applications except when repeat applications are necessary after 6-8 weeks.

Do not mow treated area until product has been watered in and turf foliage has dried.

# **Precautions**

Do not use on bentgrass, *Poa annua* and bermudagrass putting greens and tees, or on dichondra. Where bentgrass and bentgrass-containing turf and *Poa annua* are the desired species, a light rate application only is recommended.

On cool season turfgrasses avoid applications when temperatures are regularly in the 80's.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under sever stress conditions (disease, drought, low temperature – below 40°F, scalping, and excessive soil moisture).

Do not apply to dormant bentgrass and *Poa Annua* (turf that is not actively growing or has experienced frost).

9198-209 Page 5 of 8 6/9

When any preemergence herbicide is applied to dormant bermudagrass, green-up may be delayed 2 to 3 weeks in the spring. The delay in green-up may be extensive in low-cut bermudagrass and on sandy soils.

As with most other preemergence herbicides, application of this product to bermudagrass overseeded with cool season turfgrass species may injure and thin the overseeding.

Do not apply to bentgrass of Poa annua after late summer.

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water.

If metallic surfaces (golf course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5-10 MPH) due to its staining potential.

Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

# Properties of All Dinitroanaline (DNA) Products

May cause temporary slowing of turf fill-in and stolon-rhizome rooting where turf has been thinned.

Requires a waiting period before seeding or sprigging.

# Seeding or Sprigging Intervals after Treatment

Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

# Treatment Interval after Seeding or Sprigging

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

#### Notes

Do not apply more than a cumulative total of 349 lbs. of product/acre (3.0 lbs. active ingredient/acre) per year on bentgrass and *Poa annua*; 698 lbs. of product/acre (6.0 lbs. active ingredient/acre) per year on cool season turfgrass; and 1,046 lbs. of product/acre (9.0 lbs. active ingredient/acre) per year on warm season turfgrasses.

Turfgrass Species	Weeds	Rates	Comments
Cool Season Grasses  Kentucky bluegrass	Crabgrass Foxtail Poa annua	175 – 233 lbs./acre as the initial application prior to weed germination	Apply a repeat application of 145 – 175 lbs/acre after 6 –
Perennial ryegrass Fine fescue Tall fescue	Barnyardgrass Fall panicum Oxalis Prostrate spurge Purslane Knotweed Chickweed	weed germalation	8 weeks for extended control or where heavy weed infestations are expected
	Goosegrass	175 – 233 lbs./acre in the spring prior to weed germination. Apply a repeat application after 6 – 8 weeks at 175 – 203 lbs./acre	
Bentgrass and established <i>Poa Annua</i> (1/2 inch height or taller). Not for use on greens and tees	Crabgrass Foxtail Poa annua Barnyardgrass Fall panicum Oxalis Prostrate spurge Purslane Knotweed Chickweed	175 lbs./acre in the spring prior to weed germination	Apply a repeat application of 145 – 175 lbs/acre after 6 – 8 weeks where heavy weed infestations are expected
	Goosegrass	175 lbs./acre in the spring prior to weed germination with a repeat application of 175 lbs./acre after 6 – 8 weeks	

Warm Season Grasses Bermudagrass St. Augustinegrass Bahiagrass Zoysiagrass Centipedegrass (401 lbs./acre rate or less) Tall fescue (Not for use on greens and tees)	Crabgrass Foxtail Poa annua Barnyardgrass Fall panicum Oxalis Prostrate spurge Purslane Knotweed Chickweed	OR (OPTIONAL WORDING) 233 lbs./acre may be used prior to weed germination with a repeat application of 175 lbs./acre after 4 months where weeds germinate over an extended period. (Do not use this high rate on centipedegrass)	Apply a repeat application of 145 - 175 lbs./acre after 6 - 8 weeks for extended control or where heavy weed infestations are expected
	Goosegrass Cudweed	175 – 233 lbs./acre prior to weed germination with a	
	Hop clover	repeat application of 175 -	
	Evening-	233 lbs./acre after 6 – 8	
}	primrose	weeks.	
	Henbit		
		<u>OR</u>	
		OTE TO EPA - OPTIONAL W	
	Goosegrass	175 - 233 lbs./acre prior to	Apply a repeat
		weed germination	application of 175 – 233 lbs./acre to
			extend control of
			goosegrass or if the
			lower rate was used initially
	Cudweed	175 - 233 lbs./acre prior to	Apply a repeat
	Hop clover	weed germination	application of 145 –
	Evening-		203lbs./acre after 6 -
	Primrose		8 weeks if the lower
	Henbit		rate was used initially

# Storage and disposal

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Reseal opened bag boy folding top down and securing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

# **Suggested Spreader Settings**

To provide proper distribution calibrate spreader before application. Do not adjust rates in order to meet fertilizer requirements of your soil.

50 lbs. treat 10,729 ft<sup>2</sup> (1/4 acre) at the 203 lbs. product/acre rate (equivalent to 1.75 lbs. active ingredient/acre and X lbs. N/1,000 sq ft)

			RECOMMENDED RATE RANG (lbs.product/acre)			
SPREADER	GROUND OR PTO SPEAD	WIDTH OF COVERAGE	145	175	203	233
Gandy - 30"	X mph	Overlap wheels	Х	Х	X	Х
Scotts Rotaries* R-7, R-7X R-8, R-8A (cone 8)	X mph	XX feet	X	X	X	X
Lely Model	X mph	XX feet	Х	X	X	Х
Vicon	X mph	XX feet	Х	X	X	X

\*Use discharge hole cover #13030 (already attached to the PT-10 and PT-11 models).

For the R-7X model use with adjustable pattern slide ¼ closed.

Lely models WTR, WFR, HR and 1250, PT) at 450 rpm.

# **IMPORTANT: READ BEFORE USE:**

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

# WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

# **LEGAL RIGHTS:**

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

APtrans/Rynot072301

## Manufactured by:

The Andersons Lawn Fertilizer Division, Inc.

d/b/a Free Flow Fertilizer

PO Box 119

Maumee, OH 43537

Date Printed	1/10/02	1 1	
Date Approved	Transfer	• • •	,
Supersedes	Transfer		3 ' +
Label changes	Notification: new reg no/our name/warranty	1 * 4 *	3
Filename	9198-209	1 *	•
Label code	(APtrans/RVnot072301)		