



9198-223

08-28-2009

1/7

 <p style="text-align: center;">U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460</p>	EPA Reg. Number: 9198-223	Date of Issuance: AUG 28 2009
	Terms of Issuance: Unconditional	
	Name of Pesticide Product: The Andersons Insect Killer Granules with 2% Carbaryl	
<p>NOTICE OF PESTICIDE: <u> </u> Registration <u>XX</u> Reregistration (under FIFRA, as amended)</p>		
<p>Name and Address of Registrant (include ZIP Code): The Anderson Lawn Fertilizer Division P.O. Box 119 Naumee, OH 43537</p>		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p>		
<p>Based on your response to the Reregistration Eligibility Document(s), EPA has reregistered the product listed above. This action is taken under the authority of section 4(g)(2)(c) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product. Within 45 days from the date of this notice, two copies of a final printed label.</p>		
<p>Your Confidential Statement of Formula (CSF) dated May 19, 2006 is acceptable. This CSF supercedes all previous CSFs. You may distribute products bearing the old label for 12 months from receipt of this letter or your next printing of the label, whichever comes first.</p>		
<p>A copy of the stamped label is enclosed for your records.</p>		
<p>Signature of Approving Official:  Meredith Laws, Branch Chief Registration Division, Insecticide-Rodenticide Branch</p>		<p>Date: AUG 28 2009</p>

(Front panel)

THE ANDERSONS INSECT KILLER GRANULES WITH 2.0% CARBARYL

Intended For Residential Use

- Kills Listed Pests on Contact
- Kills Listed Surface & Subsurface Insects
- also kills chinch bugs, cutworms & other listed Insects
- Ready-to-Use
- Kills on Contact
- Kills Ants, Fleas, Brown Dog Ticks, Chinch Bugs, Armyworms
- Kills spiders, cutworms, scorpions and other listed insects
- Kills listed Lawn & Ornamental Nuisance Pests Around the Outside of Homes
- Ant, Flea, Tick & Grub Killer
- Kills ants, fleas, ticks and grubs
- Kills listed insects
- For use outdoors on lawns, gardens and around the outside of homes
- Kills ants, fleas, ticks & other listed insects
- Kills Listed Insects
- For use on lawns
- Kills listed home-invading pests
- Kills listed surface and foliage feeding pests
- Kills ants, fleas, ticks, cutworms, chinch bugs & other listed pests
- Ready-to-Spread Granules
- Easy to use
- Lawn Insect Killer Granules
- Easy-to-Use Granules
- Use on Lawn, Ornamentals and Around the Outside of Homes

Active Ingredient:


Carbaryl*	2.0%
Other Ingredients:	<u>98.0%</u>
Total:	100.00%

*CAS# 63-25-2

EPA REG. NO. 9198-223
 EPA Est. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A
 Underlined letter is first letter used in run code on bag

KEEP OUT OF REACH OF CHILDREN
CAUTION
 See Back Panel For Additional Precautionary Statements

NET WEIGHT _____ LBS.
 TREATS UP TO _____ SQ. FT.



(Back panel)

NET WEIGHT _____ LBS.
TREATS UP TO _____ SQ. FT.

THE ANDERSONS INSECT KILLER GRANULES WITH 2.0% CARBARYL

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear long-sleeved shirt and long pants, and shoes plus socks and chemical-resistant gloves (HOUSEHOLD OR LATEX RUBBER GLOVES) WHEN HANDLING THIS PRODUCT. REMOVE CLOTHING IMMEDIATELY if pesticide gets inside. Then wash thoroughly and put on clean clothing. Wash hands before eating, drinking, chewing gum, tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

First Aid

Carbaryl is an N-Methyl Carbamate Insecticide.

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing; Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

ANTIDOTE STATEMENT: ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN.

NOTE TO PHYSICIAN:

Carbaryl is a N-methyl carbamate that inhibits cholinesterase. Atropine is antidotal. Do not use 2-PAM (pyridene-2-aldoxime methiodide), opiates or cholinesterase inhibiting drugs. Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or allow drift to water. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in any manner inconsistent with its labeling.

Read entire label before using this product.

Do not allow people or pets to enter the treated area until dusts have settled. Do not allow people (except those involved in the water-in) or pets to enter the treated area until the water-in is completed and the area has dried.

(Back panel)

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. This product may be applied only by broadcast or drop type spreaders. Do not apply by hand, spoon or belly grinder. For outdoor use only.

Watering-in is required immediately after application to lawns

Kills turf grass insects, apply granules uniformly to the turf grass.

INSECT PEST		LBS OF PRODUCT PER 1,000 SQ. FT.	SPECIFIC DIRECTIONS
Ants (except harvester, pharaoh and carpenter ants) Armyworms Centipedes Chiggers Chinch Bugs Cutworms Darkling Ground Beetles Earwigs Essex Skipper European Chafer Fall Armyworm Fiery Skipper Grasshoppers Green June Beetle Grubs Lawn Moths (Sod Webworm)	Leafhoppers Lucrene Moth Millipedes Scarab Beetle adults (such as May Beetle, June Beetle, Japanese Beetle, Green June Beetle) Scorpions Silverfish Sowbugs Spiders Spittlebugs Springtails Yellowstriped Armyworms	2.25	For kill of listed pests, apply granules uniformly when grass is dry. Product must be watered in immediately after application. Repeat applications may be made up to a total of four times per year but not more often than once every seven days. Will only kill pests that are present at the time of application. Pest must be directly contacted by the granules to be effective.
White Grubs (such as Japanese Beetle Grubs, Chafer Beetles, <i>Phyllophaga</i> spp. Grubs and Annual White Grubs) Bluegrass billbugs larvae European crane fly larvae Black Turfgrass atanius beetle larvae Hyperodes weevil (Bluegrass weevil Larvae)		9.0	For kill of grubs, applications should be made when grubs are feeding near the soil surface, usually during late March through May or July to early September, or timing as recommended by local agricultural extension agents. Product must be watered in immediately after application.

TO KILL TICKS WHICH MAY TRANSMIT LYME DISEASE

To kill juvenile and adult Ixodes spp. Ticks (Deer Tick, Bear Tick, and Black Legged Tick). Apply granules at the rate of 1.15 lbs of product per 1,000 square feet. Begin applications in late spring or early summer (May/June). Uniformly apply granules to entire lawn, perimeter wooded areas, and property boundaries including shrub beds, ornamental plantings, and wooded areas where exposure to ticks may occur. Product must be watered in immediately after application to lawns. Will only kill ticks present at the time of application. Pest must be directly contacted by granule to be effective. Reapply as necessary, but no more than 4 times per year. Minimum reapplication interval is 7 days.

(Back panel)

NUISANCE PEST KILL IN AND AROUND FLOWER BEDS AND ORNAMENTAL PLANTINGS AND AROUND HOMES

To kill Ants (except harvester ants, pharaoh ants and carpenter ants), Armyworms, Centipedes, Crickets, Cutworms, Earwigs, Firebrats, Silverfish, Fleas, Millipedes, Sowbugs, Scorpions, and Spiders; apply granules uniformly in a 6 foot band around the building. Apply granules at the rate of 2.5 – 5.0 lbs. per 1,000 square feet to shrub and flower beds, foundations, ornamental plantings, and lawn or soil areas immediately adjacent to the building. Product must be watered in immediately after application. Repeat treatments as necessary to kill new infestations of listed pests. Do not reapply more often than 4 times per year. Minimum reapplication interval is 7 days.

HOME FRUIT AND VEGETABLE GARDEN

To kill Ants (except harvester ants, pharaoh ants and carpenter ants), Centipedes, Crickets, Cutworms, Earwigs, Grasshoppers, Millipedes, Sowbugs (Pillbugs), Springtails, refer to Use Rate chart below for application information. Apply specified rate of product evenly over the soil surface. Water lightly after application to release the chemical from the granules. Applications may be made at any time before, during and after the growing season. If applying during the growing season, observe indicated pre-harvest intervals (PHI). Do not apply within the number of days indicated in the commodity's PHI before harvesting.

Commodity	Application Rate (Lbs./1,000 sq. ft.)	Maximum Yearly Application (Lbs./1,000 sq. ft.)	Retreatment Interval Days	PHI (days)	Comments
Asparagus (Spears)	2.3	6.9	3	1	Total maximum annual application is 11 lbs per 1,000 sq. ft. per crop. The total number of application is 3 prior to harvest with 2 additional applications only to ferns. * after harvest of spears
Asparagus (Ferns*)	2.3	4.6	7	1	
Beans, dried (snap, navy and kidney), peas, dried, cowpea, lentil	1.7	6.9	7	3 (EP) 21(D)	EP = edible-podded D - = dried Total number of applications per crop per year can be 4. Use on succulent peas or beans is prohibited.
Garden beets (roots), carrots, horseradish, radish, parsnip, rutabaga, salsify, turnips (roots)	1.2 - 2.3	6.9	7	7	Total number of applications per crop per year can be 6 (some of which must be at lower than maximum single application rate)
Brussels sprouts, broccoli, cabbage, cauliflower, Chinese cabbage, collards, kale, kohlrabi, mustard greens	1.7 - 2.3	6.9	7	3 (A) 14 (B)	A = Brussels sprouts, broccoli, cabbage, kohlrabi, cauliflower B= Chinese cabbage, collards, kale, mustard greens Total number of applications per crop can be 4 (some of which must be lower than the maximum single application rate)
Cucumber, melon, squash	1.2	6.9	7	3	Total number of applications per crop per year can be 6

6
7

(Back panel)

Potato	2.3	6.9	7	7	Total number of applications per crop per year can be 3
Strawberry	2.3	11.5	7	7	Total number of applications per crop per year can be 5
Tomato, pepper, eggplant	2.3	9.2	7	3	Total number of applications per crop per year can be 4

STORAGE AND DISPOSAL

STORAGE: Store unused product in original container only, in cool, dry area out of the reach of children and animals. Keep container closed and away from food, feedstuffs and domestic water supplies. Do not store in areas where temperatures frequently exceed 100° F.

CONTAINER DISPOSAL

If empty: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.

PESTICIDE DISPOSAL

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

SPREADER SETTINGS

IMPORTANT: The above spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of terrain. Calculate the size of the area to be treated and do not over or under apply. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting, if necessary, in order to assure correct rate of application. Close spreader when filling, stopping or turning, do not hand spread.

Broadcast spreaders

For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of product will ensure complete coverage. *[diagram of path] [listing of settings]*

Drop spreaders

First apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips. *[diagram of path] [listing of settings]*

Optional:

To determine square footage of application area, follow calculation chart. *[lawn/turf diagram]*

$$\begin{aligned}
 & \text{___ length} \\
 \times & \text{___ width} \\
 = & \text{___ Total Sq. Ft.} \\
 - & \text{___ Non-lawn areas} \\
 = & \text{___ Total Sq. Ft.}
 \end{aligned}$$

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer

7
7

(Back panel)

makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Manufactured by:
THE ANDERSONS LAWN FERTILIZER DIVISION, Inc.
D/b/a Free Flow Fertilizer, Inc.
P.O. BOX 119
MAUMEE, OHIO 43537

Date Printed	5/4/09
Date Approved	1/24/05
Supersedes	New
Label Change	Amendment - RED label changes
Filename	9198-223
Label code	RVamd050409