

Net Wt. 10 Lbs. (4.5 Kg.)

Treats 2,000 Sq. Ft.

TURF CARE Lawn Insecticide

2% DIAZINON

ACTIVE INGREDIENTS
O,O-Diethyl O-2-methyl-4-phosphorothioate 2.00%
INERT INGREDIENTS 98.00%
TOTAL 100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals CAUTION

Harmful if swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of food and feed products. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

STATEMENT OF PRACTICAL TREATMENT

If swallowed - Drink plenty of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled - Remove victim to fresh air. Apply respiration if indicated.

If on skin - Remove contaminated clothing and wash affected areas with soap and water.

If in eyes - Flush eyes for at least 15 minutes with water. Call a physician if irritation persists.

First Aid Physician - If signs of cholinesterase inhibition are present, atropine is an antidote. 2-PAM is also an antidote and may be administered in conjunction with atropine.

Environmental Hazards

This product is toxic to fish, birds and other wildlife. Birds and other wildlife feeding on treated areas may be killed. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by running of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply this product directly to blooming crops or weeds having a high bee activity in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

DIRECTIONS FOR USE

This is a contact type insecticide. Use this product in a manner consistent with this label.

Turf Care Lawn Insecticide controls the following lawn insects on grass and Dichondra lawns:

Ants, Armyworms, Beetles, Mites, Brown Dog Ticks, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Hymenoptera Weevils (New York only), Lawnmoths (Sod Webworms), Leatherjackets, Millipedes, Slugs, Snails, Springtails (Chironomids).

Do not hand spread. Apply Turf Care Lawn Insecticide by broadcast or band application. For broadcast application, use a 5 foot band applicator in a circular or rectangular pattern. Apply at a rate of 5 lbs. per 1,000 sq. ft. of area. For areas 20' by 10' or 8' by 100' sq. ft. (an area 10' by 100'). Four heavy textured lawns such as St. Augustine use 7.5 lbs. per 1,000 sq. ft. for 12 oz. per 100 sq. ft. Use back panel for recommended spreader settings. Apply when grass is dry. Water lawn lightly after application. Repeat as necessary.

Turf Care Lawn Insecticide also controls Billbugs and White Grubs of Japanese Beetle, European Chaffer, Southern Chaffer and Acanthos spretus (Dung Beetle). Apply at 6 lbs. per 1,000 sq. ft. for all except Dung Beetle, which requires 7 lbs. per 1,000 sq. ft. for adequate control. For Billbugs, apply when ant activity is first observed, or when the lawn shows signs of damage. For Grub control, apply when the lawn shows signs of damage, or when the lawn shows signs of damage. Apply in spring or early summer when damage is first noticed. Thoroughly water grass with 1/4" to 2" of water after application. Repeat treatment in late summer or early fall if a second generation is in the lawn. For Japanese Beetle, apply 5 lbs. per 1,000 sq. ft. in mid April and again in mid May, watering thoroughly after each application. See back panel for recommended spreader settings.


Turf Care Lawn Insecticide can be used for localized control of ants and Digger Wasps. Apply one pint of insecticide around each ant or wasp nest. Apply to wasp nests around dusk when activity is low.

STORAGE AND DISPOSAL

Keep container closed and away from heat, fire, hot surfaces, and domestic animals. Do not use empty container. Wrap container and put in trash.

WARRANTY STATEMENT

Seal of acceptance of this material is hereby made by the undersigned, who is authorized to make such a seal of acceptance. The undersigned is a duly qualified and licensed professional person in the State of New York, and is authorized to make such a seal of acceptance. The undersigned is a duly qualified and licensed professional person in the State of New York, and is authorized to make such a seal of acceptance. The undersigned is a duly qualified and licensed professional person in the State of New York, and is authorized to make such a seal of acceptance.



Manufactured by **The Green Andersons**

Lawn Fertilizer Division • P.O. Box 119 • Maumee, Ohio 43537

EPA REG. NO. 499-61
EPA EST. NO. 499-61