KILL and PREVENT

Non Extendina Vegetal in Killer-

GENERAL INFORMATION:

A liquid actal weat killer. To be applied undiluted as a spray on non-cropland areas where bere ground is desired. The degree of weed control and duration of effect will vary with the amount of chemical applied, soil type, rainfall and other conditions.

DIRECTIONS: Apply any time of year except when the ground is frozen. Best results are obtained after predominant species are well established in late spring or early summer. Before treating, calibrate equipment to insure uniform cover of the area to be treated. Measure out the proper amount and apply.

pressed air hand sprayers, knapsack sprayers or an ordinary sprinkler can be used to apply. No dilution or agitation is required.

NET CONTENTS:

☐ U.S. Gals.

US WA THEY

KILLS EE VEGETATION WEEDS CAN BE KILLED QUICKLY

KEEP OUT OF REACH OF

WARNING:

HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN. CAUSES SKIN IRRITATION. SEE SIDE PANEL FOR FURTHER CAUTION STATEMENTS. KEEP AWAY FROM CHILDREN.

ACTIVE INGREDIENTS:

*2,4-Dichlorophenoxyacetic acid equivalent 0.70%

CHERNICA AND FREDRETS OF

CAUTION:

Do not apply figuid directly to, or otherwise permit it to come into contact with susceptible plants, such as vegetables, flowers, fruit trees, ornamentals, cotton, grapes, or other desirable plants and do not permit spray mists containing it to drift onto them. Icoarse sprays are less likely to drift). Vápors from this product may injure susceptible plants in the immediate vicinity. Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes (or do not use same spray equipment for other purposes). Do not store near fertilizers, seeds, insecticides, or fungicides. Do not contarninate waters used for irrigation or domestic purposes. Do not re-use this container for other materials without thorough cleaning.

Controlling to some

CAUTION: Harmful if swallowed or absorbed through skin! Causes Skin Irritations! Do not breathe vapor or spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. In case of contact wash skin with soap and water; for eyes, immediately flush with large amounts of water for at least 15 minutes and get medical attention. Remove and wash contaminated clothing before re-use.

Krippetteller Market Con Krippetteller Strenger Apply
the premise Apply
the premise to

GENERAL WEED CONTROL: To control most weeds for an extended period of time on noncropland areas such as parking lots, fence rows, loading ramps, railroad sidings, storage yards, a point buildings and industrial plant sites, apply 2 to 5 gals. per 1,000 sq. ft. treated (80 to 200 gals. per acre).

To control annual weeds and grasses such as foxtail, ryegrass, wild oats, crabgrass, cheatgrass, broomsedge, ragweed, lambsquarter and turkey mullein, apply 1 to 2 gals. per 1,000 sq. ft. (40 to 80 gals. per acre).

For other HARD-TO-KILL perennial weeds and grasses such as Johnson-grass, saltgrass, bouncing bet., bracken fern and horsetail apply 3 to 5 gals. per 1,000 sq. ft. (130 to 220 gals. per acre).

To control perennial weeds and grasses, such as smooth brome, Canada bluegrass, purpletop, quackgrass, broomsedge, aster, dandelion, dog fennel, goldenrod, plaintain and wild carrot, apply 2 to 4 gals. per 1,000 sq. ft. (80 to 160 gals. per acre).