

SUPER - CLO

A SANITIZER FOR FOOD PROCESSING
PLANTS AND EQUIPMENT

ACTIVE INGREDIENTS:
SODIUM HYPOCHLORITE... 6%
INERT INGREDIENTS:..... 94%
TOTAL 100%

CAUTION: Keep out of reach of children. See side panel for further cautions.
DANGER

ONE U.S. GALLON

MANUFACTURED BY
KANSAS CITY CHEMICAL CO., INC.
310 DELAWARE KANSAS CITY, MO. 64105
(816) 842-7586

EPA Reg. No. 9218-2

EPA Est. No. 9218-MO-1

7

ACCEPTED
with COMMENTS
in EPA Reg. No. 9218-2


NO. 28 1398

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this pesticide is registered under EPA Reg. No. 9218-2

9218-2

FRONT PANEL

GOLDEN ROD STOCK



DIRECTIONS FOR USE
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with it's labeling.

SANITIZATION OF NONPOROUS FOOD CONTACT SURFACES

RINSE METHOD:

A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to insure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 1 oz. of this product with 4.5 gallons of water. If no test kit is available, prepare a sanitizing solution by thoroughly mixing 2 oz. of this product with 4.5 gallons of water to provide approximately 200 ppm available chlorine by weight.

Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. If solution contains less than 50 ppm available chlorine, as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Do not rinse equipment with water after treatment and do not soak equipment overnight.

SANITIZATION OF POROUS FOOD CONTACT SURFACES

RINSE METHOD:

Prepare a sanitizing solution by thoroughly mixing this product with 4.5 gallons of water to provide 600 ppm available chlorine by weight. Clean in the normal manner. Prior to use, rinse all surfaces with the sanitizing solution, maintaining contact for at least 2 minutes. Rinse equipment thoroughly with water and do not soak equipment overnight.

Do not

DILUTION TABLE

PPM	SUPER-CLO
100	1 oz.
150	1 oz.
200	1 oz.

SUPER-CLO is recommended for the sanitizing of milk, ice cream, cheese and butter plant equipment, packers, poultry packers and food processing equipment.

SUPER-CLO is recommended for chlorinating and rinsing water. Also for sweet water tanks.

RIGHT PANEL

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS

DANGER

Corrosive, may cause severe skin irritation or chemical burns to broken skin. Causes eye damage. Do not get in eyes, on skin or clothing. Wear face shield or goggles and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until strong odors have dissipated.

FIRST AID

If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of milk or gelatin solution or, if these are not available, drink large quantities of water. DO NOT give vinegar or other acids. DO NOT induce vomiting. Get prompt medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not discharge into lakes, streams, ponds or public waterways. For guidance, contact the regional office of the Environmental Protection Agency.

PHYSICAL AND CHEMICAL HAZARDS
STRONG OXIDIZING AGENT

Mix only with water according to label directions. Mixing this product with chemicals (ammonia, acids, detergents, ect.) or organic matter (urin, feces, ect.) will release chlorine gas which is irritating to eyes, lungs and mucous membranes.

S STORAGE AND DISPOSAL

Store product in a cool dry area, away from direct sunlight. In case of spill flood areas with large quantities of water. Product or rinsate that cannot be used should be diluted with water before disposal in a sanitary sewer. Do not reuse container. Rinse thoroughly and offer for recycling or dispose of in sanitary landfill. Do not contaminate food or feed by storage, disposal or cleaning of equipment.

NOTE:

This product degrades with age. Use a chlorine test kit and increase dosage, as necessary, to obtain the required level of available chlorine.

KEEP OUT OF REACH OF CHILDREN
DANGER

LEFT PANEL