

JUN 17 1988

Huber Janitor Supplies, Inc.
114 North St. Francis
Wichita, KS 67202

Attention: Dorothy Tillery

Gentlemen:

Subject: Sani-Dis Rinse
EPA Registration No. 9228-5
Your Submission Dated September 16, 1986

BEST AVAILABLE COPY

The amendment referred to above, submitted in connection with the Label Improvement Program Notice for Antimicrobial Products Used as Food Contact Surface Sanitizers, is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling.
 - a. Expand the sanitizing use directions to include a contact time of at least 1 minute. The directions must also provide instructions to drain the use solution from the surface and air dry.
 - b. Expand the sanitizing direction to state that a fresh sanitizing solution must be prepared at least daily or more often if the solution becomes diluted or soiled.
 - c. Add the following statement under the USE DIRECTIONS heading.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

93150: I:lee: L-7: KENCO: 4/28/87: 5/7/87: de:vo


CONCURRENCES

SYMBOL	ORIGINATOR							
SURNAME								
DATE								

Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,


John H. Lee
Product Manager (31)
Disinfectants Branch
Registration Division (TS-767C)

Enclosure

BEST AVAILABLE COPY

Cold Disinfection

For disinfection of previously cleaned surgical instruments, rubber tools and dental equipment, simply submerge instruments into solution containing 1 1/2 ounce Sani-Dis Rinse per gallon of water for 10 minutes.

FOR ROOM FOGGING

DIRECTIONS:

To minimize the danger of cross-contamination from environmental surfaces apply solution of Sani-Dis Rinse as a fog as described below, before applying standard routine terminal cleaning and disinfecting practices. 1. Remove all human, animal, and plant life from room. 2. Open closed doors and drawers. 3. Set up fogging platform in center of room. 4. Mount fogging device discharging equivalent spray at 3 RPM. 5. Fill sprayer reservoir with 1 1/2 oz. of Sani-Dis Rinse to 1 gallon of water to produce 1150 ppm active ingredients. 6. Set sprayer mechanism to deliver 1 gallon of solution. 7. Fog for 15 minutes for an average 2100 cu. ft. room. For different room sizes, vary fogging time proportionately to ensure complete wetting of exposed surfaces. 8. Wait 2 hours before re-entring room after treatment. 9. Rinse fogging equipment thoroughly with clear water following use.

DANGER

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food.

FIRST AID


In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN:

Frobbable mucosa. Damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard. Storage and Disposal — Storage: Store in a dry place. Keep container closed when not in use.

Disposal: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by state or local authorities, by burning. If burned, stay out of smoke.



SANI-DIS RINSE

DISINFECTANT DEODORIZER SANITIZER

ACTIVE INGREDIENTS:

n A'ky: (60% C14, 30% C16, 5% C12, 5% C18)
dimethyl benzyl ammonium chlorides 5%
n A'ky: (65% C12, 32% C14) dimethyl
dihydrobenzyl ammonium chlorides 5%

INERT INGREDIENTS:

90%

100%

Sani-Dis Rinse is a concentrate and should be diluted before using.

NET CONTENTS:

EPA Reg. No. 9228-5 EPA Est. No. 9228-KS-1

DANGER

KEEP OUT OF REACH OF CHILDREN

(See other precautionary and first aid statements on left side panel.)

HUBER JANITOR SUPPLIES, INC.

114 NORTH ST. FRANCIS
WICHITA, KANSAS 67202

Only for Sale to, Use and Storage by Service Personnel

WARRANTY

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guaranteed expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials. Statements concerning the use of these products are not to be construed as recommending the infringement of any patent, and no liability for infringement resulting out of any such use is assumed.

USE DIRECTIONS:

Commercial

For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices and Shower Stalls. Apply solution with mop or cloth. Sanitize with 1 ounce Sani-Dis Rinse to 4 gallons of water. Disinfect with 3 ounces Sani-Dis Rinse to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

Industrial Inanimate Surfaces

Floors, walls and other hard surfaces in Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms. Use Sani-Dis Rinse at 3 1/2 ounces per 5 gallons of water. Apply with cloth, mop, or sponge. At this level, Sani-Dis Rinse will be effective against Pseudomonas aeruginosa PRD10. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

Meat and Food Processing Equipment

For sanitization of previously cleaned food processing equipment and food items, dilute 1 ounce Sani-Dis Rinse for 4 gallons of water to provide 200 ppm of active quaternaries. At this level, No Potable Water Rinse is required.

Dairies

To sanitize dairy equipment such as tanks, lines, pails and milk cans, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce Sani-Dis Rinse to 4 gallons of water (200 ppm). At this level, No Potable Water Rinse is required. Follow recommendations of local Health Board.

Restaurant and Bar Rinse

Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent, rinse thoroughly and immerse in a sanitizing solution containing 1 ounce Sani-Dis Rinse to 4 gallons of water. No Terminal Potable Water Rinse Required.

This product when used on environmental, inanimate, hard surfaces at 3 1/2 ounces per 5 gallons of water is effective against Influenza A2, Herpes Simplex, Adenovirus type 5, and Vaccinia Viruses.

CONTENTS: 1 GALLON

BEST AVAILABLE COPY