

ACCEPTED

JUN 12 1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 9275-10

Net Weight 50 Pounds

10% SEVIN DUST

ACTIVE INGREDIENTS

Carbaryl* (1-Naphthyl N-Methylcarbamate) 10.00%

INERT INGREDIENTS 90.00%

TOTAL 100.00%

*Trade mark of Union Carbide Corp. for 1-Naphthyl N-Methylcarbamate

CAUTION

Keep Out of Reach of Children

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Do not contaminate foodstuffs, feed, feeding troughs or receptacles.

DIRECTIONS

Use sufficient amount of dust to obtain full coverage. Apply when insects or their signs appear. Repeat at 5 to 7 day intervals as necessary for good control. Use in accordance with label directions and cautions. Sevin may be applied up to time of harvest.

COTTON - To control Boll Weevils, Boll Worms, Trips, Mea Beetles, Fall Army Worms, dust at rate of 15 to 25 lbs. per acre according to infestation. For Stink Bugs use 20 to 25 lbs. per acre.

BEANS - Mexican Bean Beetle, dust at rate of 6 to 8 lbs. per acre according to infestation.

PEANUTS - Velvet Bean Caterpillar (Peanut Worm), the Fall Army Worm and Trips. Apply from 10 to 15 lbs. per acre depending upon size of plant and infestation. Repeat as often as needed.

CORN - Corn Fleahopper, Corn Earworm, Fall Army Worm, Mea Beetles, Leafhoppers, apply 10 to 20 lbs. per acre.

POTATO, TOMATO, SQUASH, PEPPER - Leafhoppers, Mea Beetles, Colorado Potato Beetle, apply 10 to 15 lbs. per acre.

CUCUMBER, CUCURBIT, SQUASH - Corn Earworm, Leafhoppers, Mea Beetles, Squash Bug, Corn Fleahopper, Stink Bug, etc. per acre. Avoid excessive applications.

Do not reuse empty containers. Destroy by burying with waste.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

MANUFACTURED BY

UNION CARBIDE CORP.

DAVIDSON, GEORGIA