

WARNING: DANGEROUS LIQUID

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN, REPEATED INHALATION OR SKIN CONTACT MAY, WITHOUT SYMPTOMS, PROGRESSIVELY INCREASE SUSCEPTIBILITY TO METHYL PARATHION POISONING.

Do not swallow or get in eyes, on skin, or on clothing. Do not breathe dust or vapors.; Wear clean, natural rubber gloves; clean waterproof protective clothing, and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for Methyl Parathion protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes, immediately flush with water for at least ten minutes and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water; wash clothing before re-use. In case of spillage, wash down with large amounts of water. Decontaminate areas and equipment by washing with a concentrated solution of alkali and detergent and rinse with water.

This product will kill fish and wildlife. Keep out of any body of water. Apply this product only as specified on this label.

Do not reuse empty container. Destroy by burying in a safe place.

ANTIDOTE AND FIRST AID TREATMENT

Atropine is the emergency antidote for methyl parathion poisoning. Consult your physician about obtaining an adequate supply of 1/100 grain atropine tablets for emergency use.

ACCEPTED

AUG 22 1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 275-21 SUBJECT
TO ATTACHED COMMENTS



POISON DANGER

Keep Out of Reach of Children



BERRIEN BRAND METHYL PARATHION ENDRIN

EMULSIFIABLE CONCENTRATE

ACTIVE INGREDIENTS

* Endrin: Hexachlorocyclopentadiene-endo, endo-dimethanonaphthalene. 18.30%

**O, O-Dimethyl O-p-Nitrophenyl thiophosphate 18.30%

INERT INGREDIENTS 63.40%

TOTAL 100.00%

This is a concentrated insecticide and must be diluted with water for use. Mix by adding to spray tank containing water with pump and agitator running. Before using study safety directions. Consult your physician about obtaining an adequate supply of 1/100 gr. Atropine tablets for emergency use.

DIRECTIONS FOR USE

COTTON - For the control of the following insects use 1 to 2 pints of Methyl Parathion Endrin per acre in enough water to give uniform coverage. Repeat as necessary. Thrips, Fleahoppers, Boll Weevils, Bollworm, Cabbage Looper, Brown cotton leaf-worm, Leaf perforator, Cotton Leafworm, Lygus bugs, Tarnished plant bug, Salt Marsh Caterpillar, Cutworms, Grasshoppers, certain Spider mites and Armyworms (up to 3rd instar). Do not treat within 15 days of hand picking. Do not graze dairy animals or animals being finished for slaughter on treated fields. Workers entering fields within 5 days after treatment should wear protective clothing.

*Equivalent to 1.6 lbs. Endrin per gallon as active material.

**Equivalent to 1.6 lbs. Methyl Parathion per gallon as active material.


BERRIEN PRODUCTS CO., INC.

NET CONTENTS

NASHVILLE, GEORGIA

_____ GAL.

U.S.D.A. REG. NO. 9275-21



Comment

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES
OF SUSPECTED POISONING

1 - Methyl Parathion is a cholinesterase inhibitor and treatment of severe poisoning should include the use of Atropine. 2 - Endrin produces central nervous system excitation which responds to the barbituric acid derivatives.

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the Methyl Parathion, remove all contaminated clothing, and wash skin with soap and running water, if the material gets into the eyes, wash immediately with running water for at least ten minutes. IF WARNING SYMPTOMS APPEAR: Administer two 1/100 grain atropine tablets immediately. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.