

ACCEPTED

12-8-67

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 9275-35

- May be fatal if swallowed
- Do not breathe spray mist
- Do not get in eyes, on skin, or on clothing
- Avoid contamination of food and feedstuffs
- Wash after handling
- Keep out of reach of children

WARNING OBSERVE THE FOLLOWING USE PRECAUTIONS

Do not apply this product on feed or forage to be fed to dairy animals or to livestock being finished.

Do not apply to fish and wildlife. Keep out of any body of water. Apply this product as directed on this label. Do not apply when weather conditions favor drift of spray from the treated area. Wash all equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying them in a non-crop area. (Keep out of smoke).

Do not allow to drift to areas occupied by unprotected humans, beneficial animals, or crops. Do not apply to feed and forage crops about to be harvested.

The Federal and State Administration has set official tolerances for certain pesticide residues. Do not apply that results on food and forage crops will not exceed established tolerances, and only at the recommended rates and do not apply closer to harvest than specified. Consult State Agricultural Extension Service Authorities for additional information, as the degree of control, and rate of applications needed will vary with local conditions.

BERRIEN STROBILINE 3 E

WARNING

Keep Out of Reach of Children

See other precautions on the left side panel.

EMULSIFIABLE LIQUID

AN AGRICULTURAL INSECTICIDE

(Contains 8 lbs. Strobilene per Gallon)

ACTIVE INGREDIENTS

Terpene Polychlorinated (60% Chlorine)	1.00
Nylone-Range Aromatic Hydrocarbon Solvent	1.00
Other Ingredients	1.00
TOTAL	3.00

*Solvent, consisting of polychlorinate of
Cyclopentane, Thione, and related terpene.

**DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME
PROTECT FROM TEMPERATURES BELOW 32 DEGREES F.**


*Reg. TM Hayden Newport GAL. NET

Berrien Products Co., Inc.

NASHVILLE, GEORGIA

U.S.D.A. REG. NO. 9275-35

Cameraman's Note
Poor Copy



DIRECTIONS FOR USE

Combine the needed amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing and spraying operations. Do not allow mixture to stand. If mixture does not mix readily but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine emulsifiable liquids with wettable powders in the same spray tank unless previous use of the materials being combined has proved them to be physically compatible.

RECOMMENDATIONS

(The dosage rates given below are to be incorporated with 25 gal. of water to provide adequate coverage.)

Use for the control of Beet Weevil, Bollworm, Cotton Leaf Worm, Cotton Fleahopper and Stink Bug apply 1 1/2 to 2 2/3 qt. per acre. For Cotton Fleahopper use 1 to 3 qt. per acre. If used apply 1/2 to 2/3 qt. per acre. For Cotton Leaf Fleahopper and Stink Bug apply 2 2/3 qt. per acre. Use the higher rate for heavy insect infestations and on rank-growing cotton, repeating at 5 day intervals. Do not apply after bolls open. Do not graze dairy animals or animals being held for slaughter on cotton or cotton trash.

If applied by aircraft use recommended amount in 2 gallons of water per acre.

CAUTION

This material is recommended for use on cotton only.