

**ACCEPTED**

**MAR 2 - 1968**

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. **9275-44**

# 15% FERMATE DUST

FOR BLUE MOLD CONTROL ON TOBACCO LEAF

### DIRECTIONS

Apply at rate of 5 to 10 lbs. per 100 square yds. Make first application when plants are the size of a dime, or when Blue Mold is first reported in the area and repeat applications at 3 to 4 day intervals until transplanted. See your State Agricultural Experiment Station or State Extension Service Authorities for further instructions.

### CAUTION

#### Keep Out of Reach of Children

CAUTION: May cause irritation of eyes, nose, throat and skin!! Avoid breathing of dust. Avoid contact with eyes, skin or clothing. In case of contact, flush with plenty of water; for eyes, get medical attention.

Keep out of any body of water. This product is toxic to fish.



ACTIVE INGREDIENTS	
Ferbam (Fermoxin)	15.00%
Inert Ingredients	85.00%
<b>TOTAL</b>	<b>100.00%</b>

\* Registered Trademark  
E. I. du Pont de Nemours & Co.  
Wilmington, Delaware

MANUFACTURED BY

**DU PONT DE NEMOURS & COMPANY**  
WILMINGTON, DELAWARE

Cameraman's Note

Poor Copy