

SANTIMINE
(Quaternary Ammonium Germicide)
TABLETS

LEFT LABEL

CAUTION

HARMFUL IF SWALLOWED.
DO NOT USE WITH SOAP
OR SYNTHETIC DETER-
GENTS. AVOID CON-
TAMINATION OF FOOD.
AVOID CONTACT WITH
SKIN AND EYES. IN
CASE OF CONTACT FLUSH
WITH PLENTY OF WATER.

PHENOL COEFFICIENTS

Salmonella typhosa 250
Staphylococcus aureus 450

Santimine Tablets fulfill
the criteria of appendix F,
the Grade 'A' Pasteurized
Milk Ordinance 1965 recom-
mendations of the U. S.
Public Health Service in
water up to 500 ppm hard-
ness by the Chambers method.

USDA Reg. No.

SANTIMINE SOLUTIONS are
non-flammable, non-volatile,
non-irritating, odorless,
non-corrosive to ordinary
surfaces and non-toxic in
recommended use dilutions.
SANTIMINE destroys odors of
bacterial origin.

ALWAYS FOLLOW YOUR
HEALTH DEPARTMENT
REGULATIONS

For the SANITIZATION of
DISHES-GLASSES
SILVER-UTENSILS

ACTIVE INGREDIENTS.....50%
n--Alkyl (98% Cl₂, 2% Cl₄)
dimethyl
1--naphthylmethyl ammonium
chlorides monohydrate
INERT INGREDIENTS.....50%

CAUTION

Keep Out of Reach of Children
See Side Panel for Additional Cautions
200 Tablets Net Weight 8 ozs.
Manufacturing Chemists

SANTIMINE COMPANY
Subsidiary of Texas
Gulf Industries

HOUSTON BALTIMORE
CHICAGO

ACCEPTED

MAY 24 1968

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PEST CONTROL REGISTERED
ED UNDER NO. 9279-2 SUBJECT
TO ATTACHED COMMENTS

RIGHT LABEL

PREPARE SANTIMINE SOLU
WITH HOT WATER (clean
drinking water) AND STIR
WELL BEFORE USING. For
sanitizing Dishes, Glasses,
Silver and Utensils in
Restaurants, Taverns, etc.

1. Scrape and prewash
utensils and glasses
whenever possible.
2. Wash with a good deter-
gent or compatible
cleaner in first sink
compartment.
3. Rinse with clean drink-
ing water in second sink
compartment.
4. Sanitize in a solution of
1 tablet to 3 quarts of
water (200 ppm) in third
sink compartment.
Immerse all utensils for
at least two minutes or
for contact time specified
by governing sanitary code.
5. Rinse with clean drinking
water.
6. Place sanitized utensils
on a rack or drainboard
to air-dry.

For General Disinfection of
Sanitary Equipment, Restrooms,
Floors, Refrigerators or Cab-
inets and for Deodorizing
Garbage Pails, Coffee Urns,
Restroom Equipment, etc.,
clean and Rinse surface to be
treated. Then apply freely a
solution of 4 tablets per
gallon of water. Rinse all
utensils or surfaces coming in
contact with food with clean
drinking water before use.

ACCEPTED

6-18-73

UNDER THE FEDERAL INSECTICIDES
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 9279-3

SANTIMINE 150

(QUATERNARY AMMONIUM GERMICIDE)
TABLETS

Use 1 Tablet Per Gallon of Water
FOR THE SANITIZATION
OF DISHES - GLASSES - UTENSILS

ACTIVE INGREDIENTS 50%
n-Alkyl (95% C14, 3% C12, 2% C16) dimethyl benzyl
ammonium chlorides monohydrate.
INERT INGREDIENTS 50%

DANGER

KEEP OUT OF REACH OF CHILDREN.

See First Aid and Other precautions on left panel.
EPA Reg. No. 9279-3

150 Tablets

Net Weight: 8 Ounces

SANTIMINE DIVISION

Texas Gulf Industries, Inc.
2912 Pulaski Hwy., Baltimore, Md. 21224
(301) 235-8473

ACCURATE - CONVENIENT EFFECTIVE

This product fulfills the criteria of Appendix F of the Grade A Pasteurized Milk Ordinance 1965 Recommendations of the U.S. Public Health Service in waters up to 500 ppm of hardness calculated as CaCO₃ when tested by the ACAC Germicidal and Detergent Sanitizers-Official Method.

DANGER - CORROSIVE. Keep out of reach of children. Tablets may be harmful or fatal if swallowed. Causes eye damage. May cause skin irritation. Wash hands thoroughly after handling. Avoid prolonged skin contact. Do not get in eyes. Avoid contamination of food. **FIRST AID** In case of eye contact, immediately flush eyes with plenty of water for at least 15 minutes. Get prompt medical attention. If tablets are swallowed, drink promptly a large quantity of milk, egg whites or gelatin solution. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard.

**ALWAYS FOLLOW YOUR
HEALTH DEPARTMENT
REGULATIONS W-11-572.**

DIRECTIONS

When preparing sanitizing solutions, allow several minutes for tablets to dissolve in water. Stir well before using.

For sanitizing Dishes, Glasses, and Utensils in Restaurants, Taverns.

1. Scrape and prewash utensils and glasses whenever possible.
2. Wash with a good detergent or compatible cleaner in first sink compartment.
3. Rinse with clean water in second sink compartment.
4. Sanitize in a solution of 1 tablet per gallon of water (200 ppm) in third sink compartment. Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drain-board to air dry.

Tablet sanitizing solutions of 200 ppm concentration may be safely used on food processing equipment and utensils and on other food contact articles as specified under Subpart F of the Food Additives Amendment of the Federal Food, Drug and Cosmetic Act.