

ACCURATE-CONVENIENT-EFFECTIVE

This product fulfills the criteria of appendix F of the Grade A Pasteurized Milk Ordinance 1945 Recommendations of the U.S. Public Health Service in waters up to 500 ppm of hardness calculated as CaCO₃, when tested by the ACAC Germicidal and Detergent Sanitizer-Official Method

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER

Keep Out of Reach of Children

Causes severe eye and skin damage. Do not get in eyes, on skin or clothing. Harmful if swallowed. Wear goggles or face shield and rubber gloves when handling. Avoid contamination of food. Remove and wash contaminated clothing.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink egg whites, gelatin solution or, if these are not available, drink large quantities of water and call a physician.

SANTIMINE 150

(QUATERNARY AMMONIUM GERMICIDE) TABLETS
USE 1 TABLET PER GALLON OF WATER
FOR THE SANITATION OF DISHES-GLASSES-UTENSILS

ACTIVE INGREDIENTS 50%
N-ARYLIDIPYCLIA, 3% C12, 2% C18 dimethyl benzyl ammonium chlorides monohydrate
INERT INGREDIENTS 50%

KEEP OUT OF REACH OF CHILDREN

DANGER

See side panel for additional precautionary statements

EPA Reg. No. 9275-3

EPA Est. 9277 - MD - 3

150 Tablets

Net Weight 8 ounces

TEXAS GULF INDUSTRIES, INC.
725 N. Haven St., Baltimore, Md. 21205
(301) 327-0695

NOT REVIEWED

In Accordance with 16 CFR 302-2.
Based on Data Labeling Dated

10F1
PM 31
9279-3



DIRECTIONS FOR USE

When preparing sanitizing solution, allow several minutes for tablets to dissolve in water. Stir well before using for sanitizing dishes, glasses and utensils in restaurants, etc.

- 1 - Scrape and polish dishes and glasses when ever possible
- 2 - Wash with a good detergent or compatible cleaner in first sink compartment
- 3 - Rinse with clean water in second sink compartment
- 4 - Sanitize in a solution of 1 TABLET PER GALLON (100 ppm) in third sink compartment. Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code
- 5 - Place sanitized utensils on a rack or drain board to air dry

Tablet sanitizing solutions of 100 ppm concentration may be safely used on food processing equipment and utensils and on other food contact articles as specified under subpart F of the Food Additives Amendment of the Federal Food, Drug and Cosmetic Act

- Disposal -

Rinse empty container thoroughly with water and discard

It is a violation of federal law to use this product in a manner inconsistent with its labeling

ALWAYS FOLLOW YOUR HEALTH DEPARTMENT REGULATIONS
M - 11 - 573

BEST AVAILABLE COPY