



WOOD STAIN EVER WOOD SURFACES


YELLOW
GREEN

Reg. No. 9284-5

**FOR ANYTHING
MADE OF WOOD**



DOOR JAMBS AND SILLS



FENCES AND POSTS



WINDOW FRAMES AND SILLS

CONTAINS PENTACHLOROPHENOL

Harmful or fatal if Swallowed or Absorbed Through the Skin. If Swallowed, Induce Vomiting and Call a Physician Immediately

WARNING! COMBUSTIBLE

FIRE Avoid exposing the contents of this can or the freshly treated lumber to open flame. Treat in well-ventilated areas.

SKIN During the application avoid contact with the skin. Wear rubber gloves and protective clothing when handling the freshly treated lumber.

EYES If in contact with eyes, flush for 15 minutes with running water. Call a physician immediately.

BREATHING Use a mask when spraying. Do not spray for extended periods in a closed area.


HAZARDS Keep freshly treated articles away from plants, food stuffs, and children. Do not get in eyes, on skin, or on clothing.

ACTIVE INGREDIENTS		79.50%
PENTACHLOROPHENOL	4.50%	
OTHER CHLORINATED PHENOLS	0.50%	
PETROLEUM SOLVENTS	74.50%	
Total	79.50%	

INERT INGREDIENTS		20.50%
RESIN SOLIDS	6.50%	
OXIDE TINTING PIGMENTS (Titanium Dioxide, Chrome Oxide or Iron Oxide)	4.40%	
HEAT TREATED LINSEED OIL	9.60%	
Total	20.50%	

100.00%

This product is toxic to fish and wildlife. Do not contaminate water by cleaning of equipment or disposal of wastes.



**BRUSH IT!
DIP IT!
SPRAY IT!
ROLL IT!**

WATERLOX CHEMICAL & COATINGS CORP.

WOOD ROT

FOR ADDITIONAL WARNINGS.

ALLOWED. KEEP OUT OF REACH OF CHILDREN.

CHLORINATED PHENOLS 0.50%, PETROLEUM SOLVENTS 74.50%, INERT INGREDIENTS 20.50%


U. S. MEASURE



PATIO - FURNITURE



ROUGH CUT SIDING



BOAT BILGE

RIEDEL'S SUPERCOSAN™

FUNGICIDE-MITICIDE

READ THIS LABEL CAREFULLY

U.S.D.A. Registration No. 9288-1

DIRECTIONS FOR USE

Measure needed amount into bucket of water, stir to make thin slurry. Add to tank when part full, and before other materials are added. Add wetting agents last to avoid foaming. Keep agitated.

APPLES, PEARS: Powdery Mildew, Scab, Blister Mites, Silver Mites, Red Spider spp. 2-6 pounds per 100 gals. water. Do not apply to D'anjou Pears. Apply pre bloom and petal fall. Apply with cover sprays throughout the season.

ALMONDS, CHERRIES, PLUMS, PRUNES: Brown Rot, Powdery Mildew, Leaf Spot, Silver Mites, Flat Mites, Red Spider spp. 2-6 pounds per 100 gals. of water. Apply at bloom or early petal fall, repeat as necessary, usually 10-14 days, or after a period of wet weather.

PEACHES, NECTARINES: Brown Rot, Powdery Mildew, Silver Mites, Red Spider spp. 2-6 pounds per 100 gals. water. Apply at bloom or early petal fall, repeat as necessary, usually 10-14 days, or after a period of wet weather.

CITRUS: Flat Mite, Silver Mites, Red Spider spp. 2-6 pounds per 100 gals. of water. Apply in any spray during season — NOT CONTAINING OIL or following oil by 14 days ! ! ! !

OLIVES: Bud Mites, apply 2 lbs. per 100 gallons of water.

AVOCADO: Brown Mite, apply 2-6 lbs. per 100 gals. of water. Apply as needed.

GRAPES: Powdery Mildew, Bud Mite, Blister Mite, Red Spider spp. 2 to 6 pounds per 100 gals. of water. Apply in late dormant, or with cutworm spray, early shoot period and thru season on 14 day schedule, or with pesticides.

BLACKBERRY: (Rubis Spp): Red Berry Mites (Eriophyes Gracilis) apply 3-5 lbs. per 100 gals. water. Start in early bud swell, repeat applications as necessary, with or without pesticides.


RASPBERRY (Rubis Spp): Powdery Mildew (Sphaerotheca Humuli) spray each week, starting at first bloom-color with 3-5 lbs. per 100 gals. of water. Continue until all fruit is set, then repeat as weather requires.

Manufactured by Riedel-De Haën AG
D 3016 Seelze-Hannover, Germany



TM COSAN — Trademark of Riedel-De Haën AG Int. Reg. B.I.R.P.I. No. 152354

ACTIVE INGREDIENTS:
Micronized Sulfur
INERT INGREDIENTS:



STRAWBERRIES: Powdery Mildew, Red Spider spp. apply 3-5 lbs. per 100 gallons leaf stage; repeat applications with or without pesticides.

TOMATOES: Russet Mite, apply 3-5 lbs. per acre. Thorough coverage is required of water either by air or ground rig.

BEANS, PEAS, COLE CROPS, PEPPERS, POTATOES: Powdery Mildew, Red Spider spp. apply 3-5 lbs. per acre in 15-50 gals. of water by aircraft or ground rig. Start in early leaf stage, repeat every 10-14 days with or without pesticides, or immediately after a rain.

HOPS: Red Spider (spp) apply 2-6 lbs. per 100 gals. water. Start in early leaf stage, repeat applications as needed, usually 10-14 days.

SUGAR BEETS: Red Spider Spp. apply 3-5 lbs. per acre in 15-50 gal. water by ground rig. Apply as needed with or without pesticides.

COTTON, SEED ALFALFA: Red Spider spp. Apply 3-5 pounds per acre in 25-50 gallons of water by aircraft or ground rig.

PEANUTS: Powdery Mildew, Red Spider Mite, Leaf Spot. Apply 3 to 5 lbs. per acre in 15-50 gallons of water by aircraft or ground rig. Start in early leaf stage. Repeat applications as necessary, usually 10 to 14 days.

WARNING

KEEP OUT OF REACH OF CHILDREN

Do not use on sulfur sensitive plants. Sulfur may burn foliage very severely if applied excessively high. Do not apply with or following an oil application. Do not apply oil for 14 days following treatment.

NET WT. 56 LBS. — KEEP DRY

Conditions of Sale

Follow directions carefully. Timing and method of application are subject to local conditions, mixtures with other chemicals not specifically recommended. Buyer assumes all risks of use, storage and handling of this product. Conditions of sale in accordance with directions given herewith.



Imported by Haley & Kruger Co., Inc.
Reedley, California 93654