

9289-1

REGISTERED
4-10-67
SIDE
ACT
REGISTER
9289-1

APICIDE

Net Weight 7 Ounces

ACTIVE INGREDIENTS	3%
Technical chlordane*	2.5%
*(Equivalent to 1.5% Octachloro-4, 7-methanotetrahydroindane and 1% of related compounds)	
O,O-diethyl O-(2-isopropyl-6-methyl-6-pyrimidinyl) phosphorothioate also known as "Diazinon"	0.5%
*Diazinon is a registered Trademark of Geigy Chemical Corporation for its patented insecticide, U. S. Patent No. 2, 754, 343.	
INERT INGREDIENTS	97%
Total	100%

The user assumes all risk of use or handling of APICIDE whether or not in accordance with directions or suggestions.

DIRECTIONS

Locate the brood nest, or center of the colony, if possible by listening with your ear to the wall (inside or out). Where the buzzing and clicking sound is the loudest is generally the area of the brood comb and bees.

Drill a 1/4" hole into that section of the wall from either the inside or outside and push the bit on through to clear a hole into the brood area.

Insert the nozzle of the APICIDE dispenser into the 1/4" hole and squeeze the dispenser, thus pumping the dust into the brood area. Continue squeezing until most of the insecticide dust is pumped into the wall cavity. If dust packs in nozzle, remove and shake dispenser to loosen dust. Allow 24 hours for APICIDE to completely destroy the colony after which the 1/4" hole and the outside bee entrance can be permanently sealed with wood putty or some other similar impervious material.

If it is impossible to drill a hole into the brood nest, apply APICIDE heavily around the outside entrance to the colony. Also, pump a large quantity of APICIDE into the entrance. This method will take longer to destroy the colony and may require two or three applications in order to do so.

APICIDE will also destroy wasps and hornets but heavy protective clothing including veil and gloves should be worn in applying APICIDE to the nest. Otherwise, the hornets will attack savagely before the APICIDE can be applied effectively.

WARNING: Do not breathe APICIDE, nor allow it to contaminate food or clothing. Harmful if swallowed. Avoid contact with skin. Wash hands thoroughly with soap and water after using. Stand upwind while pumping APICIDE into the entrance of the colony. **Do not eat any honey taken out of the wall after bees have been destroyed with APICIDE.** Empty container should be destroyed by burning (outdoors) or burying a foot deep.

KEEP OUT OF REACH OF CHILDREN

Formulated and packed by the Professional Insect Control Co. of Medina, Ohio for:

THE A. I. ROOT CO. Medina, Ohio

100 lbs. Net Weight

Fletcher's Shade Tobacco Mixture WITH CHLORDANE

EPA Reg. No. 9355-1

(10 lbs. 40% Chlordane D.B. Per Ton)

This fertilizer contains .20 lbs. of technical chlordane per 100 lbs. of fertilizer.

Pesticide Ingredient Statement

ACTIVE INGREDIENTS	BY WT.
Technical Chlordane	0.20%
Inert Ingredients	99.80%
TOTAL	100.00%

*Equivalent to .12% Octachloro-4, 7 Methanotetra hydroindane and .08% of related compounds. Derived from addition of 10 lbs. 40% chlordane dust base per ton.

DIRECTIONS

Tobacco soil treatment — Against wireworms and cutworms, apply as a pre-plant broadcast application at the rate of 2,000 lbs. per acre. Thoroughly disc 4 to 6 inches deep in soil.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid inhalation, bathe thoroughly, do not leave on skin. Avoid contamination of feed and food stuffs. Do not apply to areas occupied by unprotected humans or beneficial animals. This product is toxic to fish and wild life. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label. Do not use on pastures for dairy animals. Do not use on pastures for meat animals if to be grazed within 90 days before slaughter.

Use this product only for recommended purposes and at recommended dosage.

Do not reuse empty container. Destroy when empty.

Manufactured by:
FLETCHER COMPANY, INC.
GREENSBORO, FLORIDA
EPA Reg. No. 9355-1

**Do not use on pastures
for dairy animals.**

**Do not use on pastures
for meat animals
if to be grazed
within 90 days
before slaughter.**

