

BEST SUPER COSAN

Micronized Wettable Sulfur - A Product of West Germany

You Get Results With

BEST
FERTILIZERS - CHEMICALS

Read This Label Carefully

ACTIVE INGREDIENTS

Micronized Sulfur -----

INERT INGREDIENTS: -----

TOTAL

BY WT.

80%

20%

100%

WARNING

KEEP OUT OF REACH OF CHILDREN

U. S. DA. Reg No. _____

DIRECTIONS FOR USE

Measure needed amount into bucket of water, stir to make thin slurry. Add to tank when part full, and before other materials are added. Add wetting agents last to avoid foaming.

APPLES, PEARS: Powdery Mildew, Scab, Blister Mites, Silver Mites, Red Spider spp. 2-6 pounds per 100 gals water. Do not apply to D'anjou Pears. Apply pre bloom and petal fall. Apply with cover sprays throughout the season.

ALMONDS, CHERRIES, PLUMS, PRUNES: Brown Rot, Powdery Mildew, Leaf Spot, Silver Mites, Flat Mites, Red Spider spp. 2-6 pounds per 100 gals of water. Apply at bloom or early petal fall, repeat as necessary, usually 10-14 days, or after a period of wet weather.

PEACHES, NECTARINES: Brown Rot, Powdery Mildew, Silver Mites, Red Spider spp. 2-6 pounds per 100 gals water. Apply at bloom or early petal fall, repeat as necessary, usually 10 - 14 days, or after a period of wet weather.

CITRUS: Flat Mite, Silver Mites, Red Spider spp. 2-6 pounds per 100 gals of water. Apply in any spray during season--NOT CONTAINING OIL or following oil by 14 days!!!!

GRAPES: Powdery Mildew, Bud Mite, Blister Mite, Red Spider spp. 2 to 6 pounds per 100 gals of water. Apply in late dormant, or with cutworm spray, early shoot period and thru season on 14 day schedule, or with pesticides.

BEANS, PEAS, COLE CROPS, PEPPERS: Powdery Mildew, Septoria Leaf Spot, Red Spider spp. 3-5 pounds per acre in 15-50 gals of water by aircraft or ground rig. Start in early leaf stage. Repeat application every 10-14 days with or without pesticides, or immediately after a rain.


COTTON, SEED ALFALFA: Red Spider spp. Apply 3-5 pounds per acre in 15-50 gallons of water by air craft or ground rig.

STRAWBERRIES: Powdery Mildew, Red spider spp. 3-5 pounds per 100 gals of water. Start in early leaf stage; repeat applications with or without pesticides.

TOMATOES: Russet Mite. Apply 3-5 pounds per acre. Thorough coverage is required, 20 to 50 gallons of water either by air or ground rig.

Do not use on sulfur sensitive plants. Sulfur may burn when temperature is excessively high. Do not apply with or following an oil application for 14 days. Do not apply oil for 14 days following treatment.

483:6/67



BEST DOCUMENT AVAILABLE

THE BEST FERTILIZERS CO.

DIVISION OF OCCIDENTAL PETROLEUM CORPORATION

P. O. BOX 198 • LATHROP, CALIFORNIA 95330 • PHONE (415) 858-2511

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than an officer of Seller, is authorized to make any other warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of purchase price.

Best Super Cosan

9288-1
08/14/1967

40