

Riedel's SUPERCOSAN™

FUNGICIDE - MITICIDE

73

Read This Label Carefully

U.S.D.A. Registration No. 9288-1

ACTIVE INGREDIENTS:

Micronized Sulfur

BY WT.

80 %

INERT INGREDIENTS:

20 %

TOTAL 100 %

DIRECTIONS FOR USE

Measure needed amount into bucket of water, stir to make thin slurry. Add to tank when part full, and before other materials are added. Add wetting agents last to avoid foaming.

APPLES, PEARS: Powdery Mildew, Scab, Blister Mites, Silver Mites, Red Spider spp. 2—6 pounds per 100 gals water. Do not apply to D'anjou Pears. Apply pre bloom and petal fall. Apply with cover sprays throughout the season.

ALMONDS, CHERRIES, PLUMS, PRUNES: Brown Rot, Powdery Mildew, Leaf Spot, Silver Mites, Flat Mites, Red Spider spp. 2—6 pounds per 100 gals of water. Apply at bloom or early petal fall, repeat as necessary, usually 10—14 days, or after a period of wet weather.

PEACHES, NECTARINES: Brown Rot, Powdery Mildew, Silver Mites, Red Spider spp. 2—6 pounds per 100 gals water. Apply at bloom or early petal fall, repeat as necessary, usually 10—14 days, or after a period of wet weather.

CITRUS: Flat Mite, Silver Mites, Red Spider spp. 2—6 pounds per 100 gals of water. Apply in any spray during season — **NOT CONTAINING OIL** or following oil by 14 days!!!!

GRAPES: Powdery Mildew, Bud Mite, Blister Mite, Red Spider spp. 2 to 6 pounds per 100 gals of water. Apply in late dormant, or with cutworm spray, early shoot period and thru season on 14 day schedule, or with pesticides.

Manufactured by Riedel-de Haën AG
D 3016 Seelze-Hannover, Germany

ACCEPTED

4/30/68

IN THE FEDERAL INSECTICIDE
FUNGICIDE AND MITICIDE ACT
FOR ECONOMIC PEST CONTROL, REGISTERED
UNDER NO. 9288-1 SUBJECT
TOMATOES

MORE THAN 150 YEARS



Riedel-de Haën AG Int. Reg. B.I.R.P.I. No. 152354

BEANS, PEAS, COLE CROPS, PEPPERS: Powdery Mildew, Red Spider spp. 3—5 pounds per acre in 15—50 gals of water by aircraft or ground rig. Start in early leaf stage. Repeat application every 10—14 days with or without pesticides, or immediately after a rain.

COTTON, SEED ALFALFA: Red Spider spp. Apply 3—5 pounds per acre in 25—50 gallons of water by air craft or ground rig.

STRAWBERRIES: Powdery Mildew, Red spider spp. 3—5 pounds per 100-gals of water. Start in early leaf stage; repeat applications with or without pesticides.

TOMATOES: Russet Mite, Apply 3—5 pounds per acre. Thorough coverage is required, 25 to 50 gallons of water either by air or ground rig.

WARNING

KEEP OUT OF REACH OF CHILDREN

Do not use on sulfur sensitive plants. Sulfur may burn foliage when temperature is excessively high. Do not apply with or following an oil application for 14 days. Do not apply oil for 14 days following treatment.

NET WT. 56 LBS.

KEEP DRY

Conditions of Sale

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Imported by Haley & Kruger Co., Inc.
Reedley, California 93654

BEST DOCUMENT AVAILABLE



50

9288-1

04/30/1968

1/2

RIEDEL'S "SUPERCOSAN" FUNGICIDE-MITICIDE

ACCEPTED

4/30/68

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND MITICIDE ACT
FOR ECONOMIC PESTICIDE REGISTRATION
ED UNDER NO. 9288-1. SUBJECT
TO ATTACHED COMMENTS.

22

Read This Label Carefully

DIRECTIONS FOR USE

Measure needed amount into bucket of water, stir to make thin slurry. Add to tank when part full, and before other materials are added. Add wetting agents last to avoid foaming.

APPLES, PEARS: Powdery Mildew, Scab, Blister Mites, Silver Mites, Red Spider spp. 2-6 pounds per 100 gals water. Do not apply to D'anjou Pears. Apply pre bloom and petal fall. Apply with cover sprays throughout the season.

ALMONDS, CHERRIES, PLUMS, PRUNES: Brown Rot, Powdery Mildew, Leaf Spot, Silver Mites, Flat Mites, Red Spider spp. 2-6 pounds per 100 gals of water. Apply at bloom or early petal fall, repeat as necessary, usually 10-14 days, or after a period of wet weather.

PEACHES, NECTARINES: Brown Rot, Powdery Mildew, Silver Mites, Red Spider spp. 2-6 pounds per 100 gals water. Apply at bloom or early petal fall, repeat as necessary, usually 10 - 14 days, or after a period of wet weather.

CITRUS: Flat Mite, Silver Mites, Red Spider spp. 2-6 pounds per 100 gals of water. Apply in any spray during season--NOT CONTAINING OIL or following oil by 14 days!!!!

GRAPES: Powdery Mildew, Bud Mite, Blister Mite, Red Spider spp. 2 to 6 pounds per 100 gals of water. Apply in late dormant, or with cutworm spray, early shoot period and thru season on 14 day schedule, or with pesticides.

BEANS, PEAS, COLE CROPS, PEPPERS: Powdery Mildew, Red Spider spp. 3-5 pounds per acre in 15-50 gals of water by aircraft or ground rig. Start in early leaf stage. Repeat application every 10-14 days with or without pesticides, or immediately after a rain.

COTTON, SEED ALFALFA: Red Spider spp. Apply 3-5 pounds per acre in 15-50 gallons of water by air craft or ground rig.

STRAWBERRIES: Powdery Mildew, Red spider spp. 3-5 pounds per 100 gals of water. Start in early leaf stage; repeat applications with or without pesticides.

TOMATOES: Russet Mite. Apply 3-5 pounds per acre. Thorough coverage is required, 15 to 30 gallons of water either by air or ground rig.

ACTIVE INGREDIENTS:	BY WT.
Micronized Sulfur-----	80%
INERT INGREDIENTS:-----	20%
TOTAL	100%

U.S. D. A. Registration No. 9288-1

* Cosan=European Trademark of Riedel-de Haën A G

WARNING

KEEP OUT OF REACH OF CHILDREN

Do not use on sulfur sensitive plants. Sulfur may burn when temperature is excessively high. Do not apply with or following an oil application for 14 days. Do not apply oil for 14 days following treatment.

NET WT. 56 LBS.

KEEP DRY

Manufactured By Riedel-De Haën AG
Seelze - Hannover, Germany



Imported by Haley & Kruger Co., Inc.
Reedley, California

CONDITIONS OF SALE

BEST DOCUMENT AVAILABLE

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

PEANUTS: Powdery Mildew, Red Spider spp. 3-5 pounds per acre in 15-50 gals. water by aircraft or ground rig. Start in early leaf stage. Repeat applications as necessary, usually 10-14 days.

2/2