

Riedel's SUPERCOSAN™

FUNGICIDE - MITICIDE

Read This Label Carefully

U.S.D.A. Registration No. 9288-1

ACTIVE INGREDIENTS:

Micronized Sulfur

BY WT.

80 %

INERT INGREDIENTS:

20 %

TOTAL 100 %

DIRECTIONS FOR USE

Measure needed amount into bucket of water, stir to make thin slurry. Add to tank when part full, and before other materials are added. Add wetting agents last to avoid foaming.

APPLES, PEARS: Powdery Mildew, Scab, Blister Mites, Silver Mites, Red Spider spp. 2—6 pounds per 100 gals water. Do not apply to D'anjou Pears. Apply pre bloom and petal fall. Apply with cover sprays throughout the season.

ALMONDS, CHERRIES, PLUMS, PRUNES: Brown Rot, Powdery Mildew, Leaf Spot, Silver Mites, Flat Mites, Red Spider spp. 2—6 pounds per 100 gals of water. Apply at bloom or early petal fall, repeat as necessary, usually 10—14 days, or after a period of wet weather.

PEACHES, NECTARINES: Brown Rot, Powdery Mildew, Silver Mites, Red Spider spp. 2—6 pounds per 100 gals water. Apply at bloom or early petal fall, repeat as necessary, usually 10—14 days, or after a period of wet weather.

CITRUS: Flat Mite, Silver Mites, Red Spider spp. 2—6 pounds per 100 gals of water. Apply in any spray during season — NOT CONTAINING OIL or following oil by 14 days!!!!

GRAPES: Powdery Mildew, Bud Mite, Blister Mite, Red Spider spp. 2 to 6 pounds per 100 gals of water. Apply in late dormant, or with cutworm spray, early shoot period and thru season on 14 day schedule, or with pesticides.

Manufactured by Riedel-de Haën AG

D 3013 Seelze-Hannover, Germany

Leaf Spot: 2-6 lbs. per 100 gals water by aircraft or ground rig. Start in early leaf stage, repeat applications as necessary, usually 10-14 days.



OLIVES: Bud mites. Apply 2 lbs. per 100 gallons of water. Spray as necessary.

TM COSAN — Trademark of Riedel-de Haën AG Int. Reg. B.I.R.P.I. No. 152354

HOPS: Red Spider spp. 2-6 lbs. per 100 Gals. of water. Start in early leaf stage, repeat applications as necessary, usually 10-14 days.

Potatoes, [redacted] Powdery Mildew, Red Spider spp. 3—5 pounds per acre in 15—50 gals of water by aircraft or ground rig. Start in early leaf stage. Repeat application every 10—14 days with or without pesticides, or immediately after a rain.

COTTON, SEED ALFALFA: Red Spider spp. Apply 3—5 pounds per acre in 25—50 gallons of water by air craft or ground rig.

STRAWBERRIES: Powdery Mildew, Red spider spp. 3—5 pounds per 100 gals of water. Start in early leaf stage; repeat applications with or without pesticides.

TOMATOES: Russet Mite, Apply 3—5 pounds per acre. Thorough coverage is required 25 to 50 gallons of water either by air or ground rig.

WARNING

KEEP OUT OF REACH OF CHILDREN

Do not use on sulfur sensitive plants. Sulfur may burn foliage when temperature is excessively high. Do not apply with or following an oil application for 14 days. Do not apply oil for 14 days following treatment.

NET WT. 56 LBS.


KEEP DRY

Conditions of Sale

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Imported by Haley & Kruger Co., Inc.

Reedley, California 93654



BEST DOCUMENT AVAILABLE



Best copy available

9288-1

09/03/1968

1/2

60

SUGAR BEETS: Red spider spp. Apply 3-5 lbs. per acre in 15-50 gals. of water by aircraft or ground rig. Apply as needed with or without pesticides.

BLACKBERRY: (Rubis spp) Red Berry mites (*Eriophyes fructilis*). Apply 3-5 lbs. per 100 gals. of water. Start in early bud swell, repeat applications as necessary, with or without pesticides.

RASPBERRY (Rubis spp): Powdery mildew (*sphaerotheca humuli*). Spray each week, starting at first bloom-color with 3-5 lbs. per 100 gals. of water, continue until all fruit is set, then repeat as weather requires.

ACCEPTED

9/3/68

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND MITICIDE ACT
FOR EACH OF THE FOLLOWING REGISTERED
ED 101-14 NO. 101-14... SUBJECT
TO ATTACHED COMMENTS.

BEST DOCUMENT AVAILABLE