

RIEDEL'S SUPERCOSAN™

FUNGICIDE-MITICIDE

READ THIS LABEL CAREFULLY

U.S.D.A. Registration No. 9288-1

DIRECTIONS FOR USE

Measure needed amount into bucket of water, stir to make thin slurry. Add to tank when part full, and before other materials are added. Add wetting agents last to avoid foaming.

APPLES, PEARS: Powdery Mildew, Scab, Blister Mites, Silver Mites, Red Spider spp. 2-6 pounds per 100 gals. water. Do not apply to D'anjou Pears. Apply pre bloom and petal fall. Apply with cover sprays throughout the season.

ALMONDS, CHERRIES, PLUMS, PRUNES: Brown Rot, Powdery Mildew, Leaf Spot, Silver Mites, Flat Mites, Red Spider spp. 2-6 pounds per 100 gals. of water. Apply at bloom or early petal fall, repeat as necessary, usually 10-14 days, or after a period of wet weather.

PEACHES, NECTARINES: Brown Rot, Powdery Mildew, Silver Mites, Red Spider spp. 2-6 pounds per 100 gals. water. Apply at bloom or early petal fall, repeat as necessary, usually 10-14 days, or after a period of wet weather.

CITRUS: Flat Mite, Silver Mites, Red Spider spp. 2-6 pounds per 100 gals. of water. Apply in any spray during season — **NOT CONTAINING OIL** or following oil by 14 days ! ! ! !

GRAPES: Powdery Mildew, Bud Mite, Blister Mite, Red Spider spp. 2 to 6 pounds per 100 gals. of water. Apply in late dormant, or with cutworm spray, early shoot period and thru season on 14 day schedule, or with pesticides.

POTATOES:

BEANS, PEAS, COLE CROPS, PEPPERS: Powdery Mildew, Red Spider spp. 3-5 pounds per acre in 15-50 gals. of water by aircraft or ground rig. Start in early leaf stage. Repeat application every 10-14 days with or without pesticides, or immediately after a rain.

COTTON, SEED ALFALFA: Red Spider spp. Apply 3-5 pounds per acre in 25-50 gallons of water by air craft or ground rig.

STRAWBERRIES: Powdery Mildew, Red spider spp. 3-5 pounds per 100 gals. of water. Start in early leaf stage; repeat applications with or without pesticides.

TOMATOES: Russet Mite, Apply 3-5 pounds per acre. Thorough coverage is required, 25 to 50 gallons of water either by air or ground rig.

OLIVES: Bud Mites, apply 2 lbs. per 100 gallons of water.

PEACHES, NECTARINES: Brown Rot, Powdery Mildew, Silver Mites, Red Spider, apply 2-6 lbs. per 100 gallons water. Apply at bloom or early petal fall, repeat as necessary, usually 10-14 days, or after a period of wet weather.

Manufactured by Riedel-de Haën AG
D 3016 Seelze-Hannover, Germany



OVER 150 YEARS

TM COSAN — Trademark of Riedel-de Haën AG Int. Reg. B.I.R.P.I. No. 152354

ACTIVE INGREDIENTS:

Micronized Sulfur

INERT INGREDIENTS:

BY WT.

80%

20%

TOTAL 100%

STRAWBERRIES: Powdery Mildew, Red Spider, apply 3-5 lbs. per 100 gallons of water. Start in early leaf stage; repeat applications with or without pesticides.

TOMATOES: Russet Mite, apply 3-5 lbs. per acre. Thorough coverage is required, 15 to 50 gallons of water either by air or ground rig.

BLACKBERRY: (Rubis Spp): Red Berry Mites (Eriophyes Gracilis) apply 3-5 lbs. per 100 gals. water. Start in early bud swell, repeat applications as necessary, with or without pesticides.

RASPBERRY (Rubis Spp): Powdery Mildew (Sphaerotheca Humuli) spray each week, starting at first bloom-color with 3-5 lbs. per 100 gals. of water. Continue until all fruit is set, then repeat as weather requires.

SUGAR BEETS: Red Spider Spp. apply 3-5 lbs. per acre in 15-50 gal. water, by aircraft or ground rig. Apply as needed with or without pesticides.

HOPS: Red Spider (spp) apply 2-6 lbs. per 100 gals. water. Start in early leaf stage, repeat applications as needed, usually 10-14 days.

PEANUTS: Powdery Mildew, Red Spider Mite. Apply 3 to 5 lbs. per acre in 15 to 50 gallons of water by aircraft or ground rig. Start in early leaf stage. Repeat applications as necessary, usually 10 to 14 days.

WARNING

KEEP OUT OF REACH OF CHILDREN

Do not use on sulfur sensitive plants. Sulfur may burn foliage when temperature is excessively high. Do not apply with or following an oil application for 14 days. Do not apply oil for 14 days following treatment.

NET WT. 56 LBS. — KEEP DRY

Conditions of Sale


Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

ACCEPTED

NOV 12 1968

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 9288-1 SUBJECT
TO ATTACHED COMMENTS.

BEST DOCUMENT AVAILABLE



Imported by Haley & Kruger Co., Inc.
Redley, California 93654

64

9288-1

11/12/1968