RIEDELS SUPERCUSAN'

FUNGICIDE-MITICIDE

READ THIS LABEL CAREFULLY

U.S.D.A. Registration No. 9288-1

DIRECTIONS FOR USE

Measure needed amount into bucket of water, stir to make thin sturry. Add to tank when part full, and before other materials are added. Add wetting agents last to avoid foaming. Keep agitated.

APPLES: PEARS: Powdery Mildew, Scab, Blister Mites, Silver Mites, Red Spider spp. 2-6 pounds per 100 gale, water. Do not apply to D'anjou Pears. Apply pre bloom and petal fall. Apply with cover sprays throughout the season.

ALMONDS, CHERRIES, PLUMS, PRUNES: Brown Rot, Powdery Mildew, Leaf Spot, Silver Mites, Flat Mites, Red Spider spp. 2-6 pounds per 100 gals. of water. Apply at bloom or early petal fall, repeat as necessary, usually 10-14 days, or after a period of wet weather.

PEACHES, NECTARINES: Brown Rot, Powdery Mildew, Silver Mites, Red Spider spp. 2-6 pounds per 100 gals, water. Apply at bloom or early petal fall, repeat as necessary, usually 10-14 days, or after a period of wet weather.

CITIOUS: Flat Mite, Silver Mites, Red Spider spp. 2-6 pounds per 100 gals. of water. Apply in any sprey dering season — NOT CONTAINING OIL or following oil by 14 days 1 [] [

DENESS Bud Mites, apply 2 lbs. per 100 gallons of water.

AVOCADO: Brown Mite, apply 2-6 lbs. per 100 gals. of water. Apply as needed.

GRAPES: Powdery Mildew, Bud Mite, Blister Mite, Red Spider spp. 2 to 6 pounds per 100 gals. of water. Apply in late dormant, or with cutworm spray, early shoot period and thru season on 14 day schedule, or with posticides.

BLACKBERRY: (Rubis Spp): Red Berry Mites (Eriophytes Gracilis) apply 3-5 lbs, per 100 gals, water. Start in early bud swell, repeat applications as necessary, with or without pesticides.

RASPBERRY (Rubis Spp): Powdery Mildew (Sphaerotheco Humuli) spray each week, starting at first bloom-color with 3-5 lbs. per 100 gals. of water. Continue until all fruit is set, then repeat as weather requires.

ACTIVE INGREDIENTS:

Micronized Sulfur

INERT INGREDIENTS:

TOTAL

BY WT.

80%

10%

STRAWBERRIES: Powdery Mildew, Red Spider, apply 3-5 lbs. per 100 gallons of water. Start in early leaf stage; repeat applications with or without pesticides.

TOMATOES: Russet Mite apply 3-5 lbs. per acre. Thorough coverage is required, 15 to 50 gallons of water either by air or ground rig.

POTATOES:

BEANS, PEAS, COLE CROPS, PEPPERS) Powdery Mildew, Red Spider spp. 3-5 pounds per acre in 15-50 gals, of water by aircraft or ground rig. Start in early leaf stage. Repeat application every 10-14 days with or without pesticides, or immediately after a rain.

HOPS: Red Spider (spp) apply 2-6 lbs, per 100 gals, water. Start in early leaf stage, repeat applications as needed, usually 10-14 days,

SUGAR BEETS: Red Spider Spp. apply 3-5 lbs. per acre in 15-50 gal. water, by aircraft or ground rig. Apply as needed with or without pesticides.

COTTON, SEED ALFALFA: Red Spider spp. Apply 3-5 pounds per acre in 25-50 gallons of water by air craft or ground rig.

PEANUTS: Powdery Mildew, Red Spider Miles Apply 3 to 5 lbs. per acre in 15 to 50 galloes of water by aircraft or ground rig. Short in early leaf stage. Repeat applications as necessary, essetly 10 to 14 days.


WARNING KEEP OUT OF REACH OF CHILDREN

Do not use on sulfur sensitive plants. Sulfur may burn foliage when temperature is excessively high. Do not apply with or following an oil application for 14 days. Do not apply oil for 14 days following treatment.

NET WT. 56 LBS. — KEEP DRY

Conditions of Sale

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with otherschemicals not specifically recommended, and other influencing factors in the section of the seller. Buyer assumes all risks are product are beyond the control of the seller. Buyer assumes all risks are product are beyond the control of the seller, accordance with direct therewith.



ACCEPTED

1/23/69

PUNISH ONE PEDEMAN SERVICE
FOR MEDICAL AND MEDICAL DESIGNATION
FOR MEDICAL DESIGNAT



TM COSAN --- Trademark