


RIEDEL'S "SUPERCOSAN"™

"FOR HORTICULTURE AND HOME GARDENING"

Fungicide - Miticide

USDA REGISTRATION NO. 9288-4



containing

ACTIVE INGREDIENTS:	BY WT.
Micronized Sulfur80%
INERT INGREDIENTS20%
TOTAL	100%

"FOR POWDERY MILDEW AND RED SPIDER MITE CONTROL
IN FLOWERING PLANTS, SHRUBS, TREES,
VEGETABLES AND VINES."

CAUTION! KEEP OUT OF REACH OF CHILDREN KEEP DRY
May cause irritation to eyes

IMPORTED BY

Haley & Kruger Co., Inc.

19931 EAST MANNING AVENUE

REEDLEY, CALIFORNIA 93654

Manufactured by:
RIEDEL-DE HAËN, AG
Seelze-Hannover,
West Germany.



TM COSAN — Trademark of Riedel-De Haen AG Int. Reg. B.I.R.P.I. No. 152354

RIEDEL'S "SUPERCOSAN"™ For Horticulture and Home Gardening Fungicide - Miticide

NET WT. _____

Mix the required amount of RIEDEL'S SUPERCOSAN with a small quantity of water to make a thin slurry, then dilute to desired concentration while agitating. Spray carefully to cover all parts of the plant tissue.

RIEDEL'S SUPERCOSAN is compatible with other compounds; organic fungicides and insecticides generally used in garden pest control. Do not combine with spray-oils.

DIRECTIONS FOR USE:

DISEASE AND PEST

CONCENTRATION

Powdery Mildew infesting Flowering Plants, Shrubs, Trees, Vines and Vegetables Crops.

Red Spider and "certain other mite species", on the following plants, trees, shrubs, vegetables and vines, Apples, Almonds, Cherries, Plums, Prunes, Peaches, Nectarines, Citrus, Grapes, Beans, Peas, Broccoli, Brussels sprouts, Cabbage, Cauliflower, Peppers, Cotton, Seed Alfalfa, Strawberries, and Tomatoes.

Early Applications: Use 6-7 oz per 10 gal.
(before leafing out) Add 1/2 to 1 oz RIEDEL'S 1814 WETTING AGENT

Maintenance Sprays: Use 2-3 oz per 10 gal.
Add 1/2 - 1 oz RIEDEL'S 1814 WETTING AGENT

Repeat applications at regular intervals.

Not usually more often than 7 - 10 day intervals or as needed.

The above recommendations are general in nature and will vary according to local climatic conditions, degree of infestation, sensitivity of the plant and method of application.

* Do not use on salt-tolerant plants. Salt may burn foliage when temperature is excessively high. Do not apply with or following an oil application in 14 days. Do not apply oil for 14 days following treatment.

* Some types of melons, gooseberries, apples, pears and apricots. When in doubt, it is recommended that a test be made on a few leaves before spraying to determine tolerance to SUPERCOSAN.

CONDITIONS OF SALE

Follow directions carefully. Timing and method of application, weather and temperature, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are at the control of the user. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given on label.

*There
is
no
substitute
for
Quality*

