


pop-in WEED CONTROL SPRAY

ACCEPTED
2-5-74
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND MITICIDE ACT
FOR CONTROL OF WEEDS REGISTERED
ED UNDER NO. 9307-3



FOR BEST RESULTS

Apply in early spring when plants are growing vigorously. Repeat as needed throughout growing season and when there is ample soil moisture.

Apply when air is calm or still—use coarse spray to avoid drift onto other desirable plants.

Apply only between temperatures of 50° and 85°F.

For overall coverage, carefully measure area and apply 1 gallon to 1,000 square feet. Apply correct dosage uniformly to each square foot.

To spot treat individual weeds—spray to runoff. Wet tall weeds thoroughly. Apply at double rate for hard to kill weeds, knotweed, chickweed, clover and mature weeds. Deep rooted perennials may require repeat applications.

Do not apply if heavy rain is expected within eight hours.

Avoid spraying 2-days before mowing or just after mowing when foliage is insufficient to catch spray. Seedling grass should be mowed twice before spraying. Do not re-seed for 3 weeks after spraying.

Use with caution when treating sensitive grasses—such as Bent Grass, Centipede Grass, St. Augustine Grass. When treating sensitive grasses, use half rate. Do not spray Dichondra!

A separate sprayer should be used for spraying weed killers and lawns.

CAUTION: Avoid inhaling dust or spray mist. Avoid contact with skin, eyes or clothing. Harmful if swallowed • Do not store near foodstuffs, seeds, fertilizers, insecticides or fungicides • Avoid use where there is a drift hazard.

EFFECTIVE FORMULATION

ACTIVE INGREDIENTS	LITHIUM 2,4-Dichlorophenoxyacetate* (equivalent to 52.3% of 2,4-D acid)	53.77%
INERT INGREDIENTS		46.23
		100.00%

*LITHATE • Guth Corporation

NOTICE

This product is guaranteed to conform to the ingredient statement stated on the label. Encap products are made to exacting standards to give you the best results possible. Since use of this material is beyond manufacturer's control, manufacturer is not responsible for uses and conditions beyond the recommendations of the label.

Sunflowers
Thistles
Velvetleaf
Wild Carrot
Wild Lettuce
Wild Onion
Shepherd's Purse


Knotweed
Lambsquarters
Morning Glory
Mustards
Pigweed
Primrose
Ragweed
Speedwell

Bindweed
Broadleaf Plantain
Buckhorn Plantain
Burdock
Chickweed
Chickory
Dandelion
Ground Ivy

CONTROLS ALL THESE

pop-in

CONTROLS BROADLEAF WEEDS IN
LAWNS • EACH PACKET TREATS
1,000 SQUARE FEET. A highly effective
formulation of non-volatile 2,4-D
for the control of dandelions and a
wide range of other broadleaf weeds.
Use regularly for a healthy, beautiful lawn.



CAUTION:
KEEP OUT OF REACH OF CHILDREN

POP-IN[®] WEED CONTROL SPRAY

ACTIVE INGREDIENTS
LITHIUM CHLORIDE 1.0%
INERT INGREDIENTS 99.0%

CAUTION: Avoid inhaling dust or spray mist. Avoid contact with skin, eyes or clothing. Harmful if swallowed. Do not store near foodstuffs, seeds, fertilizers, insecticides or fungicides. Avoid use where there is a drift hazard.

Please refer to outer carton for directions for acceptable uses, cautions and limitations.

E.P.A. REG. NO. 9307-3 AA
ENCAP PRODUCTS COMPANY
P.O. BOX 278
MT. PROSPECT, ILLINOIS 60056

CAUTION:
KEEP OUT OF REACH OF CHILDREN

POP-IN[®] WEED CONTROL SPRAY

ACTIVE INGREDIENTS
LITHIUM CHLORIDE 1.0%
INERT INGREDIENTS 99.0%

CAUTION: Avoid inhaling dust or spray mist. Avoid contact with skin, eyes or clothing. Harmful if swallowed. Do not store near foodstuffs, seeds, fertilizers, insecticides or fungicides. Avoid use where there is a drift hazard.

Please refer to outer carton for directions for acceptable uses, cautions and limitations.

E.P.A. REG. NO. 9307-3 AA
ENCAP PRODUCTS COMPANY
P.O. BOX 278
MT. PROSPECT, ILLINOIS 60056

CAUTION:
KEEP OUT OF REACH OF CHILDREN



POP-IN[®] WEED CONTROL SPRAY


ACTIVE INGREDIENTS
LITHIUM CHLORIDE 1.0%
INERT INGREDIENTS 99.0%

CAUTION: Avoid inhaling dust or spray mist. Avoid contact with skin, eyes or clothing. Harmful if swallowed. Do not store near foodstuffs, seeds, fertilizers, insecticides or fungicides. Avoid use where there is a drift hazard.

Please refer to outer carton for directions for acceptable uses, cautions and limitations.

E.P.A. REG. NO. 9307-3 AA
ENCAP PRODUCTS COMPANY
P.O. BOX 278
MT. PROSPECT, ILLINOIS 60056





SPECIALLY FORMULATED FOR THE POST-EMERGENCE CONTROL OF BROADLEAF AND SURFACE CREEPING WEEDS IN TURF. MAY BE USED ON KENTUCKY BLUE GRASS, MERION, BERMUDA, FESCUES AND RYE GRASSES.

WEEDS CONTROLLED:

Clover	Heal-all	Red Clover
Chickory	Knotweed	Red Sorrel*
Chickweed, common	Lambsquarters	Stitchwort
Chickweed, mouse-ear	Mallow	Speedwell
Cinquefoil	Morning Glory	Thistle
Clover	Pigweed	White Clover
Curled Deck	Plantain, buckhorn	Wild Carrot
Dandelion*	Plantain, common	Wild Garlic*
English Daisy*	Purslane	Wild Onion*
Ground Ivy	Ragsweed	Yarrow

DIRECTIONS FOR USE: Mix with water as follows:

	Per 1000 sq. ft.		Per acre	
	Product	Water	Product	Water
Low Rate	8 fl. oz.	2-4 gals.	2 1/2 gals.	80-160 gals.
High Rate	10 fl. oz.	2-4 gals.	3 gals.	80-160 gals.

Use high rate for weeds marked with an asterisk() above.

Apply mixture by means of a sprayer or sprinkling can to give uniform coverage. Repeat applications at the lower rate may be necessary, especially with heavy infestations of the less susceptible weeds indicated above with an asterisk(*).

See additional directions on the right side of this label.

Net contents gals.

ROUT

SELECTIVE WEED KILLER



Active Ingredients:
 Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid* 7.02%
 Dimethylamine salt of 2,4-dichlorophenoxyacetic acid** 6.99%
 Inert Ingredients: 85.99%
 100.00%

*Equivalent to 5.81% of 2-(2-methyl-4-chlorophenoxy) propionic acid or 0.5 lbs. MCPP acid per gallon.
 **Equivalent to 5.81% of 2,4-dichlorophenoxyacetic acid or 0.5 lbs. 2,4-D acid per gallon.
 EPA Reg. No. 9313-13

CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing of spray mist. Do not store with foodstuffs. Do not transport with or store near seeds, fertilizers, insecticides or fungicides. Do not use in or around greenhouses. Avoid contamination of water intended for irrigation or domestic purposes. Do not reuse empty containers, destroy by crushing and burying; or wash with an alkaline detergent and water and return to a drum reconditioner. Avoid spray drift to susceptible plants as injury may occur to grapes, tomatoes, beans, peas, other vegetables, legumes, ornamentals, etc. (Coarse sprays are less likely to drift) Because of danger of contamination, do not use same application equipment for applying insecticides, liquid fertilizers, etc. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes or containers.

See cautions elsewhere on container.

Mfd. by: Rose Chemical Products, Inc. Columbus, Ohio 43223

Either before or after application, it is wise to properly fertilize the lawn to encourage the grass to fill in the bare spots where the weeds have died away. Once the weeds have been removed, the bare areas should be reseeded. Bare areas may be reseeded 3 to 4 weeks after application. Apply in spring or fall when weeds are actively growing. Knotweed is more easily controlled in spring or early summer when weeds are young.

NOTE: MCP P acts more slowly than 2,4-D, so that some weeds will be controlled before others. Allow 3 to 4 weeks for complete weed killing action to occur.

CAUTION: Do not apply when turf is suffering from drought conditions or during very hot weather as damage to the grass may result. Do not mow for several days before or after application. Do not sprinkle for 24 hours after application. Do not apply during periods of heavy rainfall. Do not use on newly seeded areas. Do not use if clover is desired in the turf. In some areas, bent grasses, carpet, buffalo and St. Augustine grasses may be injured. Do not use on turf where these grasses predominate. Do not apply to Dichondra.

Local conditions may affect the use of this product. Consult Agricultural Experiment Station or Extension Service weed specialists for specific recommendations as to dosage, time of application and for control of other weeds.

NOTE: Store at temperatures above 32°F. If allowed to freeze, thaw and mix thoroughly before use.

Lab control no.