

JUL 19 1967

UNDER THE FEDERAL BUDGETING  
FUNCTIONS AND ORGANIZATION  
ACT OF 1950 (22 USC 2301-2304)  
E.O. 11649 9/9-1

**DIRECTIONS**

Use CHARDON X-TOX 404 in a good sprayer adjusted to deliver a fine mist. Where applicable, CHARDON X-TOX 404 may be used in a thermal type fogger. Regulate dosage in accordance with equipment manufacturers directions for the particular type of applicator employed. Application should be sufficient to fill the enclosure with the fog, mist or vapor. Treat infested areas thoroughly and whenever possible, keep doors and windows closed for 10 minutes after spraying.

**Flying Insects:** For control of flies, mosquitoes, gnats, wasps, and hornets, direct the spray or fog toward the upper portions of the room and saturate the atmosphere with the mist or fog, using 1.5 ounces to each 1000 cubic feet of space.

**Crawling Insects:** To control roaches, silverfish, crickets, and spiders, spray infested areas thoroughly and direct the liquid behind baseboards, moldings, cabinets, shelves, sinks, door and window frames, and pipes, toward drains and dark corners, and into cracks, crevices, recesses and similar hiding places. Repeated treatments are necessary for control of these crawling insects.

Against ant infestations, treat shelves, cabinets and storage areas; spray ant runs, trails, baseboards, thresholds and other places of entry, and treat nests liberally.

For controlling fleas and Brown Dog ticks in buildings, direct the application toward the lower wall portions, baseboards, rugs, carpets and floors of infested areas. Animal quarters, kennels and stabling should also be sprayed.

To control bedbugs effectively, apply liberally to top, ends, sides and bottom of mattresses, spray thoroughly all tufts, seams and folds. Spray springs and frame, directing the liquid into all cracks and lock joints. Treat walls, baseboards, moldings, floors, closets and bedroom furniture, picture frames, with particular attention to protected bedbug hiding and breeding places.

**CH**

CHARDON  
ants and  
buildings.  
should not  
useful for  
restaurants

CHARDON  
Roaches  
Silverfish  
Crickets  
Spiders  
Ants  
Flour Beetle  
Weevils

Active In  
Py  
N  
P  
A

\*Equivalent  
of related

Harmful if  
Remove P  
toward an  
only when

For most effective control of clothes moths, carpet beetles, remove infested articles from storage, brush thoroughly and, if possible, air for 24 hours in direct sunlight. Apply spray liberally to empty chests, closets, bureaus and storage, directing the liquid into cracks, joints and crevices. After airing, infested articles may be treated by holding the spray nozzle at least 3 feet away from the fabric and applying the liquid lightly to the seams to avoid the possibility of staining. Treat both sides of rugs, carpets and pads, and all exposed and accessible surfaces of overstuffed furniture. Do not overspray if products are made of rubber or are rubber coated, to prevent expansion or swelling. Repeat treatment monthly for articles not stored in moth-tight containers.

Restaurant and institutional pantries and storages frequently become infested with various stored food pests including flour beetles, weevils, cadelles, grain beetles, meal worms, sawtooth beetles, meal moths, skippers, grain and storage mites. To avoid sources of reinfestation, flour, grain and cereals should be destroyed before treatment. Spray thoroughly all storage shelves, bins, cupboards, cabinets, pantries and similar storages, with particular attention to cracks, niches, crevices and other hidden places. Cover all foodstuffs prior to treatment. Do not spray directly on foodstuffs. All processing surfaces should be covered and thoroughly cleaned before using.

Lab Control No



# ROSE 606

READY TO USE

## INSECTICIDE

CONTAINS "DUAL SYNERGIST"

### DIRECTIONS

Use ROSE 606 in a good sprayer adjusted to deliver a fine mist. Where applicable, ROSE 606 may be used in a thermal type fogger. Regulate dosage in accordance with equipment manufacturers directions for the particular type of applicator employed. Application should be sufficient to fill the enclosure with the fog, mist or vapor. Treat infested areas thoroughly and whenever possible, keep doors and windows closed for 10 minutes after spraying.

*VII indoors*  
*XI outdoors*  
Flying Insects: For control of flies, mosquitoes, gnats, wasps, and hornets, direct the spray or fog toward the upper portions of the room and saturate the atmosphere with the mist or fog, using 0.5 ounces to each 1000 cubic feet of space.

*XIV indoors*  
*XII outdoors*  
Crawling Insects: To control roaches, silverfish, crickets, and spiders, spray infested areas thoroughly and direct the liquid behind baseboards, moldings, cabinets, shelves, sinks, door and window frames, and pipes, toward drains and dark corners, and into cracks, crevices, recesses and similar hiding places.

*XIII indoors*  
*XV outdoors*  
Against ant infestations, treat shelves, cabinets and storage areas: spray ant runs, trails, baseboards, thresholds and other places of entry, and treat nests liberally.

*XVI indoors*  
In controlling fleas and Brown Dog ticks in buildings, direct the application toward the lower wall portions, baseboards, rugs, carpets and floors of infested areas. Animal quarters, kennels and bedding should also be sprayed.

*XVII indoors*  
To control bedbugs effectively, apply liberally to top, ends, sides and bottom of mattresses, spraying thoroughly all tufts, seams and folds. Spray springs and frame, directing the liquid into all cracks and lock joints. Treat walls, baseboards, moldings, floors, closets and bedroom furniture, picture frames, with particular attention to protected bedbug hiding and breeding places.

ROSE 606 is an effective, fast-acting insecticide for the control of roaches, ants and the other listed injurious and annoying insects found in homes, institutions and buildings. It can be used around food, food products and utensils as directed, but should not be applied directly to food or foodstuffs. ROSE 606 is highly useful for insect control in institutions, public buildings, hospitals, hotels, clubs, restaurants, stores, markets, theaters and other situations where an effective insecticide with a high degree of safety is required.

### ROSE 606 Insecticide Kills by contact:

- |            |            |               |                                 |
|------------|------------|---------------|---------------------------------|
| Roaches    | Ants       | Wasps         | Bedbugs                         |
| Silverfish | Flies      | Hornets       | Carpet Beetles                  |
| Crickets   | Mosquitoes | Sand Flies    | Fleas                           |
| Spiders    | Gnats      | Clothes Moths | Brown Dog Ticks<br>in buildings |

### ROSE 606 Insecticide kills by contact the exposed or accessible stages of:

- |               |               |                |              |
|---------------|---------------|----------------|--------------|
| Flour Beetles | Shippers      | Meal Worms     | Grain Mites  |
| Weevils       | Grain Moths   | Spider Beetles | Cheese Mites |
| Cadellies     | Grain Beetles | Meal Moths     |              |

### Active Ingredients

Fyrethrins	0.20%
N-Octyl bicycloheptene dicarboximide	0.67%
Piperonyl Butoxide, Technical*	0.40%
Aliphatic Petroleum Solvent	98.73%
	<u>100.00%</u>

\*Equivalent to 0.32% of (butylcarbityl) (5-propylpiperonyl) ether and to 0.08% of related compounds.

USDA Reg. No. 9313-2

## CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, avoid contact with skin. Do not breathe mist, fog, or vapor. Remove Pets, Bird and Fish Bowls from room before spraying. Do not spray near or toward an open flame. Use the product as a space spray in food processing plants only when plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered, thoroughly cleaned before using. Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place. Empty drums should be returned to a drum reconditioner.

*Moth moths*  
For most effective control of clothes moths and carpet beetles, remove infested articles from storage, brush thoroughly and, if possible, air several hours in direct sunlight. Apply spray liberally in empty chests, closets, bureaus and storages, directing the liquid into cracks, joints and crevices. After airing, infested articles may be treated by holding the spray nozzle at least 3 feet away from the fabric and applying the liquid lightly to the seams to avoid the possibility of staining. Treat both sides of rugs, carpets and pads, and all exposed and accessible surfaces of overstuffed furniture. Do not overspray if products are made of rubber or are rubber coated, to prevent expansion or swelling. Repeat treatment as needed for susceptible articles not stored in moth-tight containers.

*XVIII outdoors*  
Restaurant and institutional pantries and food storages frequently become infested with various stored food pests including flour beetles, weevils, cadellies, grain beetles, meal worms, spider beetles, meal moths, skippers, grain and cheese mites. To avoid sources of reinfestation, wormy flour, grain and cereals should be destroyed before treatment. Spray thoroughly all surfaces of shelves, bins, cupboards, cabinets, pantries, lockers and similar storages, with particular attention to cracks, niches, crevices and other hiding places. Cover all foodstuffs prior to treatment. Do not spray directly on foodstuffs. All food processing surfaces should be covered or thoroughly cleaned before using.

Lab Control No. \_\_\_\_\_

Net Contents \_\_\_\_\_ Gals.

**DIRECTIONS**

Use CHARDON X-TOX 606 in a good sprayer adjusted to deliver a fine mist. Where applicable, CHARDON X-TOX 606 may be used in a thermal type fogger. Required dosage in accordance with equipment manufacturers directions for the particular type of applicator employed. Application should be sufficient to fill the enclosure with the fog, mist or vapor. Treat infested areas thoroughly and whenever possible, keep doors and windows closed for 10 minutes after spraying.

**Flying Insects:** For control of flies, mosquitoes, gnats, wasps and hornets, direct the spray or fog toward the upper portions of the room and saturate the atmosphere with the mist or fog, using 0.5 ounces to each 1000 cubic feet of space.

**Crawling Insects:** To control roaches, silverfish, ants and spiders, spray infested areas thoroughly and direct the liquid behind baseboards, along pipes, shelves, sinks, doors and window frames, and pipes, toward drains and dark corners and into cracks, crevices, recesses and other hiding places.

**Application Instructions:** Treat shelves, cabinets and storage cases, spray ant runs, trails, baseboards, closets and other places of entry, and treat nests/betales.

**Controlling Fleas and Brown Dog Ticks in Buildings:** Direct the application toward the lower wall portions, baseboards, baseboards and floors of infested areas. Animal quarters, kennels and barns should also be treated.

**For the best results:** Apply liberally to top, sides and bottom of drawers, spraying into all tufts, seams and folds. Spray into all frame joints, door and baseboards, along floor closets, and under furniture. Do not use on upholstered furniture to prevent damage to the fabric.



# CHARDON X-TOX 606

READY TO USE  
**INSECTICIDE**

**CONTAINS "DUAL SYNERGIST"**

CHARDON X-TOX 606 is an effective, fast-acting insecticide for the control of roaches, ants and other listed injurious and annoying insects found in homes, institutions and buildings. It can be used around food, food products and utensils as directed, but should not be applied directly to food or foodstuffs. CHARDON X-TOX 606 is highly useful for insect control in institutions, public buildings, hospitals, hotels, clubs, restaurants, stores, markets, theaters and other situations where an effective insecticide with a high degree of safety is required.

**CHARDON X-TOX 606 Insecticide Kills by contact:**

- |            |            |               |                                 |
|------------|------------|---------------|---------------------------------|
| Roaches    | Ants       | Wasps         | Bedbugs                         |
| Silverfish | Flies      | Hornets       | Carpet Beetles                  |
| Cricket    | Mosquitoes | Sand Flies    | Fleas                           |
| Spiders    | Gnats      | Clothes Moths | Brown Dog Ticks<br>in buildings |

**CHARDON X-TOX 606 Insecticide kills by contact the exposed or accessible stages of:**

- |               |               |                |              |
|---------------|---------------|----------------|--------------|
| Flour Beetles | Shippers      | Meal Worms     | Grain Mites  |
| Weevils       | Grain Moths   | Spider Beetles | Cheese Mites |
| Caddisflies   | Grain Beetles | Meal Moths     |              |

<b>Active Ingredients</b>	
Pyrethrins	0.20%
N-Octyl bicycloheptene dicarboximide	0.07%
Piperonyl Butoxide, Technical	0.40%
Aliphatic Petroleum Solvent	98.73%
	<hr/> 100.00%

\*Equivalent to 0.32% of (butylcarbityl-β-propylpiperonyl) ether and to 0.08% of related compounds.

USDA Reg. No. 9313-2

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed, avoid contact with skin. Do not breathe mist, fog, or vapor. Remove Pets, Bird and Fish Bowls from room before spraying. Do not spray near or toward an open flame. Use the product as a space spray in food processing plants only when plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using. Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place. Empty drums should be returned to a drum reconditioner.

Manufactured for:

**Chardon Laboratories, Inc.**

P.O. Box 22401 Columbus, Ohio 43223

For most effective results, use CHARDON X-TOX 606 in a good sprayer adjusted to deliver a fine mist. Where applicable, CHARDON X-TOX 606 may be used in a thermal type fogger. Required dosage in accordance with equipment manufacturers directions for the particular type of applicator employed. Application should be sufficient to fill the enclosure with the fog, mist or vapor. Treat infested areas thoroughly and whenever possible, keep doors and windows closed for 10 minutes after spraying.

Resistant and other insects found in stored food, such as flour, grain, and other foodstuffs, should be treated by holding the spray nozzle 3 feet away from the food and spraying the liquid lightly to the seams, joints, and parts of staining. Treat both sides of the pads, and all exposed surfaces of the overstuffed furniture. Do not use on objects made of rubber or plastic to prevent expansion or swelling. Use as needed for storage of food in moth-tight containers.

Net Content

Net Content