
# ALGEX 10

## A CONCENTRATED SWIMMING POOL ALGAECIDE. WHEN USED AS DIRECTED, IT WILL KILL AND PREVENT DEVEL-OPMENT OF CHLORELLA PRYENOIDOSA (GREEN ALGAE), PHORMIDIUM INUNDATUM (BLACK ALGAE). PHORMIDIUM RETZII (SQUARE D ALGAE), LYNGBA VERSICOLOR (BLUE-GREEN ALGAE), AND OTHER SWIMMING POOL ALGAE, WILL NOT HARM POOL EQUIPMENT AT USE-DILUTION. COMPATIBLE WITH MOST COMMONLY USED SWIMMING POOL CHEMICALS, BUT SHOULD NOT BE MIXED WITH SOAPS OR CLEANERS OR PREMIXED WITH OTHER WATER-TREATING CHEMICALS. FOR BEST RESULTS, ADD DIRECTLY TO POOL WATER.

Net contents \_\_\_\_\_ ga/lon(s)

Lab control no.



## CONCENTRATED SWIMMING POOL ALGAECIDE



#### ACTIVE INGREDIENTS

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl Julizyl ammonium n-Alkyl (68% C12, 32% C14) dimethyl ethylbenzul ammonium chlorides . . 5.0%

EPA Reg. No. 9313-10

## DANGER KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed Avoid contamination of food.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Use product according to directions. Do not reuse empty container. Rinse thoroughly with

water and destroy.

FIRST AID In case of contact, immediately flush eyes or skin with plenty of water for 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse if swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol, Call a physician immediately NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock as well as oxygen and measures to support breath ing manually or mechanically may be needed. If persistent, convulsions may be controlled by the cautious intravenous injection of a short - acting barbituate drug.

Mfg. by: Rose Chemical Products, Inc. 545 Stimmel Road

Columbus, Ohio 43225

100.0%

### Initial Dose:

1 gallon to 50,000 gallons of water. (Provides 2 ppm active quaternary).

Maintenance Dose: 1 quart to 50,000 gallons of water. (Provides ½ ppm active quaternary).

DIRECTIONS

Initial dose is used when filling pool at beginning of swimming pool season. If pool has visible algae growth, treat with initial dose. Vacuum pool after 24 hours to remove algae debris. If visible algae are still present or reappear, repeat this treatment as necessary. When pool is free of visible algae, use recommended maintenance dose.

Maintenance dose should in added at least once every 3-5 days. If high temperatures prevail or pool has unusually heavy use, add maintenance dose more frequently.

If makeup water is added at the rate of more than 10% of pool capacity per week, and 1 quart for each 10,000 gallons of fresh makeup water added per week.


When the swimming season is over, add 1 gallon to 50,000 gallons of water left in pool. This dose helps provide a measure of control of algae growth during the winter months. This treatment will also help the servicing of the pool before it is returned to usage the following season.



A CONCENTRATED SWIMMING POOL ALGAECIDE. WHEN USED AS DIRECTED, IT WILL KILL AND PREVENT DEVEL-OPMENT OF CHLORELLA PRYENOIDOSA (GREEN ALGAE), PHORMIDIUM INUNDATUM (BLACK ALGAE), PHORMIDIUM RETZII (SQUARE D ALGAE), LYNGBA VERSICOLOR (BLUE-GREEN ALGAE), AND OTHER SWIMMING POOL ALGAE, WILL NOT HARM POOL EQUIPMENT AT USE-DILUTION. COMPATIBLE WITH MOST COMMONLY USED SWIMMING POOL CHEMICALS, BUT SHOULD NOT BE MIXED WITH SOAPS OR CLEANERS OR PREMIXED WITH OTHER WATER-TREATING CHEMICALS. FOR BEST RESULTS, ADD DIRECTLY TO POOL WATER.

Net contents \_\_\_\_\_ gailon(s)

Lab control no.





ACTIVE INGREDIENTS

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium n-Alkyl (68% C12, 32% C14) dimethyl ethylbenzul ammonium chlorides . . 5.0%

EPA Reg. No. 9313-10

DANGER KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food. This product is toxic to fish. Keep out of lakas, streams or ponds. Do not contaminate water

by cleaning of equipment, or disposal of wastes. Use product according to directions. Do not reuse empty container. Rinse thoroughly with

water and destroy.

FIRST AID: In case of contect, immediately flush eyes or skin with plenty of water for 15 minutes. For eyes, call a physician.Framove and wash contaminated clothing before reuse. f swallowed, drink promptly a large quantity of milk, agg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol, Call a physician immediately. NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock as well as oxygen and measures to support breathing manually or mechanically may be needed. If persistent, convulsions may be controlled by the cautious intravenous injection of a short - acting barbituate drug.

Mig. by: Rose Chemical Products, Inc.

545 Stimmel Road

Columbus, Ohio 43223

## CONCENTRATED SWIMMING POOL ALGAECIDE

100.0%

- -- ·

### Initial Dose:

1 gallon to 50,000 gallons of water. (Provides 2 ppm active quaternary).

1 quart to 50,000 gallons of water. Maintenance Dose:

DIRECTIONS

(Provides ½ ppm active quaternary). Initial dose is used when filling pool at beginning of swimming pool season. If pool has visible algae growth, treat with initial dose. Vacuum pool after 24 hours to remove algae debris. If visible algae are still present or reappear, repeat this treat-

ment as necessary. When pool is free of visible algae, use recommended maintenance dose. Maintenance dose should be added at least once every 3-5

days. If high temperatures prevail or pool has unusually heavy use, add maintenance dose more frequently.

If makeup water is added at the rate of more than 10% of pool capacity per week, add 1 quart for each 10,000 gallons of fresh makeup water added per week.

When the swimming season is over, add 1 gallon to 50,000 gallons of water left in pool. This dose helps provide a measure of control of algae growth during the winter months. This treatment will also help the servicing of the pool before it is returned to usage the following seaso .

> *ب* ۱ and the second second