

SPECIALLY FORMULATED FOR THE POST-EMERGENCE CONTROL OF BROADLEAF AND SURFACE CREEPING WEEDS IN TURF. MAY BE USED ON KENTUCKY BLUE GRASS, MERION, BERMUDA, FESCUES AND RYE GRASSES.

WEEDS CONTROLLED:

Clover	Heal-all	Red Clover
Chickory	Knotweed	Red Sorrel*
Chickweed, common	Lambsquarters	Stitchwort
Chickweed, mouse-ear	Mallow	Speedwell
Cinquefoil	Morning Glory	Thistle
Clover	Pigweed	White Clover
Curled Dock	Plantain, buckhorn	Wild Carrot
Dandelion*	Plantain, common	Wild Garlic*
English Daisy*	Purslane	Wild Onion*
Ground Ivy	Ragsweed	Yarrow

DIRECTIONS FOR USE: Mix with water as follows:

	Per 1000 sq. ft.		Per acre	
	Product	Water	Product	Water
Low Rate	8 fl. oz.	2-4 gals.	2 1/2 gals.	80-160 gals.
High Rate	10 fl. oz.	2-4 gals.	3 gals.	80-160 gals.

Use high rate for weeds marked with an asterisk () above.


Apply mixture by means of a sprayer or sprinkling can to give uniform coverage. Repeat applications at the lower rate may be necessary, especially with heavy infestations of the less susceptible weeds indicated above with an asterisk (*).

See additional directions on the right side of this label.

Net contents gals

ROUT

SELECTIVE WEED KILLER



[Faint, mostly illegible text, likely bleed-through from the reverse side of the page.]

CAUTION KEEP OUT OF REACH OF CHILDREN

[Faint, mostly illegible text, likely bleed-through from the reverse side of the page.]

Mfd. by Rose Chemical Products, Inc. Columbus, Ohio 43213

Either before or after application, it is wise to properly fertilize the lawn to encourage the grass to fill in the bare spots where the weeds have died away. Once the weeds have been removed, the bare areas should be reseeded. Bare areas may be reseeded 3 to 4 weeks after application. Apply in spring or fall when weeds are actively growing. Knotweed is more easily controlled in spring or early summer when weeds are young.

NOTE: MCPP acts more slowly than 2,4-D, so that some weeds will be controlled before others. Allow 3 to 4 weeks for complete weed killing action to occur.

CAUTION: Do not apply when turf is suffering from drought conditions or during very hot weather as damage to the grass may result. Do not mow for several days before or after application. Do not sprinkle for 24 hours after application. Do not apply during periods of heavy rainfall. Do not use on newly seeded areas. Do not use if clover is desired in the turf. In some areas, bent grasses, carpet, buffalo and St. Augustine grasses may be injured. Do not use on turf where these grasses predominate. Do not apply to Dichondra.

Local conditions may affect the use of this product. Consult Agricultural Experiment Station or Extension Service weed specialists for specific recommendations as to dosage, time of application and for control of other weeds.

Do not use at temperatures above 32 F. If allowed to freeze, thaw and mix thoroughly before use.

Lab control no