

A WATER SOLUBLE LIQUID TO BE MIXED WITH WATER AND APPLIED AS A SPRAY ON NON-CROPLAND AREAS. AN EFFECTIVE GENERAL HERBICIDE FOR CONTROL OF BRUSH AND MANY ANNUAL AND PERENNIAL WEEDS, AND IS PARTICULARLY USEFUL FOR CONTROL OF PERENNIAL GRASSES. EFFECTS ARE SLOW TO APPEAR AND MAY NOT BECOME APPARENT UNTIL THE PRODUCT HAS BEEN CARRIED INTO THE ROOT ZONE OF THE WEEDS BY MOISTURE. THE DEGREE OF CONTROL AND DURATION OF EFFECT WILL VARY WITH THE AMOUNT OF PRODUCT APPLIED, SOIL TYPE, RAINFALL AND OTHER CONDITIONS. WHEN MIXED WITH WATER, IT IS NON-VOLATILE, NON-FLAMMABLE AND NON-CORROSIVE TO METALS EXCEPT ALUMINUM. DO NOT USE WITH ALUMINUM SPRAY NOZZLES.

DIRECTIONS

Apply just before or during the period of active growth of plants to be controlled. If dense growth is present, results will be improved if vegetation is removed before treatment. Do not apply when ground is frozen.

Before spraying, calibrate equipment to determine quantity of water necessary to uniformly cover measured area to be treated. Measure the proper amount of product and mix into the necessary volume of water. Mix thoroughly. Once mixed, further agitation is not required.

Use a fixed-boom power sprayer properly calibrated to a constant speed and rate of delivery. For application with a hand-gun sprayer, use at least 200 gals. of diluted spray per acre (approx. 5 gals. of diluted spray per 1000 sq. ft.) For treating small areas, a tank-type hand sprayer or sprinkling can may be used.

WEED CONTROL: To control most weeds for an extended period of time on non-cropland areas such as railroads, highway and pipeline right of ways, petroleum tank farms, lumberyards, storage areas, industrial plant sites and drainage ditch banks.

Apply 3 to 6 qts. per 1000 sq. ft. to control ANNUAL WEEDS and GRASSES such as foxtail, ryegrass, wild oats, crabgrass, cheatgrass, bromegrass, ragweed, lambsquarters, puncturevine, orchardgrass and turkey mullein. When applied just prior to or after emergence of annuals, rates as low as 1½ qts. per 1000 sq. ft. control many annual weeds and grasses in low rainfall areas and give short term control in higher rainfall areas.

Apply 6 to 12 qts. per 1000 sq. ft. to control PERENNIAL WEEDS and GRASSES such as smooth brome, Bahiagrass, bluegrass, redtop, purpletop, quackgrass, broomsedge, aster, dandelion, dog fennel, goldenrod, plantain and wild carrot. In areas of low or seasonal rainfall, rates as low as 4½ qts. per 1000 sq. ft. control many perennial weeds and grasses.

(Continued on right panel)

ACCEPTED

FEB 23 1968

UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE AND RODENTICIDE
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 9313-18 SUBJECT
TO ATTACHED COMMENTS.



**A WATER-SOLUBLE, NON-SELECTIVE,
PERSISTENT VEGETATION KILLER**

Active Ingredient:	Lithium salt of bromacil (5-bromo-3-sec-butyl-6-methyluracil)*	1.12%
Inert Ingredients	98.88%
		100.00%

Contains 0.092 lbs. bromacil per gal.
*U.S. Pat. 3,352,862
EPA Reg. No. 9313-18

WARNING KEEP OUT OF REACH OF CHILDREN

HARMFUL OR FATAL IF SWALLOWED. Causes eye irritation. May irritate nose, throat and skin. Do not get in eyes. Avoid contact with skin and clothing. Avoid breathing spray mist. Wash thoroughly after handling. In case of contact, flush eyes or skin with plenty of water; for eyes, get medical attention.

Do not reuse empty container; crush and bury when empty, or return to a drum reconditioner. Keep container closed when not in use.

See cautions elsewhere on this label.

Mfg. by: ROSE CHEMICAL PRODUCTS, INC.
545 Stimmel Road, Columbus, Ohio 43223

Apply 12 to 15 qts. per acre to control **JOHNSONGRASS**, **HARD-TO-KILL** and other weeds such as **Barnyardgrass**, **Vaseygrass**, **crabgrass**, **bracken fern**, **fall (usually active growth)**. In the West, this is the best control of weeds such as

Use the high adsorptive matter or ca

Retreatment when annual sites where

BRUSH CONTROL plants on right of way and drainage

Apply in spots, basal (spot) ditches, app Note: For et of damage to brush ste from treated irrigation de sites where trees and sh treated area

Apply properly calibrated rate of delivery. Use as a backpack sprayer, use at 1000 sq. ft. For type hand sprayer

Use on most weeds for an non-cropland areas and pipeline right of way, orchards, storage and drainage ditch

Apply 12 to 24 qts. per 1000 sq. ft. to control ANNUALS such as foxtail, yellow top, prostrate knotweed, puncturevine, etc. When applied to annuals, rates of 12 to 24 qts. per 1000 sq. ft. control many low rainfall areas in higher rainfall

Apply 12 to 24 qts. per 1000 sq. ft. to control PERENNIALS such as smooth crabgrass, redtop, purpletop, dandelion, dogwood, wild carrot. In high rainfall, rates as low as 12 qts. per 1000 sq. ft. control many perennial

panel)

ACCEPTED

FEB 23 1968

UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTRATION UNDER NO. 9313-18 SUBJECT TO ATTACHED COMMENTS.



A WATER-SOLUBLE, NON-SELECTIVE, PERSISTENT VEGETATION KILLER

Active Ingredient:	Lithium salt of bromacil (5-bromo-3-sec-butyl-6-methyluracil)*	1.12%
Inert Ingredients		98.88%
		100.00%

Contains 0.092 lbs. bromacil per gal.
*U.S. Pat. 3,352,862
EPA Reg. No. 9313-18

WARNING KEEP OUT OF REACH OF CHILDREN

HARMFUL OR FATAL IF SWALLOWED. Causes eye irritation. May irritate nose, throat and skin. Do not get in eyes. Avoid contact with skin and clothing. Avoid breathing spray mist. Wash thoroughly after handling. In case of contact, flush eyes or skin with plenty of water; for eyes, get medical attention.

Do not reuse empty container; crush and bury when empty, or return to a drum reconditioner. Keep container closed when not in use. See cautions elsewhere on this label.

Mfg. by: ROSE CHEMICAL PRODUCTS, INC.
545 Stimmet Road, Columbus, Ohio 43223

(Continued from left panel)

Apply 12 to 24 qts. per 1000 sq. ft. to control **JOHNSONGRASS**; use at the same rate for **OTHER HARD-TO-KILL PERENNIAL WEEDS and GRASSES** such as Bermudagrass, Dallisgrass, nutsedge, Vaseygrass, saltgrass, bouncingbet, dogbane, bracken fern and horsetail. Where limited rainfall (usually less than 4 inches) occurs during the active growth period, such as some areas of the West, this product will not provide satisfactory control of hard-to-kill, deep-rooted perennial weeds such as Johnsongrass.

Use the higher levels of the dosage ranges on adsorptive soils (usually those high in organic matter or carbon).

Retreatment - Apply 1 1/2 to 6 qts. per 1000 sq. ft. when annual weeds and grasses reappear on sites where weed growth has been controlled.

BRUSH CONTROL: To control undesirable woody plants on non-cropland areas such as railroad right of ways, storage areas, industrial plant sites and drainage ditches.

Apply in spring or summer as a broadcast or basal (spot) treatment; for use on drainage ditches, apply as a basal (spot) treatment only. Note: For effective brush control and prevention of damage to desirable vegetation: do not apply to brush standing in water; do not use water from treated ditches for irrigation; do not use in irrigation ditches nor on right of ways or other sites where marketable timber or other desirable trees and shrubs are immediately adjacent to the treated area.

Broadcast Treatment - Apply 4 1/2 to 10 qts. per 1000 sq. ft. to control oak, willow, sweet gum and pine; apply 12 to 24 qts. per 1000 sq. ft. to control brush such as American elm, winged elm, hackberry, sumac and cottonwood. Use the higher levels of the dosage ranges on adsorptive soils (those high in organic matter or carbon).

Basal (Spot) Treatment - Apply undiluted at the rate of 4 to 8 fl. oz. per stem 2" to 4" in basal diameter; wet base of stem to run-off. Treatment controls woody plants such as cottonwood, hackberry, maple, oak, poplar, red bud, sweet gum, wild cherry, willow and winged elm.

IMPORTANT NOTE - Injury to or loss of desirable trees or other plants may result from failure to observe the following: Do not apply, or drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, driveways, tennis courts or similar areas. Prevent drift of spray to desirable plants. Do not contaminate domestic waters. Keep from contact with fertilizers, insecticides, fungicides and seeds. Thoroughly clean all traces of product from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

Lab control no. _____

Net contents _____ gal(s).