

AT-90

AMINO TRIAZOLE WEEDKILLER

Active ingredient:

Amitrole* 90.0%
Inert ingredients .. 10.0%
*3-amino-1,2,4-triazole

Cameraman's Note
Poor Copy

DIRECTIONS

THOROUGHLY SPRAY the amount water for 100 gallons per acre to Add 1 to 2 pts. of UNIFILM 707 thoroughly wet soil foliage. In highway spraying, spray 8 ft wide and 1 mph being a constant to 1 mph. Shake the required amount of AT 90 in water then add the Unifilm.

WOODY PLANTS AND VINES IN NONCROPLANDS

Weeds on Roadside, Drainage Ditchbanks, Fencerows
Amount Per Acre When to Apply

<p><u>Railroad-vine</u> <u>Poison Oak</u></p>	<p>2 lbs. in 100 gals. water (3 level tablespoons) in 1 gallon water</p>	<p>Apply when this growth is young and growing actively.</p>
<p><u>Honeysuckle</u> <u>Kudzu Vine</u></p>	<p>5 1/2 lbs. in 125 gals.</p>	<p>Treat from the time foliage is fully developed until plants begin to go dormant. For effective kill, all leaves, stems and suckers must be thoroughly wet to ground line.</p>
<p><u>Western dewberry</u> <u>(blackberry)</u> <u>Big leaf maple</u> <u>Wild cherry</u> <u>Ash, Locust, Sumac</u></p>	<p>2 lbs. in 100 gals. water</p>	<p>Spot treat any regrowth.</p> <p><i>Feb 12, 1974</i></p> <p><i>9319-4</i></p>
<p><u>Cattails and tules</u> <u>PHRASMITES</u></p>	<p>8 to 10 lb. in 300 gal. water</p>	<p>Apply after catkins are fully formed but before frost. Do not disturb sprayed plants. Spot treat any regrowth.</p>

WEED CONTROL IN NONCROPLANDS

Weeds on Roadside, Drainage Ditchbanks, Fencerows
Amount Per Acre When to Apply

<p>CONTROL annual grasses and broadleaf weeds (wild radish, yellow grass, timothy, alfalfa, lucerne, pigweed, horseweed, etc.)</p>	<p>2 lbs. in 30-60 gals water</p>	<p>Spray when most weeds are 3-4 inches tall.</p>
<p>CONTROL annual grasses, annual and perennial broadleaf weeds (whiteclover, Canada clover, etc.) SUPPRESS perennial grasses (legume grass, meadowgrass)</p>	<p>4 lb. in 100 gal. water</p>	<p>Spray when most weeds are 6-10 inches tall.</p>
<p>SUPPRESS PERENNIAL GRASSES Bermuda grass (quackgrass) Nutgrass (sedge)</p>	<p>4 lbs. in 100 gals. water</p>	<p>Spray when weeds are 3-6 inches tall.</p>

Weeds on Roadside, Fencerows with Simazine 80W

		Amounts per Acre	
		AT 90	Simazine
a - For seasonal control of most weeds where safety to nearby desirable ornamental trees, shrubs is important.	4 lbs.	12 1/2 lbs. (80W)	
b - For economical control of easy-to-kill (annual) weeds in Roadside, Drainage Ditchbanks, Fencerows	2 lbs.	6 1/2 lbs. (80W)	

WARNING!

DO NOT SPRAY OR ALLOW SPRAY DRIFT TO CONTAMINATE EDIBLE CROPS OR WATER WHICH WILL BE USED FOR IRRIGATION, DRINKING OR FISHING OR OTHER DOMESTIC PURPOSES.

AS NO RESIDUE OF AMITROLE IS PERMISSIBLE ON CROPS. DO NOT SPRAY OR PERMIT DRIFT OF SPRAY ON LAWNS, SHRUBS, FLOWERS AND OTHER DESIRABLE VEGETATION SINCE THIS PRODUCT WILL CAUSE INJURY. DO NOT USE IN GREENHOUSES Lilac, privet, honeysuckle, Boxwood, and barberry are susceptible to this mixture and may be injured if contacted by spray.

CUSTOM CHEMICIDES INC.

BERKELEY, CALIFORNIA 94710

NET WEIGHT:

..... pounds

WARNING!

KEEP OUT OF REACH OF CHILDREN

Coarse sprays are less likely to drift.

KEEP LIVESTOCK OFF TREATED AREAS
WARNING - KEEP OUT OF THE REACH OF CHILDREN
DOMESTIC ANIMALS AND FOODSTUFFS

MAY CAUSE IRRITATION OF NOSE AND THROAT MAY BE HARMFUL IF INHALED OR SWALLOWED DO NOT BREATHE SPRAY MIST USE WITH ADEQUATE VENTILATION AVOID CONTACT WITH EYES SKIN OR CLOTHING WASH THOROUGHLY AFTER HANDLING IN CASE OF CONTACT FLUSH WITH PLENTY OF WATER FOR EYES GET MEDICAL ATTENTION

DO NOT STORE NEAR FERTILIZERS, SEEDS, OR PESTICIDES OR USE IN A GREENHOUSE DO NOT SPRAY OR ALLOW SPRAY DRIFT TO CONTAMINATE NEARBY CROPS

DO NOT REUSE EMPTY CONTAINER DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

IMPORTANT DO NOT ALLOW SPRAY SOLUTION TO REMAIN IN THE SPRAYER ANY LONGER THAN NECESSARY RINSE ALL SPRAY EQUIPMENT WITH CLEAN WATER IMMEDIATELY AFTER EACH USE.

EPA Reg. No. 9319-11

Observe all cautions and limitations on labeling of all products used in mixtures.

Cameraman's Note
Poor Copy

AT-90

AMINO TRIAZOLE WEEDKILLER

Active Ingredient:

Amitrole[®] 90.0%

Inert Ingredients . . 10.0%

[®]3-amino-1,2,4-triazole

WEED CONTROL
Weeds on Roadsides, Drain
Amount Per Acre

DIRECTIONS

MIXING SPRAY: Use enough water (50 to 100 gallons per acre) to thoroughly wet weed foliage. In highway spraying, a strip 8 ft. wide and 1 mile long is equivalent to 1 acre. **Directions:** Measure the required amount of A.T. 90 in water then add the Simazine.

Add 1 to 2 pts. of UNIFILM 707 spreader sticker per 100 gallons to assure complete wetting.

WOODY PLANTS AND VINES IN NONCROPLANDS

Weeds on Roadsides, Drainage Ditchbanks, Fencerows.
Amount Per Acre When to Apply

<p><u>Rhizon Ivy</u> <u>Poison Oak</u></p>	<p>2 lbs. in 100 gals. water (3 level tablespoons) in 1 gallon water</p>	<p>Apply when this growth is young and growing actively.</p>	<p>CONTROL annual grasses and broadleaf weeds (cutting-grass, chestnut grass, barnyard grass, pigweed, foxtail, sunflower, wild barley)</p>	<p>2 lbs. in 30-60 gals water</p>
<p><u>Honeysuckle</u> <u>Kudzu Vine</u></p>	<p>5½ lbs. in 125 gals.</p>	<p>Treat from the time foliage is fully developed until plants begin to go dormant. For effective kill, all leaves, stems and suckers must be thoroughly wet to ground line.</p>	<p>CONTROL annual grasses, annual and perennial broadleaf weeds (whiteton, Canada thistle, milkweed) SUPPRESS perennial grasses (rigid grass, quackgrass)</p>	<p>4 lb. in 100 gal. water</p>
<p><u>Western dewberry (blackberry)</u> <u>Big leaf maple</u> <u>Wild cherry</u> <u>Ash, Locust, Sumac</u></p>	<p>2 lbs. in 100 gals. water</p>	<p>Spot treat any regrowth. <i>Feb. 12, 1974</i></p>	<p>SUPPRESS PERENNIAL GRASSES Bermuda grass (quackgrass) Nutgrass (sedge)</p> <p>Weeds on Roadsides, Fencerows with Simazine 80W</p>	<p>4 lbs. in 100 gals. water</p>

Honeysuckle Kudzu Vine	5½ lbs. in 125 gals.	<p>Treat from the time foliage is fully developed until plants begin to go dormant. For effective kill, all leaves, stems and suckers must be thoroughly wet to ground line.</p> <p>Spot treat any regrowth.</p> <p><i>Feb. 12, 1974</i></p> <p><i>9319-11</i></p>
Western dewberry (blackberry) Big leaf maple Wild cherry Ash, Locust, Sumac	2 lbs. in 100 gals. water	
cattails and tules <u>PARASMITES</u>	8 to 10 lb. in 300 gal. water	

CONTROL annual grasses, annual and perennial broadleaf weeds (white top, Canada thistle, milium) SUPPRESS perennial grasses (rigid grass, quackgrass)	4 lb. in 100 gal. water
SUPPRESS PERENNIAL GRASSES Bermuda grass quackgrass) Nutgrass (sedge)	4 lbs. in 100 gals. water
Weeds on Roadsides, Fencerows with Simazine 80W	
<p>a - For seasonal control of most weeds where safety to nearby desirable ornamental shrubs is important.</p> <p>b - For economical control of easy-to-kill (annual) weeds in Weeds on Roadsides, Drainage Ditchbanks, Fencerows</p>	

WARNING!
DO NOT SPRAY OR ALLOW SPRAY DRIFT TO CONTAMINATE EDIBLE CROPS OR WATER WHICH WILL BE USED FOR IRRIGATION, DRINKING OR FISHING OR OTHER DOMESTIC PURPOSES.

AS NO RESIDUE OF AMITROLE IS PERMISSIBLE ON CROPS. DO NOT SPRAY OR PERMIT DRIFT OF SPRAY ON LAWNS, SHRUBS, FLOWERS AND OTHER DESIRABLE VEGETATION SINCE THIS PRODUCT WILL CAUSE INJURY. DO NOT USE IN GREENHOUSES Lilac, privet, honeysuckle, Boxwood, and barberry are susceptible to this mixture and may be injured if contacted by spray.

WARNING!
KEEP OUT OF REACH OF CHILDREN
Coarse sprays are
KEEP LIVESTOCK OFF TREATED AREAS.
WARNING: KEEP OUT OF THE REACH OF CHILDREN DOMESTIC ANIMALS AND FOODSTUFFS.

CUSTOM CHEMICIDES INC.
BERKELEY, CALIFORNIA 94710

NET WEIGHT:
..... pounds

MAY CAUSE IRRITATION OF NOSE AND THROAT. MAY BE HARMFUL IF INHALED OR SWALLOWED. DO NOT BREATHE SPRAY MIST. USE WITH ADEQUATE VENTILATION. AVOID CONTACT WITH EYES, SKIN OR CLOTHING. WASH THOROUGHLY AFTER HANDLING. IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. FOR EYES, GET MEDICAL ATTENTION.

EPA Reg. No. 9319-11

Observe all cautions and limitations on labeling of all

Cameraman's Note
Poor Copy

AT-90

AMINO TRIAZOLE WEEDKILLER

Active Ingredient:

Amitrole* 90.0%

Inert Ingredients . . 10.0%

*3-amino-1,2,4-triazole

Cameraman's Note

Poor Copy

Add 1 to 2 pts. of UNIFILM 707

spreader sticker, per 100 gallons
to assure complete wetting.

VINES IN NONCROPLANDS

on Roadsides, Drainage Ditchbanks, Fencerows

When to Apply

WEED CONTROL IN NONCROPLANDS

Weeds on Roadsides, Drainage Ditchbanks, Fencerows

Amount Per Acre

When to Apply

s.
bons)
water
Apply when this growth is young and growing actively.

ils.
Treat from the time foliage is fully developed until plants begin to go dormant. For effective kill, all leaves, stems and suckers must be thoroughly wet to ground line.

water
Spot treat any regrowth.

Feb. 12, 1974

9319-11

CONTROL annual grasses and broadleaf weeds (nutgrass, grass, chestnutgrass, barnyard grass, pigweed, techie, sunflower, wild barley)	2 lbs. in 30-60 gals water	Spray when most weeds are 3-4 inches tall.
CONTROL annual grasses, annual and perennial broadleaf weeds (whiteton, Canada thistle, milkweed) SUPPRESS perennial grasses (rigid grass, quackgrass)	4 lb. in 100 gal. water	Spray when most weeds are 6-10 inches tall.
SUPPRESS PERENNIAL GRASSES (Bermuda grass quackgrass) Nutgrass (sedge)	4 lbs. in 100 gals. water	Spray when weeds are 3-6 inches tall.

Weeds on Roadsides, Fencerows with Simazine 80W

Amounts per Acre
A T. 90 Simazine

a - For seasonal control of most weeds 4 lbs. where safety to nearby desirable ornamental trees, 12 1/2 lbs. (80W)

Treat from the time foliage is fully developed until plants begin to go dormant. For effective kill, all leaves, stems and suckers must be thoroughly wet to ground line.

Spot treat any regrowth.

Feb. 12, 1974

9319-11

Apply after catkins are fully formed but before frst. Do not disturb sprayed plants. Spot treat any regrowth.

	water	
CONTROL annual grasses, annual and perennial broadleaf weeds (white top, Canada thistle, milkweed) SUPPRESS perennial grasses (rigid grass, quackgrass)	4 lb. in 100 gal. water	Spray when most weeds are 6-10 inches tall.
SUPPRESS PERENNIAL GRASSES Bermuda grass (quackgrass) Nutgrass (sedge)	4 lbs. in 100 gals. water	Spray when weeds are 3-6 inches tall.

Weeds on Roadsides, Fencerows with Simazine 80W **Amounts per Acre**
A T. 90 Simazine

a - For seasonal control of most weeds where safety to nearby desirable ornamental trees, shrubs is important.	4 lbs.	12 1/2 lbs. (80W)
b - For economical control of easy-to-kill (annual) weeds in Woods on Roadsides, Drainage Ditchbanks, Fencerows	2 lbs.	6 1/4 lbs. (80W)

AS NO RESIDUE OF AMITROLE IS PERMISSIBLE ON CROPS. DO NOT SPRAY OR PERMIT DRIFT OF SPRAY ON LAWNS, SHRUBS, FLOWERS AND OTHER DESIRABLE VEGETATION SINCE THIS PRODUCT WILL CAUSE INJURY. DO NOT USE IN GREENHOUSES Lilac, privet, honeysuckle, Boxwood, and barberry. Susceptible to this mixture and may be injured if contacted by spray.

MICIDES INC. **NET WEIGHT:**
..... **pounds**
ORNIA 94710

WARNING!
KEEP OUT OF REACH OF CHILDREN
Coarse sprays are less likely to drift.

KEEP LIVESTOCK OFF TREATED AREAS.
WARNING: KEEP OUT OF THE REACH OF CHILDREN DOMESTIC ANIMALS AND FOODSTUFFS.

MAY CAUSE IRRITATION OF NOSE AND THROAT MAY BE HARMFUL IF INHALED OR SWALLOWED. DO NOT BREATHE SPRAY MIST. USE WITH ADEQUATE VENTILATION. AVOID CONTACT WITH EYES, SKIN OR CLOTHING WASH THOROUGHLY AFTER HANDLING IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. FOR EYES, GET MEDICAL ATTENTION.

DO NOT STORE NEAR FERTILIZERS, SEEDS, OR PESTICIDES OR USE IN A GREENHOUSE. DO NOT SPRAY OR ALLOW SPRAY DRIFT TO CONTAMINATE NEARBY CROPS.

DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

IMPORTANT: DO NOT ALLOW SPRAY SOLUTION TO REMAIN IN THE SPRAYER ANY LONGER THAN NECESSARY. RINSE ALL SPRAY EQUIPMENT WITH CLEAN WATER IMMEDIATELY AFTER EACH USE.

Observe all cautions and limitations on labeling of all products used in mixtures.

Cameraman's Note
Poor Copy