


9319-11

pm25 10F2

RESTRICTED USE PESTICIDE
FOR RETAIL SALE TO AND APPLICATION ONLY BY CERTIFIED
APPLICATORS OR PERSONNEL UNDER THEIR DIRECT SUPERVISION.

ACCEPTED
AUG - 8 1986
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 9319-11



AT - 90
AMINO TRIAZOLE

WEED CONTROL IN NONCROPLANDS
WEEDS ON ROADSIDES, DRAINAGE DITCHBANKS, FENCEROWS

Contains: SPREADER
Active Ingredient:
 amitrole*90.0%
Inert Ingredients:10.0%
 *3-amino 1,2,4 triazole

DIRECTIONS FOR USE

AS NO RESIDUE OF AMITROLE IS PERMISSIBLE ON CROPS, DO NOT SPRAY OR PERMIT DRIFT OR SPRAY ON LAWNS, SHRUBS,
FLOWERS AND OTHER DESIRABLE VEGETATION SINCE THIS PRODUCT WILL CAUSE INJURY. DO NOT USE IN GREENHOUSES.
LILAC, PRIVET, HONEYBUCKLE, BOXWOOD, AND BARBERRY SUSCEPTIBLE TO THIS MIXTURE AND MAY BE INJURED IF
CONTACTED BY SPRAY.

DO NOT SPRAY OR ALLOW SPRAY DRIFT TO CONTAMINATE EDIBLE CROPS OR WATER WHICH WILL BE USED FOR IRRIGA-
TION, DRINKING OR FISHING OR OTHER DOMESTIC PURPOSES.

OBSERVE ALL CAUTIONS AND LIMITATIONS OF LABELING OF PRODUCTS USED IN MIXTURES.

COARSE SPRAYS ARE LESS LIKELY TO DRIFT.

IMPORTANT: DO NOT ALLOW SPRAY SOLUTION TO REMAIN IN THE SPRAYER ANY LONGER THAN NECESSARY. RINSE ALL
SPRAY EQUIPMENT WITH CLEAN WATER IMMEDIATELY AFTER USE.

**"It is a violation of Federal law to use this product in a
manner inconsistent with its labeling."**

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Avoid contact with skin, eyes, or clothing.
The use of this product may be hazardous to your health. This product contains amitrole which has been determined
to cause cancer in laboratory animals. Wear a respirator during mixing and loading of this product.
Wear lightweight waterproof clothing (jumpsuit or coveralls, boots or shoes, gloves, and a wide-brimmed plastic hard-
hat) when applying all products to control dense, tall vegetation. Workers applying this product in all other situations
must wear lightweight waterproof gloves and boots (or shoes).

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting immediately. Get medical attention.
IF ON SKIN: Wash thoroughly with soap and water. Remove contaminated clothing and wash before reuse.
IF IN EYES: Flush eyes with plenty of water.

NOTE TO PHYSICIAN

This product may affect thyroid function. Overexposure can be determined by monitoring thyroid function. If worker
is under treatment for hypo- or hyperthyroidism, he should be removed from spray crew until the euthyroid state is
achieved.

STORAGE AND DISPOSAL

STORAGE

Do not store near fertilizers, seeds, insecticides, or fungicides.

PESTICIDE DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
Open dumping is prohibited.

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal
law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or
Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for
guidance.

CONTAINER DISPOSAL

Do not reuse empty container. Triple rinse or equivalent. Then offer for recycling or reconditioning or puncture and
dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned,
stay out of smoke.

NET WEIGHT _____ pounds

CUSTOM CHEMICIDES

P.O. BOX 11218 • FRESNO, CA 93772
(209) 264-0441

ESTABLISHMENT NO. 9319-CA-2 EPA REGISTRATION NO. 9319-11

MADE IN U.S.A.

2 of 2

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues, and varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT SUCH LIABILITY IS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

WEEDCONTROL IN NONCROPLANDS

	Amount Per Acre	When to Apply
Weeds on Roadsides, Fencerows		
CONTROL annual grasses and broadleaf weed (foxtail, ryegrass, chestgrass, banyard grass, pigweed, lechla, sunflower, wild barley)	2 lbs. in 30 - 60 gals. water	Spray when most weeds are 3 - 4 inches tall.
CONTROL annual grasses, annual and perennial broadleaf weeds (whit-top, Canada thistle, milkweed) SUPPRESS perennial grasses (ripout, quackgrass)	4 lb. in 100 gal. water or 1 cup in 2 gals. water	Spot treat any regrowth. Spray when most weeds are 6 - 10 inches tall.
SUPPRESS PERENNIAL GRASSES Bermuda grass (quackgrass) Nutgrass (sedge)	4 lb. in 100 gal. water	Spray when most weeds are 3 - 6 inches tall.

WOODY PLANTS AND VINES IN NONCROPLANDS

	Amount Per Acre	When to Apply
Honeysuckle Kudzu Vine	5 1/2 lbs. in 125 gals.	Treat from the time foliage is fully developed until plants begin to go dormant. For effective kill all leaves, stems, and suckers must be thoroughly wet to ground line. Spot treat any regrowth.
Poison Ivy Poison Oak Western dewberry (blackberry) Big leaf maple Wild cherry Ash, Locust, Sumac	2 lbs. in 100 gals. water (3 level tablespoons) in 1 gallon water.	

Weeds on Roadsides, Fencerows with Simazine 80W	Amounts per acre	
	AT 80	Simazine
a. For seasonal control of most weeds where safety to nearby desirable trees, shrubs & ornamentals is important.	4 lbs.	12 1/2 lbs. (80 W)
b. For economical control of easy to kill (annual) weeds for Weeds on Roadsides, Fencerows	2 lbs.	6 1/2 lbs. (80 W)

ADD 1 TO 2 PINTS OF UNIFILM 707 SPREADER STICKER PER 100 GALLONS OF WATER SUGGESTED TO ASSURE COMPLETE WETTING.

A second treatment may be necessary to control weeds not thoroughly sprayed or weeds coming from dormant seeds and nuts or old roots. Apply when regrowth is young and growing actively.