

around

buildings storage areas fences pumps machinery fuel tanks rights-of-wav

roadways guard rails airports military installations highway medians pipelines railroads lumber yards industrial sites recreational areas

USTOM

ENVIRONMENTAL HAZARDS

Fish Caution: This product is toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes and containers.

> PHYSICAL OF CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

DISPOSAL

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing or burying in a safe place.

For weed control on industrial sites and noncrop areas around-the-farm.

Active Ingredients:

Total:

Prometon (2.4-bislisopro	-
pylaminol-6-inethoxy-s-	
triazine)	370
Pentachio uphenol	1.7
Other Chlorophenols	2
Petroleum Hydrocarbons	790

Inert Ingredients:

100%

WARNING! **KEEP OUT OF REACH OF CHILDREN** HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN. CAUSES SKIN IRRITATION

to not get in eyes on skin or on clothing.

STATEMENT OF PRACTICAL TREATMENT

Wash thoroughly after handling. Avoid prolonged or repeated breathing of vapor

IF ON SKIN In case of contact, flush with large amounts of water FIN EYES get medical attention IF SWALLOWED DO NOT INDUCE VOMITING.

Remove and wash contaminated clothing before re-use. tion of with adequate ventilation.

ESTABLISHMENT NO 09319-CA-1 EPA REGISTRATION NO 9319-34



SAN LEANDRO, CA 94577

CLOVIS, CA 93612

WEEDICIDE

0% 9•/0 1º/0 3 4 % 15.26 %

DIRECTIONS FOR USE

is a violation of Fadera ten to NE DROCKALT IN A dia ita indefini

Uniformly spray one gallon of Custom WEEDICIDE mixed in ten gallons of water over 1000 sq, ft. of soil surface. Thoroughly wet all foliage to insure contact action. Best results are obtained when applications are made during or soon after weed emergence while growth is succulent. Where vegetation is tall and dense, use $1\frac{1}{2}$ gallons of Custom WEEDICIDE In ten gallons of water per 1000 sq. ft. For control of hard - to - kill perennial weeds and grasses such as Johnsongrass, Bindweed and wild Carrot, use 3 - 41/2 gallons per 1000 sq. ft. Although CUSTOM WEEDICIDE has considerable activity through foliar contact, much of its activity is through the roots and therefore, adequate rainfall to move the chemical into the root zone is necessary

Under conditions where the rainfall is high and the growing season long, a second application may be necessary to maintain a weed - free soil surface

Care must be taken to keep this herbicide away from desirable flower beds, lawns and shrubs. Do not apply over roots of valuable trees or woody ornamental plants. Only apply herbicide outside of the drip line of trees.

TEPTED