

50 LBS.

NET WEIGHT


PROMICO™

FREE CHOICE NO. 9 MINERAL

With Rabon® Oral Larvicide

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENTS:

BY WEIGHT

* 2-Chloro 1-(2,4 5-trichlorophenyl) vinyl dimethyl phosphate	1.25%
INERT INGREDIENTS**	98.75%
TOTAL	100.00%

Each pound contains 5.625 g RABON

**Refers only to ingredients which are not larvicidal.

EPA REG. NO. 9329-16

EPA EST. NO. 9329-1A-1

FEED INGREDIENTS

Dicalcium Phosphate, ground yellow corn, cottonseed meal, salt, cane molasses, iron oxide, bentonite (not more than 5%), iron carbonate, sulphur, magnesium oxide, copper oxide, zinc oxide, mineral oil, D-Activated animal sterol (source of Vitamin D₃), Vitamin A acetate, artificial flavor added

GUARANTEED ANALYSIS

Crude Protein, not less than	5.00%
Calcium (Ca), maximum	11.00%
Calcium (Ca), minimum	9.00%
Phosphorus (P), minimum	9.00%
Salt (NaCl), maximum	16.00%
Salt (NaCl), minimum	14.00%
Zinc (Zn), not less than	0.20%
Magnesium (Mg), minimum	2.00%
Copper (Cu), minimum	0.50%
Iron (Fe), minimum	0.15%
Sulfur (S), not less than	9.70%
Vitamin D ₃ , minimum	50,000 U.S.P. Units Per Lb
Vitamin A, minimum	50,000 U.S.P. Units Per Lb

KEEP OUT OF REACH OF CHILDREN.

CAUTION

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

1. **PROHIBITIONS** — Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.
2. **PESTICIDE DISPOSAL** — Pesticide that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
3. **CONTAINER DISPOSAL** — Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.
4. **GENERAL** — Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Allow free choice to cattle. Cattle should consume an average of 6 oz. of the mineral per 100 pounds of body weight per month. This will supply the recommended average daily intake of 20 mg. RABON® Oral Larvicide per hundred pounds of body weight. Animals should not be started for minerals before initiation of the Promico Free Choice No. 9 Mineral with RABON® program but until all salt in any other feed after the program is started. Do not feed other free choice sources of salt or mineral containing salt in any form. Allow one mineral feeder for each 10-15 head and place near watering and feeding areas. Put out no more than one inch of supply at a time and protect from rain. If consumption of Promico Free Choice No. 9 Mineral with RABON® is above 6 oz. per 100 lbs. of body weight per month, reduce the number of feeding locations or reduce the consumption to below 6 oz. per 100 lbs. of body weight per month to reduce the number of feeding locations. Start for the Promico Free Choice No. 9 Mineral with RABON® early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall. For the Free Choice No. 9 Mineral with RABON®, only be fed up to strength and to lactating dairy cows. Do not feed to dry cows or to other animals. Do not feed to horses or to other animals. Do not feed to poultry or to other animals. Do not feed to swine or to other animals. Do not feed to fish or to other aquatic animals. Do not feed to birds or to other animals. Do not feed to reptiles or to other animals. Do not feed to amphibians or to other animals. Do not feed to insects or to other animals. Do not feed to plants or to other organisms. Do not feed to any other animals or organisms.

FEED INGREDIENTS

Dicalcium Phosphate, ground yellow corn, cottonseed meal, salt, cane molasses, iron oxide, bentonite (not more than 5%) iron carbonate, sulphur, magnesium oxide, copper oxide, zinc oxide, mineral oil, D-Activated animal sterol (source of Vitamin D₃), Vitamin A acetate, artificial flavor added

GUARANTEED ANALYSIS

Crude Protein, not less than	5.00%
Calcium (Ca), maximum	11.00%
Calcium (Ca), minimum	9.00%
Phosphorus (P), minimum	9.00%
Salt (NaCl), maximum	16.00%
Salt (NaCl), minimum	14.00%
Zinc (Zn), not less than	0.20%
Magnesium (Mg), minimum	2.00%
Copper (Cu), minimum	0.53%
Iron (Fe), minimum	0.15%
Sulfur (S), not less than	0.70%
Vitamin D ₃ , minimum	50,000 U.S.P. Units Per Lb.
Vitamin A, minimum	50,000 U.S.P. Units Per Lb.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

- 1 PROHIBITIONS** — Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.
- 2 PESTICIDE DISPOSAL** — Pesticide that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
- 3 CONTAINER DISPOSAL** — Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.
- 4 GENERAL** — Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

DIRECTIONS FOR USE: GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Allow free choice to cattle. Cattle should consume an average of 6 oz. of the mineral per 100 pounds of body weight per month. This will supply the recommended average daily intake of 70 mg. RABON® Oral Laxative per hundred pounds of body weight. Animals should not be starved for minerals before initiation of the Promico Free Choice No. 9 Mineral with RABON® program but until all salt from other feed after the program is started. Do not feed other free choice sources of salt or mineral containing salt in any form. Allow one mineral feeder for each 10-15 head and place near watering and feeding areas. Put out no more than one month's supply at a time and protect from rain.

If consumption of Promico Free Choice No. 9 Mineral with RABON® is above 6 oz. per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below 6 oz. per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding Promico Free Choice No. 9 Mineral with RABON® early in the spring as horn flies begin to appear, and continue feeding throughout the summer and into the fall.

Promico Free Choice No. 9 Mineral with RABON® may be fed up to slaughter and to lactating dairy cows.

Promico Free Choice No. 9 Mineral with RABON® prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control Promico Free Choice No. 9 Mineral with RABON® should be used in conjunction with other good management and sanitation practices.

Manufactured by

Promico, Inc., Eagle Grove, Iowa 50533

A division of North Central Holdings Corp.

SEE BACK OF TAG FOR MIXING DIRECTIONS

281

43

MIXING DIRECTIONS FOR FINISHED FEEDS

The following table shows the correct amount of **Promico Free Choice No. 9 Mineral** to be mixed into a complete daily ration in order to provide the recommended average daily intake of 70 mg of Rabon® Oral Larvicide per hundred pounds of body weight.

Determine your cattle's average body weight, find this figure in the left column. Then determine the average consumption of your complete daily ration and find this figure in the top column. Cross reference and you will find amount of **Promico Free Choice No. 9 Mineral** to be mixed into your ration. Mixing time should be adequate to assure uniform dispersion. Optimum performance can be assured only if **Promico Free Choice No. 9 Mineral** is dispersed uniformly with the feedstuff at the recommended level. Common feed mixing equipment (i.e., vertical mixers, horizontal blenders, mixer/feeder trucks) may be used to prepare formulated feed.

CONSUMPTION PER HEAD PER DAY OF FINISHED FEED (Pounds)

AVERAGE WEIGHT OF ANIMALS (Pounds)	LBS.												POUNDS OF Promico FREE CHOICE NO. 9 MINERAL WITH RABON® TO BE MIXED PER TON OF FINISHED FEED
		4#	5#	6#	7#	8#	9#	10#	15#	20#	25#	30#	
	400	25	20	17	14	13	11	10	7	5	4	4	
	500	32	25	22	18	16	14	13	9	7	5	5	
	600	37	30	25	22	19	17	15	10	8	7	5	
	700	44	35	29	25	22	20	17	12	9	7	7	
	800	50	40	33	29	25	22	20	13	10	8	7	
	900	56	45	37	32	29	25	23	15	12	10	8	
	1000	62	50	42	34	32	28	25	17	13	10	9	
	1100	68	55	46	40	34	31	28	18	14	11	10	
	1200	74	59	50	43	37	33	30	20	15	12	10	
	1300	81	65	54	47	40	37	32	22	17	13	11	
	1400	87	70	58	50	44	39	35	24	17	14	12	
	1500	84	74	62	53	47	42	37	25	19	15	13	
	1600	99	79	67	57	50	44	40	27	20	16	13	
	1700	106	85	70	60	53	47	43	29	22	17	14	
	1800	112	89	74	64	56	50	45	30	23	18	15	
	1900	118	94	79	67	59	52	47	32	24	19	16	
	2000	124	99	83	71	62	55	50	33	25	20	17	

Rabon is a registered trademark of Searl, Inc. U.S. Pat. 3,102,842