

PROMICO

BAR FLY

CATTLE MINERAL MEDICATED

Prevents the breeding of horn flies and face flies in the manure of treated cattle, when used as directed. Aids in the control of common stomach worms (*Haemonchus* spp.), lesser stomach worms (*Osteraia* spp.), hair worms, bankrupt worms (*Trichostrongylus* spp.), hook worms (*Bunostomum* spp.), nodular worms (*Oesophagostomum* spp.), in cattle and aids in the prevention of foot rot in cattle and soft tissue lump, jaw, nutritional source of iodine, when fed to cattle as directed below.

PESTICIDALLY ACTIVE DRUG	
INGREDIENT Phenothiazine *	2.65%
PESTICIDALLY INERT	
INGREDIENTS	97.25%

Total	100.00%
OTHER ACTIVE DRUG INGREDIENT	
Ethylene Diamine Dihydroiodide	0.066%
*Biologically Assayed	

GUARANTEED ANALYSIS	
Calcium (Ca), not more than	15.0%
Calcium (Ca), not less than	13.0%
Phosphorus (P), not less than	6.0%
Salt (NaCl), not more than	42.0%
Salt (NaCl), not less than	38.0%
Iodine (I), not less than	0.05%

INGREDIENTS
Dicalcium Phosphate, Defluorinated Phosphate, Steamed Bone Meal, Calcium Carbonate, Soybean Meal, Salt, Cane Molasses, Iron Sulfate, Red Iron Oxide, Manganous Oxide, Cobalt Carbonate, Magnesium Oxide, Zinc Oxide, Copper Oxide, D Activated Animal Sterol (source of Vitamin D3), Vitamin A Palmitate (stability improved), Mineral Oil, Artificial Flavor added

**CAUTION: Keep Out Of Reach
Of Children.**

Do not feed to lactating dairy animals. Occasional sensitivity to phenothiazine has been reported.

DIRECTIONS: Follow feeding directions on reverse side of this label.

Manufactured By 071272

PROMICO

EAGLE GROVE, IOWA 50531
50 POUNDS NET

FEEDING DIRECTIONS

Self-feed continuously, as the sole source of salt and mineral PROMICO BAR-FLY CATTLE MINERAL MEDICATED providing one-third ounce per day, for each 100 pounds of body weight. This is equivalent to 2 2/3 ounces per day for each 800 pound animal and, at the same time, will provide 1/4 gram of Phenothiazine per 100 lbs. of body weight, and 50 milligrams per head daily of Ethylene Diamine Dihydriodide for the prevention of foot rot in cattle and soft tissue lumpy jaw. Locate CATTLE MINERAL MEDICATED near the animal's watering or loafing area. Allow one pound per 6 head daily. Covered mineral feeders are recommended if fed in bunks. Protect from the weather.

NOTE: For maximum protection against horn flies, it is desirable to spray cattle once when first starting on a continuous feeding program with PROMICO BAR-FLY CATTLE MINERAL MEDICATED. PROMICO BAR-FLY CATTLE MINERAL MEDICATED prevents the breeding of horn flies and face flies in the manure, but does not affect any existing adult flies. Since face flies migrate from farm to farm, it is essential that the larvicide program be carried out on an area basis to assist in the reduction of the adult face fly population. The usefulness of this product in preventing the breeding of horn flies is limited to periods of adult fly activity. For internal parasite control, this product should be fed on a year round basis to cattle.