

ACTIVE INGREDIENTS:


*2-chloro-1(2,4, 5-trichlorophenyl vinyl dimethyl phosphate

INERT INGREDIENTS**

Each Pound contains 5.625 g RABON

**Refers only to ingredients which are not listed

EPA REG. NO. 9329-14
EPA EST. NO. 9329-1A-1



1.25%
98.75%
100.00%

FEED INGREDIENTS

Dicalcium phosphate, cane molasses, salt, processed grain by products, dried molasses, iron oxide, bentonite, iron carbonate, zinc oxide, manganous oxide, copper oxide, cobalt carbonate, magnesium oxide, Vitamin A acetate, D-Activated animal stero. (source of Vitamin D3), Vitamin E supplement, mineral oil, artificial flavors

GUARANTEED ANALYSIS

Calcium (Ca), maximum	12.00%
Calcium (Ca), minimum	10.00%
Phosphorus (P), minimum	10.00%
Salt (NaCl), maximum	11.00%
Salt (NaCl) minimum	9.00%
Magnesium (Mg), minimum	1.00%
Manganese (Mn), minimum	.10%
Cobalt (Co), minimum	.001%
Copper (Cu), minimum	.01%
Iron (Fe), minimum	.15%
Vitamin A, minimum	300,000 U.S.P. Units Per Lb.
Vitamin D ³ , minimum	45,000 U.S.P. Units Per Lb.
Vitamin E, minimum	150 IU/Per Lb.

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

25

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS**

CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

- PROHIBITIONS** — Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited
- PESTICIDE DISPOSAL** — Pesticide that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
- CONTAINER DISPOSAL** — Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.
- GENERAL** — Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

**DIRECTIONS FOR USE:
GENERAL CLASSIFICATION**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Allow free choice to cattle. Cattle should consume an average of 6 oz. of the mineral per 100 pounds of body weight per month. This will supply the recommended average daily intake of 70 mg RABON* Oral Larvicide per hundred pounds of body weight. Animals should not be starved for minerals before initiation of the Promico MIN-A-ROL MIX 10 with RABON* program but omit all salt from other feed after the program is started. Do not feed other free choice sources of salt or mineral containing salt in any form. Allow one mineral feeder for each 10-15 head, and place near watering and loafing areas. Put out no more than one month's supply at a time, and protect from rain.

If consumption of Promico MIN-A-ROL MIX 10 with RABON* is above 6 oz. per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below 6 oz. per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding Promico MIN-A-ROL MIX 10 with RABON* early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

Promico MIN-A-ROL MIX 10 with RABON* may be fed up to slaughter and to lactating dairy cows.

Promico MIN-A-ROL MIX 10 with RABON* prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control Promico MIN-A-ROL MIX 10 with RABON* should be used in conjunction with other good management and sanitation practices.

Manufactured by
Promico, Inc., Eagle Grove, Iowa 50533
A division of North Central Holdings Corp

"SEE BACK OF TAG FOR MIXING DIRECTIONS" 12-89
26