Kanch FOICATED -

Aids in preventing the breeding of hom flies and face flies in the manure of treated cattle, when used as directed. Aids in the center of stamachworms (Hasmonchus, Ostertagla, and Trichostrongylus spp.), nadular worms (Ossaphagastamum spp.), large-mouth bewelworms (Chabertia spp.), and hookwarms (Bunastamum spp.), in cattle when fed in accordance with the directions for use.

Aids in the prevention of foot not caused by Spherophonus Necrophonus, and of soft tissue lumpy jaw caused by Actinabacillus Lignierisi in cattle when fed as directed.



Prevents the breeding of HORN FLIES and FACE FLIES in the manure of treated cattle.

Aids in the control of INTERNAL PARASITES listed on the label. Aids in the prevention of GATTLE FOOT ROTand soft tissueLUMPY JAW.

Provides nutritional supplementation of Vitamins A & D . Xine. phosphorus and trace minerals.

Pesticidally active drug ingredient:	
***************************************	1.25%
Inert**ingradients 9	8.75%
10	0.00%
(Other) active drug ingredient:	
Ethylene Diamine Dihydriadide	0.03%
*Biologically assayed	
**Refers only to ingredients which are not insecticides.	
BHARANTEED AMALYSIS	

Crude Protein, not less than 14.0%	
Crude Fat, not less than 1.5%	
Crude Fiber, not more than 7.0%	
Calcium (Ca), not less than 2.0%	
Calcium (Ca), not more than 3.0%	
Phosphorus (P), not less than	
lodine (I), not less than	
Salt (NaC1), not less than 15.0%	
Salt (NaC1), not more than 17.0%	
Vitamin A Potency not less than 30,000 USP units per pound.	
Vitamin D3 Potency not less than 8,000 USP units per pound.	

INGUERIEMTS

Dried Cane Molasses, Cottonseed Meal, Saybean Millfeed, Cane Molasses, than the Vegetable OII, Saybean Meal, Wheat Middlings, Solt, Calcium Carbonate, Defluorinated Phosphate, Di-Calcium Phosphate, Vitamin A Palmitate (stability improved), D-Activated Animal Sterol (source of Vitamin D3), Iron Oxide, Iron Sulfote, Iron Carbonate, Zinc Oxide, Magnasium Oxide, Magnasium Oxide, Copper Oxide, Cabalt Carbonate, Natural and Artificial flavors added. PEEDING INSTRUCTIONS

RANCH-O Bar-Fly Blocks are to be self fed as a mineral supplement. Do not feed any other solt. Allow one block for each ten (10) head of cattle. The Blacks should be distributed over a large area to allow livestock good access to blacks, Legate the supplement near the animals' water, or near their loafing area. RANCH-O Bar-Fly Blocks eaten at the rate of 3/4 ounces daily per 100 lbs. of body weight will supply the minimum amount of phenothiazine necessary to aid in the control of the breeding of face files and horn flies and to aid in the control of certain gastrointestinal parasites such as common stamach worms, nadular worms, and hookwarms af cottle.

When eaten at the rate of 6 aunces per head per day, cattle will receive 50 mgs. of Ethylene Diamine Dihydriodide daily, which aldsin the prevention of foot rot in cattle and soft tissue lumpy jaw.

For maximum protection against hom flies, cattle should be sprayed at the start of the fly season, and fed RANCH-O Bar-Fly continuously, beginning several weeks before the fly season, until all adult hom flies have disappeared in the fall. RANCH-O Bar-Fly prevents the preeding of horn flies and face flies in the minure of treated cattle b "Since face flies in grate from farm to farm it is essential that the larvicide program be carried out adult tace fly pay listing."

CONSULT YOUR VESERINARIAN FOR ASSISTANCE IN THE DIAGNOSIS, TREATMENT AND CONTROL OF PARASITISM.

WARNING: Keep Out Of Reach Of Childhug **4** 1977

THIS PRODUCT IS NOT TO BE FED TO LACTATING BAIRY AHIMALS. OCCA PHENGINIAZINE HAS BEEN REPORTED IN CATTLE.

A MANTHULY Effecticide. Fungicide, and Rodenticide Act. cm amended, for the pesticide registered under 9374 -

MANUFACTURED AND GUARANTEED B

DAGLAND MILLS. INCLEASE No.

"Home of RANCA-O Blocks and RANCA-O Mineral"

EPA EST. 09374M001 EPA REG. No. 9374-1

ROUTE 2 BOX 55A NEOSHO, MISSOURI 64850

PHONE AC 417 451-2510