
P.M14 9374-2

# Ranch-O BAR-FLY 16.2 MINERAL SUPPLEMENT BLOCK 16.2 MEDICATED

Aids in preventing the breeding of hom files and face riles in the manure of treated cettle, when used as directed. Aids in the sentral of stamachwarms (Hossenchus, Ostertagia and Trichastrongylus spp.), nadular warms (Ossephagostamum spp.), large - mouth bawel warms (Chabertia spp.) and hookwarms (Bunasiamum spp.), in cattle when fed in accordance with the directions for use.



phosphorus and trace minerals.

-			
	Pesticidally active drug ingredient:	1.2	5%
i	Inert** ingredients	98.75	5%
1		100.00%	
Į	*Biologically assayed		
1	**Refers only to ingredients which are not insecticides.		
	, , , , , , , , , , , , , , , , , , ,		
	CUARANTEED ANALYSIS		
	Crude Protein, not less than	5.0	%
	Crude Fat, not less than		
	Crude Fiber, not more than		
Į	Calcium (Ca), not less than		
	Calcium (Ca), not more than		
1	Phosphorus (P), not less than		
1	lodine (I), not less than		
	Salt (NaC1), not less than	18.00	%
	Salt (NaC1), not more than	20.00	%
	Fitamin A Potency not less than 30,000 USP units per pound.  Fitamin D3 Potency not less than 8,000 USP units per pound.		
į			
	otal Sugars (as invert), not less than 15.00 %		
, ;	Sugars derived from 25% Liquid Cone Molasses, 1		
	Cone Molasses	J D.	

#### MPREBIENTS

Dried Cane Molasses, Cottoneed Meal, Soybean Millfeed, Cone Molasses, Animal and Vegetable Fat, Soybean Meal, Wheat Middlings, Sait, Calcium Corbonate, Defluorinated Phosphate, Di-Calcium Phosphate, Vitamin A Palmitate (stability improved), D-Activated Animal Steral (source of Vitamin D3), Iron Oxide, Iron Sulfate, Iron Carbonate, Zinc Oxide, Manganous Oxide, Magnesium Oxide, Capper Oxide, Cabalt Carbonate, Natural and Artificial flavors added.

#### FEEDING MISTRUCTIONS

RANCH-O Bar-Fly Blocks are to be self fed as a mineral supplement. Do not feed any other solt. Allow one block for each ten (10) head of cattle. The blocks should be distributed over a large area to allow livestock good access to blocks. Locate the supplement near the animals' water or near their loafing area. RANCH-OBar-Fly Blocks eaten at the rate of 3/4 ounces daily per 100 lbs. of body weight will supply the minimum amount of phenothiozine necessary to aid in the control of the breeding of face flies and horn flies and to aid in the control of certain gastrointestinal parasites such as common stomach warms, nodular warms and hookwarms of cattle.

For maximum protection against hom flies, cattle should be sprayed at the start of the fly season and fed RANCH-O Rec-Fly continuously, beginning several weeks before the fly season, until all adult horn flies have disappeared in the fall. RANCH-O Bar-Fly prevents the breeding of hom flies and face flies in the manure of treated cattle but does not affect any existing adult flies. "Since face flies migrate from farm to farm it is essential that the larvicide program be carried out on an area basis to assist in reduction of the adult face fly population."

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

### <u>WARNING</u>

1

## KEEP OUT OF THE REACH OF CHILDREN. DO NOT FEED TO LACTATING DAIRY ANIMALS.

Individual animals are occasionally sensitive to phenothiazine. Certain parasites may become resistant to phenothiazine therapy.

RMII-

MANUFACTURED AND GUARANTEED BY

RAGEAND MILLS, INC.

"Home of Ranch-O Blocks and Ranch-O Mineral"

EPA EST. 09374M001 EPA REG. No. 9374-2

ROUTE 8 BOX 168 NEOSHO, MISSOURT 64850

PHONE 417-451-2510

MAR 30 1979
Und of and Insecticide,
Fungi 110 and Fred Act,
On Care 1

10/