BU15 9374-2-1

Ranch-O BAR-FLY NO.2 MINERAL SUPPLEMENT BLOCK NO.2

Aids in preventing the breeding of horn flies and face flies in the manure of treated cattle, when used as directed. Aids in the control of stamachwarms (Haemanchus, Ostertagia and Trichostrongylus app.), nadulasyroms (Oesaphagostamum app.), large - mouth bowel worms (Chabertia app.) and haokwarms (Bunastamum app.), in cattle when led in accordance with the directions for use.

ACCE Support of the breeding of HORN FLIES and FACE FLIES in the manure of treated cattle. Aids in the control of INTERNAL PARASITES listed on the label. Provides nutritional supplementation of Vitamins A & D phosphorus and trace minerals.	Pessicially active drug ingredient: Thenothicrine* *Biologically assysed *COUNTRICTION TO THE ANALYSIS *Biologically assysed *COUNTRICTION TO THE ANALYSIS *Biologically assysed *COUNTRICTION TO THE ANALYSIS *COUNTRICTION TO THE ANALYSI
phosphorus and trace minerals.	Total Sugars (as invert), not less than

WEREDIENTS

Dried Cane Malasses, Cottonseed Meal, Soybean Millfeed, Cane Malasses, Animal and Vegetable Fat, Soybean Meal, Wheat Middlings, Salt, Calcium Carbonate, Defluorinated Phosphate, Di-Calcium Phosphate, Vitamin A Palmitate (stability improved), D-Activated Animal Sterol (source of Vitamin D3), Ethylene Diamine Dihydriodide, Iron Oxide, Iron Sulfate, Iron Carbonate, Zinc Oxide, Manganous Oxide, Magnesium Oxide, Copper Oxide, Cobalt Carbonate, Natural and Artificial flavors added.

FEEDING WSTRUCTIONS

RANCH-O Bor-Fly Blocks are to be self fed as a mineral supplement. Do not feed any other salt. Allow one block for each ten (10) head of cottle. The blocks should be distributed over a large area to allow livestock good access to blocks. Locate the supplement near the animals: water or near their loafing area. RANCH-O Bor-Fly Blocks eaten at the rate of 3/4 ounces doily per 100 lbs. of body weight will supply the minimum amount of phenothiazine necessary to aid in the control of the biceding of ince and hornflies and to aid in the control of certain gastraintestinal parasites such as common stomach worms, nodular worms and worksmaps of cattle.

For maximum protection against horn flies, cattle should be sprayed at the start of the fly season and fed RANCH-O Bar-Ffy continuously, beginning several weeks before the fly season, until all adult horn flies have disappeared in the fall. RANCH-O Bar-Ffy prevents the breeding of horn flies and face flies in the manure of treated cattle but does not affect any existing adult flies. "Since face flies migrate from farm to farm it is essential that the farvicide program be carried out on an area basis to assist in reduction of the adult face fly population."

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

WARNING KEEP OUT OF THE REACH OF CHILDREN. DO NOT FEED TO LASTATING DAIRY ANIMALS.

Individual animals are occasionally sensitive to phenothiazine. Certain parasites may become resistant to phenothiazine therapy.

MANUFACTURED AND GURANTED BY
DAGLAND MIRES: BYC

EPA EST. 09374M001 EPA REG. No. 9374-2

"Home of Ranch-O Blocks and Ranch-O Mineral"

ROUTE 8 BOX 168 NEOSHO, MISSOURI 64850

PHONE 417-451-2510



10//