

9374-8

06/13/1979

1/2

NET WEIGHT 33.3 LBS.

# Ranch-O FLY BLOCK

## WITH RABON® ORAL LARVICIDE

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENTS:	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl)	
vinyl dimethyl phosphate.....	0.30%
INERT INGREDIENTS:*	99.70%
TOTAL.....	100.00%

\* Refers only to ingredients which are not larvicidal.

EPA REG. NO. 9374-8  
EPA EST. NO. 09374M001

RABON® is the registered trademark of Shell Chemical Company, U.S. Patent 3,102,842

### FEED INGREDIENTS:

Cane molasses, soy millfeed, soybean meal, wheat middlings, dicalcium phosphate, low fluorine phosphate rock, lignin sulphate, bentonite, salt, magnesium oxide, calcium carbonate, ethylene diamine dihydriodide, cobalt carbonate, zinc oxide, sulfur, copper oxide, iron oxide, manganese oxide, Animal and Vegetable Fat, BHA (a preservative), Vitamin A Supplement, D-Activated Animal Sterol (Source of Vitamin D3), Artificial flavors added.

### GUARANTEED ANALYSIS:

Crude Protein, minimum.....	5.00%
Crude Fat, minimum.....	1.50%
Crude Fiber, maximum.....	11.00%
Calcium (Ca), maximum.....	5.50%
Calcium (Ca), minimum.....	4.50%
Phosphorus (P), minimum.....	1.00%
Salt (NaCl), maximum.....	21.00%
Salt (NaCl), minimum.....	18.00%
Iodine (I), minimum.....	.003%
Vitamin D3, minimum.....	7,500 U.S.P. Units Per Lb.
Vitamin A, minimum.....	30,000 U.S.P. Units Per Lb.

**ACCEPTED**

**JUN 13 1979**

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 9374-8

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**SEE REVERSE SIDE OF TAG FOR USE DIRECTIONS**

2/2

RANCH-O FLY BLOCK WITH RABON ® ORAL LARVICIDE

---

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
CAUTION**

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

**STORAGE AND DISPOSAL**

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

---

DIRECTIONS FOR USE:  
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Allow free choice to cattle. Cattle should consume an average of 0.8 ounces of the Fly Block per 100 pounds of body weight per day, which is equivalent to 8 ounces for a 1000 lb. animal daily. This will supply the recommended average daily intake of 70 mgs. RABON ® Oral Larvicide per hundred pounds of body weight. Animals should not be starved for minerals before initiation of the Ranch-O Fly Block with RABON ® program but OMIT all salt from other feed after the program is started. Do not feed other free choice sources of salt or mineral containing salt in any form. Allow one block for each 10 head of cattle and place near watering and loafing areas. Put out no more than two weeks supply at a time.

If consumption of Ranch-O Fly Block with RABON ® is above 0.8 ounce per 100 lbs. of body weight per day, reduce the number of feed locations or relocate. If consumption is below 0.8 ounce per 100 lbs. of body weight per day, increase the number of feeding locations. Start feeding Ranch-O Fly Block with RABON ® early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

Ranch-O Fly Block with RABON ® may be fed up to slaughter and to lactating dairy cows.

Ranch-O Fly Block with RABON ® prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control Ranch-O Fly Block with RABON ® should be used in conjunction with other good management and sanitation practices.

Manufactured by: RAGLAND MILLS, INC.  
NEOSHO, MO. 64850  
(417) 451-2510