

With Rabon[®] Oral Larvicide

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle and horses.

Active Ingredient	BY WEIGHT
Tetrachlorvinphos: 2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate.....	0.30%
Inert Ingredients	99.70%*
TOTAL	
	100.00%

* Refers only to ingredients which are not larvicidal.

EPA REG. NO. 9374-8
EPA EST. NO. 09374M001

GUARANTEED ANALYSIS:

Crude Protein, minimum.....	5.00%
Crude Fat, minimum.....	1.50%
Crude Fiber, maximum.....	11.00%
Calcium (Ca), maximum.....	5.50%
Calcium (Ca), minimum.....	4.50%
Phosphorus (P), minimum.....	1.00%
Salt (NaCl), maximum.....	21.00%
Salt (NaCl), minimum.....	15.00%
Iodine (I), minimum.....	0.003%
Vitamin D3, minimum.....	7,500 I Units Per Lb.
Vitamin A, minimum.....	30,000 I Units Per Lb.

FEED INGREDIENTS:

Cane molasses, soy millfeed, soybean meal, wheat middlings, dicalcium phosphate, lignin sulphate, bentonite, salt, magnesium oxide, calcium carbonate, ethylene diamine dihydriodide, cobalt carbonate, zinc oxide, sulfur, copper oxide, iron oxide, manganese oxide, animal and vegetable fat, BHA (a preservative), vitamin A supplement, D-activated animal steral (source of vitamin D3), artificial flavors added.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE REVERSED SIDE OF TAG FOR ADDITIONAL
PRECAUTIONARY STATEMENTS AND USE DIRECTIONS.

Manufactured by

Ragland Mills, Inc.

Rt 8, Box 168

Neosho, MO 64850

(417) 451-2510

NOTE - Label Draft identical with that submitted to EPA on March 18, 1991.
[Site of Application request - horses]

ACCEPTED

JUN 19 1991

U.S. DEPARTMENT OF AGRICULTURE
Animal and Plant Health Inspection Service
Pesticide Administration
Pest. No. 9374-8

BEST AVAILABLE COPY

W259 P81

FEB-21-'91 THU 17:55 ID:RAGLAND MILLS

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, get medical attention. Wear long-sleeved shirt and pants; chemical resistant gloves; shoes and socks for protection when handling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water when disposing of equipment wash water.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a dry place in original container.

Disposal: Completely empty container into application (mixer/feeder) equipment. Then dispose of empty bag in a sanitary landfill or by incineration or if allowed, by State and local authorities by burning. If burned, stay out of smoke. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Allow free choice to cattle and horses. Cattle and horses should consume an average of 0.8 ounces of the Fly Block per 100 pounds

of body weight per day, which is equivalent to 8 ounces for a 1000 lb. animal daily. This will supply the recommended average daily intake of 70 mgs. RABON Oral Larvicide per hundred pounds of body weight. Animals should not be starved for minerals before initiation of the Ranch-O Fly Block with RABON program but OATMEAL salt from other feed after the program is started. Do not feed other free choice sources of salt or mineral containing salt in any form. Allow one block for each 10 head of cattle or horses and place near watering and loafing areas. Put out no more than two weeks supply at a time.

If consumption of Ranch-O Fly Block with RABON is above 0.8 ounce per 100 lbs. of body weight per day, reduce the number of feed locations or relocate. If consumption is below 0.8 ounce per 100 lbs. of body weight per day, increase the number of feeding locations. Start feeding Ranch-O Fly Block with RABON early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall

Ranch-O Fly Block with RABON may be fed up to slaughter and to lactating dairy cows.

Ranch-O Fly Block with RABON prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, or horses, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in enclosed cattle or horse lots and buildings to control adult house flies and stable flies which can breed not only in cattle and horse manure but in other decaying vegetable matter or sludge on the premises.

In order to achieve optimum fly control Ranch-O Fly Block with RABON should be used in conjunction with other good management and sanitation practices.

BEST AVAILABLE COPY

#259 P02

FEB-21-'91 THU 17:56 IDIRAGLAND MILLS