

WARNING: KEEP OUT OF REACH OF CHILDREN. See other warnings on back panel

PROLIN

NET WT 5 LBS

PRICE \$2.90



**Rat & Mouse
KILLER**

Ready-to-Use

NEW IMPROVED WARFARIN

ACTIVE INGREDIENTS		
3-(o-acetylbenzyl)-4-hydroxycoumarin		0.025%
N-(2 Quinoxaliny) Sulfanilamide		0.025%
INERT INGREDIENTS		99.950%
	TOTAL	100.000%

TERMA is a Trademark of the Albany Pest Control Company
PROLIN Warfarin is a registered trademark of the Albany Pest Control Company
Albany Pest Control Company, Albany, Oregon 97321

PACKED BY XTERMA PEST CONTROL ALBANY, OREGON

USDA REG. NO. 9392-1

DIRECTIONS FOR RATS: Place not less than 1/4 to 1/2 pound of XTERMA bait in a shallow container in recognized runways or near rat holes or burrows. An uninterrupted supply of bait should be maintained for at least 10 days.

FOR MICE: The same general directions should be fol-

ACTIVE INGREDIENTS:		
3-(o-acetylbenzyl)-4-hydroxycoumarin		0.025%
N-(2 Quinoxaliny) Sulfanilamide		0.025%
INERT INGREDIENTS		99.950%
	TOTAL	100.000%

*PROLIN is a Trademark of the Wisconsin Alumni Research Foundation.
 PROLIN Warfarin plus an antibacterial agent is manufactured under license from the Wisconsin Alumni Research Foundation, U.S. Patent Nos. 2,477,775, 2,477,776, 2,477,777, 2,477,778, 2,477,779, 2,477,780, 2,477,781, 2,477,782, 2,477,783, 2,477,784, 2,477,785, 2,477,786, 2,477,787, 2,477,788, 2,477,789, 2,477,790, 2,477,791, 2,477,792, 2,477,793, 2,477,794, 2,477,795, 2,477,796, 2,477,797, 2,477,798, 2,477,799, 2,477,800, 2,477,801, 2,477,802, 2,477,803, 2,477,804, 2,477,805, 2,477,806, 2,477,807, 2,477,808, 2,477,809, 2,477,810, 2,477,811, 2,477,812, 2,477,813, 2,477,814, 2,477,815, 2,477,816, 2,477,817, 2,477,818, 2,477,819, 2,477,820, 2,477,821, 2,477,822, 2,477,823, 2,477,824, 2,477,825, 2,477,826, 2,477,827, 2,477,828, 2,477,829, 2,477,830, 2,477,831, 2,477,832, 2,477,833, 2,477,834, 2,477,835, 2,477,836, 2,477,837, 2,477,838, 2,477,839, 2,477,840, 2,477,841, 2,477,842, 2,477,843, 2,477,844, 2,477,845, 2,477,846, 2,477,847, 2,477,848, 2,477,849, 2,477,850, 2,477,851, 2,477,852, 2,477,853, 2,477,854, 2,477,855, 2,477,856, 2,477,857, 2,477,858, 2,477,859, 2,477,860, 2,477,861, 2,477,862, 2,477,863, 2,477,864, 2,477,865, 2,477,866, 2,477,867, 2,477,868, 2,477,869, 2,477,870, 2,477,871, 2,477,872, 2,477,873, 2,477,874, 2,477,875, 2,477,876, 2,477,877, 2,477,878, 2,477,879, 2,477,880, 2,477,881, 2,477,882, 2,477,883, 2,477,884, 2,477,885, 2,477,886, 2,477,887, 2,477,888, 2,477,889, 2,477,890, 2,477,891, 2,477,892, 2,477,893, 2,477,894, 2,477,895, 2,477,896, 2,477,897, 2,477,898, 2,477,899, 2,477,900, 2,477,901, 2,477,902, 2,477,903, 2,477,904, 2,477,905, 2,477,906, 2,477,907, 2,477,908, 2,477,909, 2,477,910, 2,477,911, 2,477,912, 2,477,913, 2,477,914, 2,477,915, 2,477,916, 2,477,917, 2,477,918, 2,477,919, 2,477,920, 2,477,921, 2,477,922, 2,477,923, 2,477,924, 2,477,925, 2,477,926, 2,477,927, 2,477,928, 2,477,929, 2,477,930, 2,477,931, 2,477,932, 2,477,933, 2,477,934, 2,477,935, 2,477,936, 2,477,937, 2,477,938, 2,477,939, 2,477,940, 2,477,941, 2,477,942, 2,477,943, 2,477,944, 2,477,945, 2,477,946, 2,477,947, 2,477,948, 2,477,949, 2,477,950, 2,477,951, 2,477,952, 2,477,953, 2,477,954, 2,477,955, 2,477,956, 2,477,957, 2,477,958, 2,477,959, 2,477,960, 2,477,961, 2,477,962, 2,477,963, 2,477,964, 2,477,965, 2,477,966, 2,477,967, 2,477,968, 2,477,969, 2,477,970, 2,477,971, 2,477,972, 2,477,973, 2,477,974, 2,477,975, 2,477,976, 2,477,977, 2,477,978, 2,477,979, 2,477,980, 2,477,981, 2,477,982, 2,477,983, 2,477,984, 2,477,985, 2,477,986, 2,477,987, 2,477,988, 2,477,989, 2,477,990, 2,477,991, 2,477,992, 2,477,993, 2,477,994, 2,477,995, 2,477,996, 2,477,997, 2,477,998, 2,477,999, 2,478,000.

PACKED BY XTERMA PEST CONTROL ALBANY, OREGON

USDA REG. NO. 9392-1

DIRECTIONS FOR RATS: Place not less than $\frac{1}{4}$ to $\frac{1}{2}$ pound of XTERMA bait in a shallow container in recognized runways or near rat holes or burrows. An uninterrupted supply of bait should be maintained for at least 10 days.

FOR MICE: The same general directions should be followed except placements should be in tablespoon amounts ($\frac{1}{4}$ to $\frac{1}{2}$ ounce). Bait placements should be more frequent (8 to 10 foot intervals) as mice have a limited range of travel. An uninterrupted supply of bait should be maintained for at least 15 days.

CAUTION: Baits should be placed in locations not accessible to children, pets and domestic animals or in tamper-proof bait boxes, cages or other suitable means such as wide boards slanted over baits and against an adjacent wall. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains an antibacterial agent which inhibits bacteria that produce vitamin K, an antidote for warfarin. It hereby increases effectiveness against hard-to-kill rats and mice and gives better and more thorough control.

Baits are consumed regularly with no signs of "bait shyness." Rodents continue to eat until they die. **KILL TIME** averages 5-8 days and over 2-3 weeks whole colonies are destroyed. For satisfactory control, baiting should be maintained continuously as long as evidence of feeding is observed.

Baits should be protected from the weather. Wet, mouldy or weevil infested baits should be replaced with fresh bait.