SUNNILAB

-LOWER POLICY

. . .

s 500 m

(m) D1-3:

6.5

. 4 94 - 1

SUNNILAND FLOWER POWER

8-8-8 Plant Food with Systemic Insecticide

Florida Registration No.

MINIMUM GUARANTEED ANALYSIS

Total Nitrogen		8.00%
Nitrate Nitrogen	0.00%	
Ammoniacal Nitrogen	6.00%	
Water Soluble Organic		
Nitrogen	1.75%	
Water Insoluble Nitrogen	.25%	
Available Phosphoric Acid		8.00%
Water Soluble Potash		8.00%
Total Available Primary Plant	Food	24.00%
Chlorine, not more than		7.00%

Primary Plant Food Sources: Sulphate of Ammonia, Urea, Urea Formaldehyde, Ammoniated Superphosphate, Triple Superphosphate, Muriate of Potash, Sulphate of Potash Magnesia.

Secondary Plant Foods:			
Total Magnesium	as	MgO	1.00%
Water Sol Mag	as	MgO	1.00%
Iron	as	Fe ₂ 0 ₃	0.50%
Manganese	as	MnŌ	0.09%

Secondary Plant Food Sources: Sulphate of Potash Magnesia, Iron Sulphate, Manganese Sulphate.

Ingredient Statement as an Insecticide:

Active Ingredients

0,0-diethyl S-(Z-(ethylthio)ethyl) phosphorod; thioate* 1%
Inert Ingredients

99%

***U.S.P**at. No. 2,759.010

Equivalent 20 lbs Di-Systor per ton.
(R) Di-Syston is a registered trademark of Farbenfabriken Bayer A.G., Chemagro Corporation

UNDER THE FEE CHAL INSECTIONS
UNDER THE FEE CHAL INSECTIONS
POR SEC SOME POINTS REGISTERS
UNDER NO 9404-3

$$P_{P_{n}}(x) = \frac{1}{2} \left(\frac{1}{2} \sum_{i=1}^{n} \frac{1}{2} \sum_{i=1}^{n$$

. :



Sunniland Flower Power Copy for left panel

Sunniland Flower Power Rose & Ornamental Care is a complete fertilizer containing all necessary essential plant food elements combined with a systemic insecticide to control sucking insects. This insection is absorbed through the root of the plants along with the food and protects the plant from within. It can not be washed off and will protect new shoots as well as underside of leaves.

DIRECTIONS

When to apply:

Start regular program of fertilizing as needed and continue at six week intervals. When used as directed, this material will furnish adequate plant food and protect plants from sucking insects for six weeks or longer.

For Fungus or Black Spot on Roses: Use Sunniland Fore or Sunniland Coppercide.

How to Use:

On established roses, spread 1/3 cup (2 ozs) evenly around the base of each plant and work into top 1 to 2 inches of soil. Water thoroughly. Avoid contact with hands.

When planting new roses, place 1/4 cup in hole and mix with prepared soil. Apply 1/4 cup around plant when hole is filled, working into top 2 inches. Water Well. Additional fertilizer will not be needed for six weeks.

For Annuals (petunias, snapdragons, calendulas, marigolds, zimias etc), bulbs, flower beds. Apply 2-3 lb. per each 100 sq. ft. of bed area. Work into top 1 to 2 inches of soil. Plant sends or set plants and water thoroughly. For individual plants, use I teaspoonful around each plant and water well.

On Potted Plants: Use one level teaspoonful per six inch pot and scratch into top soil. Water well after application.

On Ornamental Shrubs: Apply 2/3 cupful (4 oz.) for each foct of shrub height. Distribute the required amount evenly from the base of the plant to the drip line all around the plant. Work into the top 1 to 2 inches of soil and water thorquefuly.