

Kelthane<sup>®</sup> EC is compatible with most other commonly used insecticides and fungicides and may be used in regular or concentrated sprayers. It should not be used with lime or material not compatible with solvent based insecticides.

**CAUTION**

Harmful if swallowed. Avoid breathing of spray mist and skin contact. In case of contact, wash thoroughly with soap and water. Avoid contamination of feed or food products. Do not smoke while spraying. Keep out of reach of children. Do not use on humans or pets. In case of contact with eyes call physician and flush with water for 15 minutes. Keep away from heat or open flame. Do not reuse container. Destroy when empty.

**FISH AND WILDLIFE CAUTION:**

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME**

**WARRANTY:**

Conditions of use are beyond seller's control. Seller's liability is limited to products replacement or purchase price refund. Seller makes no warranty, express or implied, concerning products or beyond claims on label.

U.S. Patent Nos. 2,812,280, 2,812,302.

Patented by Chase & Sanford Co.

EPA Reg. No. 1404-4  
EPA Est. No. 1404 FL 1

© 1974 Chase & Sanford Co.



**CONTROLS  
RED SPIDER  
AND OTHER  
PLANT MITES**

**KELTHANE<sup>®</sup>  
EC  
MITICIDE**

<b>ACTIVE INGREDIENTS</b>	
1,1 Bis (chlorophenyl) ...	18.5%*
2,2 trichloro-ethanol ...	72.0%
Xylene ...	3.5%
<b>INERT INGREDIENTS</b> ...	100.0
<b>TOTAL</b> ... 100.0	

\*Equivalent to 1.6 lbs. Active Ingredient Per Gallon

**CAUTION: Keep Out of Reach of Children**  
See left panel for additional cautions.

Kelthane<sup>®</sup> EC gives high initial kill and against most species of agricultural mites: red, red spider, six-spotted, Pacific, privet, citrus red (purple), Schoene, y plum, peach silver, clover, tomato tropical and Yuma. Kelthane EC is a sp not kill bees and beneficial insect predit

**DIRECTIONS FOR USE**

Wet foliage thoroughly using sufficient parts of the plant, especially the undersi

<b>Recommended On</b>	<b>To Control</b>
Ornamentals, Flowers, Nursery Stock and shade trees and Orchids	Control of Mites common on green- house or out- door plants.

Changes, Grape- fruit, Lemons, Limes and Other Citrus.	Citrus red (purple) Citrus flat, six spotted, and Rust mites.
---	--

Vegetables, tomatoes, cucum- bers, squash, eggplant, peppers, and melons.	Two-spotted, Pacific, tropical and tomato russet mites.
---	--

**NOTE:** For large sprayers the rate is gal. water.

**NET WEIGHT 8 OUN**



Manu  
CHASE  
Sanford


most other commonly used miticides should be used in regular rotation. Do not be used with lime or other insecticides.

Use of spray mist and wash thoroughly with water before use. Keep out of reach of children. In case of contact with skin, wash with water for 15 minutes. Do not reuse container.

Do not use near lakes, streams and ponds. Do not apply to occur. Do not apply to areas treated. Do not use on equipment, or dis- Do not use as specified on this label.

DO NOT USE NEAR HEAT OR

Control. Seller's liability limited to purchase price refund. No warranty implied, concerning



**CONTROLS  
RED SPIDER  
AND OTHER  
PLANT MITES**

# KELTHANE<sup>®</sup> EC MITICIDE

ACTIVE INGREDIENTS:	
1, 1-Bis (chlorophenyl)-	
2, 2 trichloro-ethanol	18.5%*
Xylene	73.0%
INERT INGREDIENTS	8.5%
TOTAL 100.0%	

\*Equivalent to 1.0 lbs. Active Ingredient Per Gallon.

**CAUTION: Keep Out of Reach of Children.**  
See left panel for additional cautions.

Kelthane<sup>®</sup> EC gives high initial kill and long residual activity against most species of agricultural mites including European red, red spider, six-spotted, Pacific, strawberry, cyclamen, privet, citrus red (purple), Schoene, yellow McDaniel, rust, plum, peach silver, clover, tomato russet, spruce, desert, tropical and Yuma. Kelthane EC is a specific miticide and will not kill bees and beneficial insect predators.

**DIRECTIONS FOR USE**

Wet foliage thoroughly using sufficient gallonage to cover all parts of the plant, especially the underside of leaves.

Recommended On	To Control	Use
Ornamentals, Flowers, Nursery Stock and shade trees and Orchids.	Control of Mites common on green- house or out- door plants.	1-2 Tablespoons in 3 gal. water. Apply when mites are present, repeat when necessary.
Oranges, Grape- fruit, Lemons, Limes and Other Citrus.	Citrus red (purple) Citrus flat, six spotted, and Rust mites.	1-2 Tablespoonsful in 3 gal. water. Do not spray within 7 days of harvest of fruit.
Vegetables, tomatoes, cucum- bers, squash, eggplant, peppers, and melons.	Two-spotted, Pacific, tropical and tomato russet mites.	1-2 Tablespoonsful in 3 gal. water. Apply at first sign of mite build up and repeat as necessary. Do not spray within 7 days of harvest.

NOTE: For large sprayers the rate is 1 1/2 to 2 pints per 100 gal. water

**NET WEIGHT 8 OUNCES**



Manufactured For  
**CHASE & COMPANY**  
Sanford, Florida 32771