

SHAKE WELL BEFORE USING.

GENERAL DIRECTIONS FOR USE ON VEGETABLE INSECTS

Apply Sunniland SEVIN Insect Spray when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surfaces of all foliage for complete coverage. Repeat as necessary, generally at 7-day intervals. Use one tablespoon to one gallon water. Use 5 to 7 oz. to 10 gallons of water for control of insects listed below.

BEANS: (green beans, lima beans, snap beans, cowpeas, black-eyed) To control Mexican bean beetle, bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhopper, velvetbean caterpillar, western bean cutworm, armyworm, corn earworm, stink bugs and tarnished plant bugs.

CUCUMBER, MELONS, PUMPKIN, SQUASH: To control pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers and squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on watermelons in Florida.

POTATO, TOMATO, EGGPLANT, PEPPER: To control Colorado potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworms, lace bugs, tomato fruit worm, tomato hornworms, tarnished plant bugs, stink bugs.

OKRA: To control corn earworms and stink bugs.

HEAD LETTUCE, CARROTS, CHINESE CABBAGE, COLLARDS, GARDEN BEETS, KALE, MUSTARD GREENS, PARSLEY, RADISHES, RUTABAGAS, SPINACH, SWISS CHARD AND TURNIPS: To control flea beetles, harlequin bugs, leafhoppers, armyworms, corn earworms, spittlebugs, tarnished plant bugs and stink bugs.


Do not apply to Collards, Chinese cabbage, beet greens, kale, mustard greens, parsley, spinach, swiss chard and turnip tops within 14 day period of harvest.

Do not treat radishes, rutabagas, head lettuce or turnips (if tops are not to be used for food) within 3 days of harvest. Do not apply to beets within 3 days of harvest.

EPA NO. 9404-9

CP-6M-1354-6M-3-73

Contains 2 pounds Sevin per gallon.



Insecticide for
Lawns, Vegetables,
and Ornamental Plants

SEVIN®

INSECT SPRAY

ACTIVE INGREDIENTS:	
Carbaryl (1-naphthyl N-methyl-carbamate).....	23.3%
INERT INGREDIENTS.....	76.7%
TOTAL	100.0%

Caution: Keep Out of Reach of Children
SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS.

Sunniland
ornament
gladiolus,
elm, map

Apply to
to contro
beetles, b
gypsy mo
aphid, ros
and willow

DIRECTI
off Use 5

Lawn Inse
ticks, leaf
of water
500 sq. ft.

SEVIN®
are encour
bees expo
tion may
Service. In
plants are
following

CAUTION
breathing
feed or fo
contamina
cover on
thoroughly
WARAN
liability is
Seller mak
beyond cla

® Tradema
U.S. Pat



ACCEPTED

SECTS

age appear.
reat upper
Repeat as
one gallon
s listed be-

e-ved) To
etles, flea
stern bean
plant bugs.

ckleworm,
uash bugs.
tments are
onged high


do potato
myworms,
plant bugs,

LLARDS,
ADISHES
o control
earworms,

, mustard
day period

are not to
ets within

per gallon.



Insecticide for

Lawns, Vegetables,

and Ornamental Plants

SEVIN®

INSECT SPRAY

ACTIVE INGREDIENTS:	
Carbaryl (1-naphthyl N-methyl-carbamate).....	23.3%
INERT INGREDIENTS.....	76.7%
TOTAL	100.0%

Caution: Keep Out of Reach of Children
SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS.

ORNAMENTAL INSECT CONTROL

Sunniland SEVIN® Insect Spray can be used on a wide range of ornamental plants, flowers, shrubs and small trees, such as rose, carnation, gladiolus, chrysanthemum, lilac, arbor vitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. Do not use on Boston ivy.

Apply to upper and lower leaf surface of all foliage for complete coverage to control such pests as: apple aphid, bagworm, birch leaf miner, blister beetles, boxwood leaf miner, elm leaf aphid, elm leaf beetle, flea beetles, gypsy moth, Japanese beetle, June bugs, puss caterpillars, psyllids, rose aphid, rose-slug, scale insects, tent caterpillars, thorn bugs, thrips (exposed) and willow leaf beetles.

DIRECTIONS: Use 1 to 2 tablespoons per gallon – Wet foliage to run off Use 5 to 7 oz. to 10 gallons of water.

Lawn Insects – To control ants, chinch bugs, earwigs, armyworms, fleas, ticks, leafhoppers, sod webworms. Use 2 tablespoons per one gallon of water to treat 32 sq. ft. Use one pint to 20 gallons of water to treat 500 sq. ft.

SEVIN® Insect Spray does not control spider mites. Where infestations are encountered, use Sunniland Kelthane. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Injury to tender foliage may occur if applications are made when plants are wet or in the presence of high humidity of several days duration following treatment.

CAUTION: Harmful if swallowed or inhaled. Do not take internally. Avoid breathing spray mist. Avoid contact with skin. Avoid contamination of feed or foodstuffs. Wash thoroughly after handling. Change clothing if contaminated. Protect from temperatures below 32 degrees F. Keep cover on container to avoid drying. After using, rinse empty container thoroughly with water and discard it.

WARRANTY: Conditions of use are beyond Seller's control. Seller's liability is limited to products replacement or purchase price refund. Seller makes no warranty, express or implied, concerning products use beyond claims on label

® Trademark of Union Carbide Corp.
U.S. Pat. No. 2-903,478



NET CONTENTS – ONE QUART

MANUFACTURED BY
CHASE & COMPANY
SANFORD, FLORIDA 32771 Product No. 2314

ACCEPTED