

1/2 to
1/24
1/1

JUL 19 1990

Mr. H.R. McLane
Sunniland Corporation
P.O. Box 1697
Sanford, FL 32771

Dear Mr. McLane:

Subject: Revised Labeling and Confidential Statement of Formula
Sunniland Sevin Insect Spray
EPA Registration No. 9404-9
Your Submission Dated September 1, 1990

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling.

1. Change the percentage for the active ingredient to 21.3%, and inerts to 78.7%.
2. Add the following to the Container Disposal statement:

Do not reuse empty container.

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Dennis H. Edwards, Jr.
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

51676:1:WP5:0:Kumar:C:Disk:KENC:07/17/90:DDIVOTER:DD
CONCURRENCES

SYMBOL	DESCRIPTION	DATE	INITIALS	REMARKS	DATE	INITIALS	REMARKS

ACCEPTED

JUL 19 1990

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 8404-1

(Vegetables Continued)

OKRA: Use 8 tsp./gal. to control Corn Earworm and Stink Bugs. HEAD LETTUCE, CARROTS, CHINESE CABBAGE, COLLARDS, GARDEN BEETS, KALE, MUSTARD GREENS, PARSLEY, RADISHES, RUTABAGAS, SPINACH, SWISS CHARD, AND TURNIPS. Use the low rate (4 tsp./3 gal.) for control of Flea Beetles, Harlequin Bugs, and Leafhoppers. Use higher rate for control of Armyworms, Corn Earworms, Spittlebugs, Tarnished Plant Bugs and Stink Bugs.

Observe the Following Pre-Harvest Spray Interval for Each Vegetable Do not apply within 14 days of harvest on Collards, Chinese Cabbage, Beet Greens(Tops), Kale, Mustard Greens, Parsley, Spinach, Swiss Chard and Turnip tops. Do not treat within 3 days of harvest on Beets, Radishes, Rutabagas, Head Lettuce or Turnips (if tops are not to be used for food). May be applied day of harvest on Beans, Cucumbers, Melons, Pumpkins, Squash, Potato, Tomato, Eggplant, Pepper, Okra, Carrots. Remove viable residues by wiping, washing or other means.

ORNAMENTAL FLOWERS, SHRUBS AND TREES

SUNNILAND SEVIN INSECT SPRAY can be used on a wide range of ornamental plants, flowers, shrubs and small trees, such as Roses, Carnations, Chrysanthemum, Lilac, Arborvitae, Juniper, Hydrangeas, Azaleas, Elm, Maple, Oak, Dogwood, Birch and *Pines. Do not apply to Boston Ivy, Virginia Creeper or Maiden Hair Fern. *It may injure Virginia and Sand Pines(treat small branch first; observe for damage). Mix at the rate of 4 teaspoonfuls per gallon of water and apply to upper and lower leaf surfaces of all foliage for complete coverage to control such pest as: Apple Aphid, Bagworm, Birch Leaf Miner*, Blister Beetles, Boxwood Leaf Miner*, Elm Leaf Aphid, Elm Leaf Beetle, Flea Beetles, Gypsy Moth, Japanese Beetles, June Beetles, Puss Caterpillar, Pyralids, Rose Aphid, Rose Slug, Scale Insects, Tent Caterpillar, Thorn Bugs, Thrips (exposed), and Willow Leaf Beetle. *(Exposed Adults)

LAWNS AND RECREATIONAL TURF

For control of Ants, Earwigs, Fleas, Ticks, Leafhoppers and Sod Webworms use 8 teaspoonfuls in 3 gallon of water to treat 100 Sq.Ft. of lawn, recreational area, roadway or path border (1 Pint will treat 1,200 Sq.Ft.). For best results mow lawn and apply after watering or rain. Insure good penetration of turf. Garden Hose or pressure applicators may be used. Repeat as necessary. For control of Chinch Bugs use 12 teaspoonfuls per 3 gallons of water to cover 100 Sq.Ft. (1 pint treats 800 Sq.Ft.) Treat the entire area to prevent Chinch Bug migration.

SEVIN INSECT SPRAY does not control spider mites. Where "mite" infestations are encountered, use a suitable SUNNILAND product approved for Spider Mite Control. NOTE: Tender "new" foliage may be injured if applications are made when plants are wet or in the presence of high humidity of several days duration following treatment. NOTICE: Conditions of use are beyond seller's control. Seller's liability is limited to products replacement or purchase price refund. Seller makes no warranty, express or implied concerning products use beyond claims on label. 09-01-89



Sevin INSECT SPRAY

Contains Sevin Brand Carbaryl Insecticide

INSECTICIDE FOR LAWNS, VEGETABLES AND ORNAMENTAL PLANTS

ACTIVE INGREDIENT:	
Carbaryl: 1-Naphthyl N-methylcarbamate	22.5%
INERT INGREDIENTS:	77.5%
	Total 100.0%

SEVIN is a Registered Trademark of Rhone-Poulenc for Carbaryl Insecticides. Contains 2 Pounds Carbaryl Insecticide / Gallon

KEEP OUT OF REACH OF CHILDREN CAUTION

"PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente"

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician of Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. IF IN EYES: Flush eyes with plenty of water. Get medical attention. IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Get medical attention.

SEE NOTE TO PHYSICIAN ON SIDE PANEL

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 8404-1

Net Contents 1 Quart (0.946 L)

EPA Est. No. 8404-FL-1

SUNNILAND CORPORATION

P.O. BOX 1697 SANFORD, FL 32771