

GENERAL DIRECTIONS FOR USE ON VEGETABLE INSECTS
 Apply dust thoroughly to infested plants at first sign of damage and repeat at 7 day intervals as necessary or for good insect control. Use 1 to 2 ounces per 50 feet of row on vegetables.

BEANS: (green beans, lima beans, snap beans, cowpeas, black-eyed) To control Mexican bean beetle, bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhopper, velvetbean caterpillar, western bean cutworm, armyworm, corn earworm, stink bugs and tarnished plant bugs.

CUCUMBER, MELONS, PUMPKIN, SQUASH: To control pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers and squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on watermelons in Florida.

POTATO, TOMATO, EGGPLANT, PEPPER: To control Colorado potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworms, lace bugs, tomato fruit worm, tomato hornworms, tarnished plant bugs, stink bugs.

OKRA: To control corn earworms and stink bugs.


HEAD LETTUCE, CARROTS, CHINESE CABBAGE, COLLARDS, GARDEN BEETS, KALE, MUSTARD GREENS, PARSLEY, RADISHES, RUTABAGAS, SPINACH, SWISS CHARD AND TURNIPS: To control flea beetles, harlequin bugs, leafhoppers, armyworms, cabbage caterpillars, corn earworms, spittlebugs, tarnished plant bugs and stink bugs. To control cabbage loopers, use on a 5 to 7-day schedule.

Do not apply to Collards, kale, leaf lettuce, mustard greens, parsley spinach, swiss chard and turnip tops within 14 day period of harvest.

ORNAMENTAL INSECT CONTROL

Sunniland SEVIN DUST can be safely used on roses, carnations, gladioli, chrysanthemums, lilacs, arbor vitae, junipers, hydrangeas, azaleas, elms, maples, oaks, dogwoods, birches and pines. Do not use on Boston ivy.

Apply to leaf surface of all foliage for complete coverage to control: apple aphids, bagworms, birch leaf miners, blister beetles, boxwood leaf miners, elm leaf aphids, elm leaf beetles, flea beetles, gypsy moths, Japanese beetles, June bugs, puss caterpillars, psyllids, rose aphids, rose-slugs, scale insects, tent caterpillars, thorn bugs, thrips (exposed) and willow leaf beetles.



**SEVIN[®] DUST
 FOR GARDEN AND PETS**

Caution Keep Out of Reach of Children
 SEE BACK PANEL FOR ADDITIONAL CAUTIONS

DOGS AND CATS

To control Brown Dog Ticks & Fleas apply liberally to animal and rub thoroughly into skin. Apply to feet, between toes and on legs. Dust sleeping quarters and all areas where ticks and fleas occur. Repeat weekly if needed. Do not treat kittens or puppies less than 4 weeks old.

CAUTION

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing spray mist. Avoid contact with skin. Avoid contamination of feed or foodstuffs. Wash thoroughly after handling. Change clothing if contaminated. Do not re-use container. Destroy when empty.

WARHANTY:

Conditions of use are beyond Seller's control. Seller's liability is limited to products replacement or purchase price refund. Seller makes no warranty, express or implied, concerning products use beyond claims on label.

® Trademark of Union Carbide Corp.
 U. S. Pat. No. 2-903,478

| | |
|--|---------------|
| ACTIVE INGREDIENTS | |
| Carbaryl (1-naphthyl N-methylcarbamate)..... | 5.0% |
| INERT INGREDIENTS | 95.0% |
| TOTAL | 100.0% |

NET WEIGHT - 1 POUND

EPA Registration Number 9404-10



Manufactured by
CHASE & COMPANY
 Sanford, Florida 32771

Product No. 2296

CCP-3M-G1209-9-71

9404-10
 ACCEPTED
 OCT 6 1971
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 ED UNDER NO. SUBJECT
 TO ATTACHED COMMENTS.

9404-10
 10-6-1971

1/1