

945-1-10

GENERAL DIRECTIONS FOR USE ON VEGETABLE INSECTS

Apply dust thoroughly to infested plants at first sign of damage and repeat at 7 day intervals as necessary or for good insect control. Dust the plants on bottom and top of leaf surface thoroughly using 1.2 ounces per 50 ft. of row on vegetables.

BEANS (green beans, lima beans, snap beans, cowpeas, black eyed) To control Mexican bean beetle, bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhopper, velvetbean caterpillar, western bean cutworm, armyworm, corn earworm, stink bugs and tarnished plant bugs.

CUCUMBER, MELONS, PUMPKIN, SQUASH: To control pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers and squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on watermelons in Florida.

POTATO, TOMATO, EGGPLANT, PEPPER: To control Colorado potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworms, lace bugs, tomato fruit worm, tomato hornworms, tarnished plant bugs, stink bugs.

OKRA: To control corn earworms and stink bugs.

HEAD LETTUCE, CARROTS, COLLARDS, GARDEN BEETS, KALE, MUSTARD GREENS, PARSLEY, RADISHES, RUTABAGAS, SPINACH, SWISS CHARD AND TURNIPS: To control flea beetles, harlequin bugs, leafhoppers, armyworms, corn earworms, spittlebugs, tarnished plant bugs and stink bugs.

Do not apply to collards, kale, mustard greens, parsley, spinach, swiss chard and turnip tops within 14 day period of harvest.

Do not apply to head lettuce, radishes and rutabagas within 3 days of harvest. Do not apply to garden beets or turnips within 3 days of harvest. Do not apply within 14 days of harvest if tops are to be used as food or feed.

E.P.A. REG. NO. 9404-10-539

E.P.A. EST. NO. 9404-FL-1

CCP-2M-B343-2-75

ORNAMENTAL INSECT CONTROL

Use Sears Garden & Pet Dust at the rate of 1.2 ounces per 50 sq. ft. Apply evenly to both upper and lower leaf surfaces. Repeat when necessary at 7 day intervals for proper insect control.

Apply to leaf surface of all foliage for complete coverage to control: apple aphids, bagworms, birch leaf miners, blister beetles, boxwood leaf miners, elm leaf aphids, elm leaf beetles, flea beetles, gypsy moths, Japanese beetles, June bugs, puss caterpillars, psyllids, rose aphids, rose slugs, scale insects, tent caterpillars, thorn bugs, thrips (exposed) and willow leaf beetles.

DOGS AND CATS

To control Brown Dog Ticks & Fleas apply liberally to animal and rub thoroughly into skin. Apply to feet, between toes and on legs. Dust sleeping quarters and all areas where ticks and fleas occur. Repeat weekly if needed. Do not treat kittens or puppies less than 4 weeks old.

CAUTION

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust drift. Avoid contact with skin. Avoid contamination of feed or foodstuffs. Wash thoroughly after handling. Change clothing if contaminated. Do not re-use container. Destroy when empty. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

WARRANTY

Conditions of use are beyond Seller's control. Seller's liability is limited to products replacement or purchase price refund. Seller makes no warranty, express or implied concerning products use beyond claims on label.

- Trademark of Union Carbide Corp.
U.S. Pat. No. 2,903,478

ACTIVE INGREDIENTS

Carbaryl (1-naphthyl N-methylcarbamate) 5.0%

INERT INGREDIENTS 95.0%

TOTAL 100.0%

GA

A SCIENTI
USE ON D

CAUTI
See L

ITEM NO. 71

ORNAMENTAL INSECT CONTROL

Use Sears Garden & Pet Dust at the rate of 1/2 ounces per 50 sq. ft. Apply evenly to both upper and lower leaf surfaces. Repeat when necessary at 7 day intervals for proper insect control.

Apply to leaf surface of all foliage for complete coverage to control apple aphids, bagworms, birch leaf miners, blister beetles, boxwood leaf miners, elm leaf aphids, elm leaf beetles, flea beetles, gypsy moths, Japanese beetles, June bugs, puss caterpillars, psyllids, rose aphids, rose slugs, scale insects, tent caterpillars, thorn bugs, thrips (exposed) and willow leaf beetles.

DOGS AND CATS

To control Brown Dog Ticks & Fleas apply liberally to animal and rub thoroughly into skin. Apply to feet, between toes and on legs. Dust sleeping quarters and all areas where ticks and fleas occur. Repeat weekly if needed. Do not treat kittens or puppies less than 4 weeks old.

CAUTION

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust drift. Avoid contact with skin. Avoid contamination of feed or foodstuffs. Wash thoroughly after handling. Change clothing if contaminated. Do not re-use container. Destroy when empty. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

WARRANTY:

Conditions of use are beyond Seller's control. Seller's liability is limited to products replacement or purchase price refund. Seller makes no warranty, express or implied, concerning products use beyond claims on label.

® Trademark of Union Carbide Corp.
U.S. Pat. No. 2,903,478

ACTIVE INGREDIENTS

Carbaryl (1-naphthyl N-methylcarbamate) 5.0%

INERT INGREDIENTS 95.0%

TOTAL 100.0%



**GARDEN AND
PET DUST**

5% SEVIN®

**A SCIENTIFICALLY BLENDED PESTICIDE FOR
USE ON DOGS, CATS, ORNAMENTAL SHRUBS
and VEGETABLES**

**CAUTION: Keep out of reach of children.
See Left Panel For Additional Cautions.**

**SEARS, ROEBUCK AND CO.
CHICAGO, ILLINOIS 60684**

ITEM NO. 7111411

NET WEIGHT - 1 POUND